

Product datasheet for **MC226750**

Npm1 (NM_001252260) Mouse Untagged Clone

Product data:

Product Type: Expression Plasmids
Product Name: Npm1 (NM_001252260) Mouse Untagged Clone
Tag: Tag Free
Symbol: Npm1
Synonyms: B23; NO38; Npm
Vector: pCMV6-Entry (PS100001)
E. coli Selection: Kanamycin (25 ug/mL)
Cell Selection: Neomycin
Fully Sequenced ORF: >MC226750 representing NM_001252260
Red=Cloning site **Blue**=ORF **Orange**=Stop codon

TTTTGTAATACGACTCACTATAGGGCGCCGGAATTCGTCGACTGGATCCGGTACCGAGGAGATCTGCC
GCC**GCGATCGCC**

ATGGAAGACTCGATGGATATGGACATGAGTCCTCTTAGGCCTCAGAACTACCTTTTCGTGGATAATGATG
AAAATGAGCACCAGTTGTCATTAAGAACGGTCAGTTTAGGAGCAGGGGCAAAGATGAGTTACACATCGT
AGAGGCAGAAGCAATGAAGCAATGAAGGCAGTCCAATTAAGTAACACTGGCAACTTTGAAAATGTCTGTA
CAACCAACAGTTTCCCTAGGGGGCTTTGAAATTACACCACCTGTGGTCTTACGGTTGAAGTGTGGTTCAG
GGCCTGTGCACATTAGTGGACAGCATCTAGTAGCTGTAGAGGAAGATGCAGAGTCTGAAGATGAAGATGA
GGAGGACGTAAAACCTTAGGCATGTCTGGAAAGCGATCTGCTCCTGGAGGTGGTAAACAGGTTCCACAG
AAAAAAGTAAAACCTTGAAGATGATGAGGACGATGATGAGGACGATGAGGATGATGAGGATGATGATG
ATGATGATTTTGAAGAGGAAACTGAAGAAAAGTCCCAGTGAAGAAATCTGTACGAGATACCCACAGC
CAAAAATGCACAAAAATCAAACAAAAATGGAAAAGACTTAAACCATCAACACCGAGATCAAAGGGTCAA
GAGTCCTTCAAAAACAGGAAAAGACTCCTAAAACACAAAAGGACCTAGTTCTGTAGAAGACATTAAGG
CAAAAATGCAAGCAAGTATAGAAAAGGGGTTCTCTTCCAAAGTGAAGCCAAGTTTCAATTAATTATGT
GAAGAATTGTTCCGGATGACTGACCAGGAGGCTATCAAGATCTCTGGCAGTGGAGGAAATCTCTT**TAA**

ACGCGTACGCGGCCGCTCGAGCAGAAAACATCTCAGAAGAGGATCTGGCAGCAAATGATATCCTGGATT
ACAAGGATGACGACGATAAGGTTTAA

Restriction Sites: SgfI-MluI
ACCN: NM_001252260
Insert Size: 840 bp



[View online »](#)

OTI Disclaimer:	Our molecular clone sequence data has been matched to the reference identifier above as a point of reference. Note that the complete sequence of our molecular clones may differ from the sequence published for this corresponding reference, e.g., by representing an alternative RNA splicing form or single nucleotide polymorphism (SNP).
OTI Annotation:	Clone contains native stop codon, and expresses the complete ORF without any c-terminal tag.
Components:	The ORF clone is ion-exchange column purified and shipped in a 2D barcoded Matrix tube containing 10ug of transfection-ready, dried plasmid DNA (reconstitute with 100 ul of water).
Reconstitution Method:	<ol style="list-style-type: none">1. Centrifuge at 5,000xg for 5min.2. Carefully open the tube and add 100ul of sterile water to dissolve the DNA.3. Close the tube and incubate for 10 minutes at room temperature.4. Briefly vortex the tube and then do a quick spin (less than 5000xg) to concentrate the liquid at the bottom.5. Store the suspended plasmid at -20°C. The DNA is stable for at least one year from date of shipping when stored at -20°C.
RefSeq:	<u>NM_001252260.1, NP_001239189.1</u>
RefSeq Size:	1401 bp
RefSeq ORF:	840 bp
Locus ID:	18148
Cytogenetics:	11 A4
Gene Summary:	<p>Involved in diverse cellular processes such as ribosome biogenesis, centrosome duplication, protein chaperoning, histone assembly, cell proliferation, and regulation of tumor suppressors p53/TP53 and ARF. Binds ribosome presumably to drive ribosome nuclear export. Associated with nucleolar ribonucleoprotein structures and bind single-stranded nucleic acids. Acts as a chaperonin for the core histones H3, H2B and H4. Stimulates APEX1 endonuclease activity on apurinic/aprimidinic (AP) double-stranded DNA but inhibits APEX1 endonuclease activity on AP single-stranded RNA. May exert a control of APEX1 endonuclease activity within nucleoli devoted to repair AP on rDNA and the removal of oxidized rRNA molecules. In concert with BRCA2, regulates centrosome duplication. Regulates centriole duplication: phosphorylation by PLK2 is able to trigger centriole replication. Negatively regulates the activation of EIF2AK2/PKR and suppresses apoptosis through inhibition of EIF2AK2/PKR autophosphorylation. Antagonizes the inhibitory effect of ATF5 on cell proliferation and relieves ATF5-induced G2/M blockade. In complex with MYC enhances the transcription of MYC target genes.[UniProtKB/Swiss-Prot Function]</p> <p>Transcript Variant: This variant (2) lacks an in-frame segment in the 5' coding region, compared to variant 1. The resulting isoform (2) lacks an internal segment, compared to isoform 1.</p>