

Product datasheet for MC226866

Sirpa (NM_001291021) Mouse Untagged Clone

Product data:

Product Type: Expression Plasmids

Product Name: Sirpa (NM_001291021) Mouse Untagged Clone

Tag: Tag Free Symbol: Sirpa

Synonyms: Al835480; Bit; CD172a; P84; Ptpns1; SHP-1; SHPS-1; SIRP

Vector:pCMV6-Entry (PS100001)E. coli Selection:Kanamycin (25 ug/mL)

Cell Selection: Neomycin

Fully Sequenced ORF: >MC226866 representing NM_001291021

Red=Cloning site Blue=ORF Orange=Stop codon

TTTTGTAATACGACTCACTATAGGGCGGCCGGGAATTCGTCGACTGGATCCGGTACCGAGGAGATCTGCC

GCCGCGATCGCC

ACGCGTACGCGGCCGCTCGAGCAGAAACTCATCTCAGAAGAGGATCTGGCAGCAAATGATATCCTGGATT

ACAAGGATGACGACGATAAGGTTTAA

Restriction Sites: Sgfl-Mlul

ACCN: NM_001291021

Insert Size: 888 bp



OriGene Technologies, Inc. 9620 Medical Center Drive, Ste 200

CN: techsupport@origene.cn

Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com

Sirpa (NM_001291021) Mouse Untagged Clone - MC226866

OTI Disclaimer: Our molecular clone sequence data has been matched to the reference identifier above as a

point of reference. Note that the complete sequence of our molecular clones may differ from the sequence published for this corresponding reference, e.g., by representing an alternative

RNA splicing form or single nucleotide polymorphism (SNP).

OTI Annotation: Clone contains native stop codon, and expresses the complete ORF without any c-terminal

tag.

Components: The ORF clone is ion-exchange column purified and shipped in a 2D barcoded Matrix tube

containing 10ug of transfection-ready, dried plasmid DNA (reconstitute with 100 ul of water).

Reconstitution Method: 1. Centrifuge at 5,000xg for 5min.

2. Carefully open the tube and add 100ul of sterile water to dissolve the DNA.

3. Close the tube and incubate for 10 minutes at room temperature.

4. Briefly vortex the tube and then do a quick spin (less than 5000xg) to concentrate the liquid

at the bottom.

5. Store the suspended plasmid at -20°C. The DNA is stable for at least one year from date of

shipping when stored at -20°C.

RefSeq: <u>NM 001291021.1</u>, <u>NP 001277950.1</u>

2 63.19 cM

 RefSeq Size:
 3389 bp

 RefSeq ORF:
 888 bp

 Locus ID:
 19261

 UniProt ID:
 P97797

Cytogenetics:

Gene Summary: Immunoglobulin-like cell surface receptor for CD47. Acts as docking protein and induces

translocation of PTPN6, PTPN11 and other binding partners from the cytosol to the plasma membrane. Supports adhesion of cerebellar neurons, neurite outgrowth and glial cell attachment. May play a key role in intracellular signaling during synaptogenesis and in synaptic function. Involved in the negative regulation of receptor tyrosine kinase-coupled cellular responses induced by cell adhesion, growth factors or insulin. Mediates negative regulation of phagocytosis, mast cell activation and dendritic cell activation. CD47 binding prevents maturation of immature dendritic cells and inhibits cytokine production by mature

dendritic cells (By similarity).[UniProtKB/Swiss-Prot Function]

Transcript Variant: This variant (6) lacks two alternate in-frame exons in the central coding region, compared to variant 4, resulting in an isoform (4) that is shorter than isoform 3.