



<b>OTI Disclaimer:</b>	Our molecular clone sequence data has been matched to the reference identifier above as a point of reference. Note that the complete sequence of our molecular clones may differ from the sequence published for this corresponding reference, e.g., by representing an alternative RNA splicing form or single nucleotide polymorphism (SNP).
<b>OTI Annotation:</b>	Clone contains native stop codon, and expresses the complete ORF without any c-terminal tag.
<b>Components:</b>	The ORF clone is ion-exchange column purified and shipped in a 2D barcoded Matrix tube containing 10ug of transfection-ready, dried plasmid DNA (reconstitute with 100 ul of water).
<b>Reconstitution Method:</b>	<ol style="list-style-type: none"><li>1. Centrifuge at 5,000xg for 5min.</li><li>2. Carefully open the tube and add 100ul of sterile water to dissolve the DNA.</li><li>3. Close the tube and incubate for 10 minutes at room temperature.</li><li>4. Briefly vortex the tube and then do a quick spin (less than 5000xg) to concentrate the liquid at the bottom.</li><li>5. Store the suspended plasmid at -20°C. The DNA is stable for at least one year from date of shipping when stored at -20°C.</li></ol>
<b>RefSeq:</b>	<a href="#">NM_001301713.1</a> , <a href="#">NP_001288642.1</a>
<b>RefSeq Size:</b>	1925 bp
<b>RefSeq ORF:</b>	948 bp
<b>Locus ID:</b>	12775
<b>UniProt ID:</b>	<a href="#">P47774</a>
<b>Cytogenetics:</b>	11 D
<b>Gene Summary:</b>	Receptor for the MIP-3-beta chemokine.[UniProtKB/Swiss-Prot Function] Transcript Variant: This variant (2) lacks an alternate exon, and it thus differs in its 5' UTR and initiates translation at a downstream in-frame start codon, compared to variant 1. The encoded isoform (b) is shorter at the N-terminus, compared to isoform a.