

Product datasheet for MC227077

Calcoco2 (NM 001271018) Mouse Untagged Clone

Product data:

Product Type: Expression Plasmids

Product Name: Calcoco2 (NM_001271018) Mouse Untagged Clone

Tag: Tag Free
Symbol: Calcoco2

Synonyms: 2410154J16Rik; C77254; Ndp52; Ndp52l1

Vector:pCMV6-Entry (PS100001)E. coli Selection:Kanamycin (25 ug/mL)

Cell Selection: Neomycin

Fully Sequenced ORF: >MC227077 representing NM_001271018

Red=Cloning site Blue=ORF Orange=Stop codon

TTTTGTAATACGACTCACTATAGGGCGGCCGGGAATTCGTCGACTGGATCCGGTACCGAGGAGATCTGCC

GCCGCGATCGCC

ACGCGTACGCGGCCGCTCGAGCAGAAACTCATCTCAGAAGAGGATCTGGCAGCAAATGATATCCTGGATT

ACAAGGATGACGACGATAAGGTTTAA

Restriction Sites: Sgfl-Mlul

ACCN: NM 001271018



OriGene Technologies, Inc. 9620 Medical Center Drive, Ste 200

CN: techsupport@origene.cn

Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com

Calcoco2 (NM_001271018) Mouse Untagged Clone - MC227077

Insert Size: 996 bp

OTI Disclaimer: Our molecular clone sequence data has been matched to the reference identifier above as a

point of reference. Note that the complete sequence of our molecular clones may differ from the sequence published for this corresponding reference, e.g., by representing an alternative

RNA splicing form or single nucleotide polymorphism (SNP).

OTI Annotation: Clone contains native stop codon, and expresses the complete ORF without any c-terminal

tag.

Components: The ORF clone is ion-exchange column purified and shipped in a 2D barcoded Matrix tube

containing 10ug of transfection-ready, dried plasmid DNA (reconstitute with 100 ul of water).

Reconstitution Method: 1. Centrifuge at 5,000xg for 5min.

2. Carefully open the tube and add 100ul of sterile water to dissolve the DNA.

3. Close the tube and incubate for 10 minutes at room temperature.

4. Briefly vortex the tube and then do a quick spin (less than 5000xg) to concentrate the liquid

at the bottom.

5. Store the suspended plasmid at -20°C. The DNA is stable for at least one year from date of

shipping when stored at -20°C.

RefSeq: <u>NM 001271018.1</u>, <u>NP 001257947.1</u>

 RefSeq Size:
 2029 bp

 RefSeq ORF:
 996 bp

 Locus ID:
 76815

 UniProt ID:
 A2A6M5

Cytogenetics: 11 59.4 cM

Gene Summary: Xenophagy-specific receptor required for autophagy-mediated intracellular bacteria

degradation (By similarity). Acts as an effector protein of galectin-sensed membrane damage that restricts the proliferation of infecting pathogens upon entry into the cytosol by targeting

LGALS8-associated bacteria for autophagy (By similarity). Initially orchestrates bacteria

targeting to autophagosomes and subsequently ensures pathogen degradation by regulating

pathogen-containing autophagosome maturation (By similarity). Bacteria targeting to autophagosomes relies on its interaction with MAP1LC3A, MAP1LC3B and/or GABARAPL2, whereas regulation of pathogen-containing autophagosome maturation requires the interaction with MAP3LC3C (By similarity). May play a role in ruffle formation and actin cytoskeleton organization and seems to negatively regulate constitutive secretion (By

similarity).[UniProtKB/Swiss-Prot Function]