

Product datasheet for **MC227613**

Mknk1 (NM_001285487) Mouse Untagged Clone

Product data:

Product Type:	Expression Plasmids
Product Name:	Mknk1 (NM_001285487) Mouse Untagged Clone
Tag:	Tag Free
Symbol:	Mknk1
Synonyms:	2410048M24Rik; Mnk; Mnk1
Mammalian Cell Selection:	Neomycin
Vector:	pCMV6-Entry (PS100001)
E. coli Selection:	Kanamycin (25 ug/mL)
Restriction Sites:	Sgfl-Mlul
ACCN:	NM_001285487
Insert Size:	1266 bp
OTI Disclaimer:	Our molecular clone sequence data has been matched to the reference identifier above as a point of reference. Note that the complete sequence of our molecular clones may differ from the sequence published for this corresponding reference, e.g., by representing an alternative RNA splicing form or single nucleotide polymorphism (SNP).
OTI Annotation:	Clone contains native stop codon, and expresses the complete ORF without any c-terminal tag.
Components:	The ORF clone is ion-exchange column purified and shipped in a 2D barcoded Matrix tube containing 10ug of transfection-ready, dried plasmid DNA (reconstitute with 100 ul of water).
Reconstitution Method:	<ol style="list-style-type: none">1. Centrifuge at 5,000xg for 5min.2. Carefully open the tube and add 100ul of sterile water to dissolve the DNA.3. Close the tube and incubate for 10 minutes at room temperature.4. Briefly vortex the tube and then do a quick spin (less than 5000xg) to concentrate the liquid at the bottom.5. Store the suspended plasmid at -20°C. The DNA is stable for at least one year from date of shipping when stored at -20°C.
RefSeq:	<u>NM_001285487.1</u> , <u>NP_001272416.1</u>



[View online »](#)

RefSeq Size: 2704 bp

RefSeq ORF: 1266 bp

Locus ID: 17346

Cytogenetics: 4 D1

Gene Summary: This gene encodes a serine-threonine protein kinase that is activated by extracellular signal-regulated kinase or p38 mitogen-activated protein kinases, and it may function in cytokine and environmental stress responses. This kinase is required for phosphorylation of eukaryotic translation initiation factor 4E but it is not required for cell growth during development. Alternative splicing of this gene results in multiple transcript variants. [provided by RefSeq, Oct 2013]

Transcript Variant: This variant (1) represents the longest transcript and encodes the longer isoform (a). Sequence Note: This RefSeq record was created from transcript and genomic sequence data to make the sequence consistent with the reference genome assembly. The genomic coordinates used for the transcript record were based on transcript alignments.