

Product datasheet for **MC227729**

Nr1h2 (NM_001285519) Mouse Untagged Clone

Product data:

Product Type:	Expression Plasmids
Product Name:	Nr1h2 (NM_001285519) Mouse Untagged Clone
Tag:	Tag Free
Symbol:	Nr1h2
Synonyms:	A1194859; LXR; LXRβ; LXRbeta; LXRBSV; NER1; OR-1; RIP15; Unr; Unr2; UR
Vector:	pCMV6-Entry (PS100001)
E. coli Selection:	Kanamycin (25 ug/mL)
Cell Selection:	Neomycin



[View online »](#)

Fully Sequenced ORF: >MC227729 representing NM_001285519
Red=Cloning site Blue=ORF Orange=Stop codon

TTTTGTAATACGACTCACTATAGGGCGGCCGGAATTCGTCGACTGGATCCGGTACCGAGGAGATCTGCC
GCC**CGATCGCC**

ATGTCCTCCCCACAAGTTCTCTGGACACTCCCGTGCCTGGGAATGGTTCTCCTCAGCCCAGTACCTCCG
CCACGTACCCACTATTAAGGAAGAGGGGCAGGAGACTGATCCTCCTCCAGGCTCTGAAGGGTCCAGCTC
TGCTACATCGTGAGGCCAGAGGATGAGCCTGAGCGCAAGCGGAAGAAGGGGCCGCCCCGAAGATGCTG
GGCCATGAGCTGTGCCGCGTGTGCGGAGACAAGGCTTCGGGCTTCCACTACAACGTGCTCAGCTGTGAAG
GCTGCAAAGGCTTCTCCGGCGCAGTGTGGTCCACGGTGGGGCCGGGCGCTATGCCTGTCGGGGCAGCGG
AACCTGCCAGATGGATGCCTTCATGCGGCGCAAGTGCCAGCTCTGCCGGTGCAGCAAGTGAAGGAGGCT
GGCATGCGGGAGCAGTGCCTGCTCTGAGGAGCAGATTCGGAAGAAAAGGATTCAGAAGCAGCAACAGC
AGCAGCCACCACCCCATCTGAGCCAGCAGCCAGCAGCTCAGGCCGGCCAGCGGCTCCCCTGGCACTTC
GGAAGCAAGCAGCCAGGGCTCCGGGAAGGAGAGGGCATCCAGCTGACCGCGGCTCAGGAGCTGATGATC
CAGCAGTTAGTTGCCGCGCAGCTGCAAGCAAAACGATCTTTCTCCGACCAGCCAAAAGTCACGCCCT
GGCCCTGGGTGCAGACCCTCAGTCCCAGATGCCCGTCAGCAACGCTTTGCCCACTTACCAGGAGTAGC
CATCATCTCGGTCCAGGAGATTGTGGACTTTGCCAAGCAGGTGCCAGGGTTCTTGCAAGTGGGCCGGGAG
GACCAGATCGCCCTCCTGAAGGCGTCCACCATTGAGATCATGTTGCTAGAAAACAGCCAGACGCTACAACC
ACGAGACAGAATGCATCACGTTCTGAAGGACTTACCTACAGCAAGGACGACTTCCACCGTGCAGGCTT
GCAGGTGGAATTCATCAATCCCATCTTCGAGTTCTCGCGGGCCATGCGCGGCTGGGCTGGACGATGCA
GAGTATGCCTTGCTTATCGCCATCAACATCTTCTCAGCCGATCGGCCTAATGTGCAGGAGCCAGCCGTG
TGGAGCCCTGCAGCAGCCCTACGTGGAGGCGCTCCTCTCCTACACGAGGATCAAGCGCCACAGGACCA
GCTCCGCTTCCCACGCATGCTCATGAAGCTGGTGAAGCTGAGCCTGCGCACCCCTCAGCTCCGTGCACTCGGAGCAG
GTCTTTGCATTGCGACTCCAGGACAAGAAGCTGCCGCCCTTGCTGTCGAGATCTGGGATGTGCACGAGT
AG

ACGCGTACGCGGCCGCTCGAGCAGAACTCATCTCAGAAGAGGATCTGGCAGCAAATGATATCCTGGATT
ACAAGGATGACGACGATAAGGTTTAA

Restriction Sites: SgfI-MluI

ACCN: NM_001285519

Insert Size: 1332 bp

OTI Disclaimer: Our molecular clone sequence data has been matched to the reference identifier above as a point of reference. Note that the complete sequence of our molecular clones may differ from the sequence published for this corresponding reference, e.g., by representing an alternative RNA splicing form or single nucleotide polymorphism (SNP).

OTI Annotation: Clone contains native stop codon, and expresses the complete ORF without any c-terminal tag.

Components: The ORF clone is ion-exchange column purified and shipped in a 2D barcoded Matrix tube containing 10ug of transfection-ready, dried plasmid DNA (reconstitute with 100 ul of water).

Reconstitution Method:	<ol style="list-style-type: none">1. Centrifuge at 5,000xg for 5min.2. Carefully open the tube and add 100ul of sterile water to dissolve the DNA.3. Close the tube and incubate for 10 minutes at room temperature.4. Briefly vortex the tube and then do a quick spin (less than 5000xg) to concentrate the liquid at the bottom.5. Store the suspended plasmid at -20°C. The DNA is stable for at least one year from date of shipping when stored at -20°C.
RefSeq:	<u>NM_001285519.1</u> , <u>NP_001272448.1</u>
RefSeq Size:	2013 bp
RefSeq ORF:	1332 bp
Locus ID:	22260
UniProt ID:	<u>Q60644</u>
Cytogenetics:	7 B3
Gene Summary:	<p>Nuclear receptor that exhibits a ligand-dependent transcriptional activation activity (PubMed:18055760, PubMed:19520913, PubMed:20427281). Binds preferentially to double-stranded oligonucleotide direct repeats having the consensus half-site sequence 5'-AGGTCA-3' and 4-nt spacing (DR-4) (PubMed:18055760, PubMed:19520913, PubMed:20427281). Regulates cholesterol uptake through MYLIP-dependent ubiquitination of LDLR, VLDLR and LRP8; DLDLR and LRP8 (PubMed:18055760, PubMed:19520913, PubMed:20427281). Interplays functionally with RORA for the regulation of genes involved in liver metabolism (PubMed:18055760, PubMed:19520913, PubMed:20427281). Plays an anti-inflammatory role during the hepatic acute phase response by acting as a corepressor: inhibits the hepatic acute phase response by preventing dissociation of the N-Cor corepressor complex (By similarity).[UniProtKB/Swiss-Prot Function]</p> <p>Transcript Variant: This variant (4) differs in the 5' UTR and uses an alternate in-frame splice junction at the 5' end of an exon compared to variant 1. The resulting isoform (b) has the same N- and C-termini but is shorter compared to isoform a. Variants 3 and 4 both encode the same isoform (b).</p>