

## Product datasheet for **MC228247**

### **Nf2 (NM\_001252253) Mouse Untagged Clone**

#### **Product data:**

Product Type:	Expression Plasmids
Product Name:	Nf2 (NM_001252253) Mouse Untagged Clone
Tag:	Tag Free
Symbol:	Nf2
Synonyms:	merlin
Vector:	pCMV6-Entry (PS100001)
E. coli Selection:	Kanamycin (25 ug/mL)
Cell Selection:	Neomycin



[View online »](#)

**Fully Sequenced ORF:** >MC228247 representing NM\_001252253  
 Red=Cloning site Blue=ORF Orange=Stop codon

TTTTGTAATACGACTCACTATAGGGCGGCCGGAATTCGTCGACTGGATCCGGTACCGAGGAGATCTGCC  
 GCC**CGGATCGCC**

ATGAAATGGAAGGGAAGGACCTGTTTGTATTTGGTGTGCCGACTGGGGCTTCGGAAACCTGGTTCT  
 TTGGACTGCAGTATACAATCAAGGACACGGTGGCTGGCTCAAAATGGACAAGAAGGTGTTGGATCATGA  
 TGTTTCGAAGGAAGAACCAGTTACCTTTCACTTCTGGCCAAATTTTATCCTGAAATGCTGAGGAGGAG  
 CTAGTTCAAGAGATCACGCAACCTTATTTTTCTTACAGGTGAAGAAGCAGATTTTGGATGAAAAGGTCT  
 ACTGCCCTCCCGAGGCGTCCGTGCTCTTGGCGTCATATGCTGTCCAGGCCAAGTATGGCGACTATGACCC  
 CTCTGTGCACAAGCGGGATTTTAGCCCAAGAGGAATTGCTCCCGAAAAGGGTGATAAATCTCTATCAG  
 ATGACTCCGAAATGTGGGAGGAGAGAATTACGGCTTGGTATGCGGAGCACCGGGGCAGAGCCAGGGATG  
 AAGCTGAAATGGAGTATTTGAAGATAGCTCAGGACCTGGAGATGTATGGTGTGAACACTTTACAATCCG  
 GAATAAAAAGGGCACAGAGTTGCTGCTTGGAGTGGATGCTCTTGGGCTTCATATCTATGACCCTGAGAAC  
 AGGCTGACCCCAAGATCTCCTTCCCATGGAATGAAATCCGAAACATCTCTACAGCGACAAGGAGTTTA  
 CTATTAACCCTGGATAAGAAAATTGATGCTTCAAATTTAACTCCTCAAAGCTTCGTGTTAATAAGCT  
 GATTCTTCAGCTATGTATTGGGAACCATGACCTATTTATGAGGCGACGAAAAGCTGACTCTTTAGAAGTT  
 CAGCAGATGAAAGCCAGGCCAGGGAAGAGAAGGCTAGAAAGCAGATGAAAGGCCAGCGCTGGCTCGAG  
 AGAAGCAGATGCGGGAGGAGGCCGAGCGTACAAGAGATGAGTTAGAGAGGAGGCTCCTGCAGATGAAAGA  
 AGAAGCAACGATGGCCAATGAAGCTCTGATGCGCTCTGAGGAGACAGCTGATCTGTTGGCTGAAAAGGCT  
 CAGATCACAGAGGAGGAGGCCAAGCTTTTGGCACAAAAGGCTGCAGAGGCTGAGCAAGAGATGCAGCGAA  
 TCAAGGCCACGGCATTTCGGACAGAGGAGGAGAAGCGCCTGATGGAGCAGAAGGTCTGGAGGCTGAAGT  
 GCTGGCATTGAAGATGGCTGAGGAGTCAAGAGAGGAGGCCAAGGAGGCTGATCAGTTAAAGCAAGACTTG  
 CAAGAAGCCAGAGAAGCAGAGCGAAGAGCCAAGCAGAAGCTCTTAGAAATCGCCACCAAGCCACCTATC  
 CACCCATGAACCAATTCCACCACCTGCCTCCTGACATACCGAGCTTCGACATTATTGCTGACAGCTT  
 GTCATTTGACTTCAAGGATACGGACATGAAGCGACTTTCCATGGAGATAGAGAAAAGAAAAGTGGAGTAC  
 ATGGAGAAGAGCAAGCACCTGCAGGAGCAGCTCAACGAGCTCAAGACGGAGATCGAGGCCTTGAACTCA  
 AAGAGCGGAGACGGCTTGGACGCTCTACACAGCGAGAGCTCAGACAGAGGCGGCCCCAGCAGCAAGCA  
 TAATACCATTA AAAAGGTACCTGAAAT**GTA**

**ACGCGT**ACGCGGCCGCTCGAGCAGAAACTCATCTCAGAAGAGGATCTGGCAGCAAATGATATCCTGGATT  
 ACAAGGATGACGACGATAAGGTTTAA

- Restriction Sites:** SgfI-MluI
- ACCN:** NM\_001252253
- Insert Size:** 1641 bp
- OTI Disclaimer:** Our molecular clone sequence data has been matched to the reference identifier above as a point of reference. Note that the complete sequence of our molecular clones may differ from the sequence published for this corresponding reference, e.g., by representing an alternative RNA splicing form or single nucleotide polymorphism (SNP).
- OTI Annotation:** Clone contains native stop codon, and expresses the complete ORF without any c-terminal tag.
- Components:** The ORF clone is ion-exchange column purified and shipped in a 2D barcoded Matrix tube containing 10ug of transfection-ready, dried plasmid DNA (reconstitute with 100 ul of water).

<b>Reconstitution Method:</b>	<ol style="list-style-type: none"><li>1. Centrifuge at 5,000xg for 5min.</li><li>2. Carefully open the tube and add 100ul of sterile water to dissolve the DNA.</li><li>3. Close the tube and incubate for 10 minutes at room temperature.</li><li>4. Briefly vortex the tube and then do a quick spin (less than 5000xg) to concentrate the liquid at the bottom.</li><li>5. Store the suspended plasmid at -20°C. The DNA is stable for at least one year from date of shipping when stored at -20°C.</li></ol>
<b>RefSeq:</b>	<u><a href="#">NM_001252253.1</a></u> , <u><a href="#">NP_001239182.1</a></u>
<b>RefSeq Size:</b>	1846 bp
<b>RefSeq ORF:</b>	1641 bp
<b>Locus ID:</b>	18016
<b>UniProt ID:</b>	<u><a href="#">P46662</a></u>
<b>Cytogenetics:</b>	11 3.0 cM
<b>Gene Summary:</b>	<p>Probable regulator of the Hippo/SWH (Sav/Wts/Hpo) signaling pathway, a signaling pathway that plays a pivotal role in tumor suppression by restricting proliferation and promoting apoptosis. Along with WWC1 can synergistically induce the phosphorylation of LATS1 and LATS2 and can probably function in the regulation of the Hippo/SWH (Sav/Wts/Hpo) signaling pathway. May act as a membrane stabilizing protein. May inhibit PI3 kinase by binding to AGAP2 and impairing its stimulating activity. Suppresses cell proliferation and tumorigenesis by inhibiting the CUL4A-RBX1-DDB1-VprBP/DCAF1 E3 ubiquitin-protein ligase complex (By similarity). Plays a role in lens development and is required for complete fiber cell terminal differentiation, maintenance of cell polarity and separation of the lens vesicle from the corneal epithelium.[UniProtKB/Swiss-Prot Function]</p> <p>Transcript Variant: This variant (5) differs in the 5' UTR, 3' UTR, and coding region, compared to variant 1. The resulting isoform (4) is shorter at the N-terminus and has a shorter and distinct C-terminus, compared to isoform 1.</p>