

Product datasheet for **MC228313**

Eps8 (NM_001271589) Mouse Untagged Clone

Product data:

Product Type:	Expression Plasmids
Product Name:	Eps8 (NM_001271589) Mouse Untagged Clone
Tag:	Tag Free
Symbol:	Eps8
Synonyms:	AW261790
Vector:	pCMV6-Entry (PS100001)
E. coli Selection:	Kanamycin (25 ug/mL)
Cell Selection:	Neomycin



[View online »](#)

Fully Sequenced ORF: >MC228313 representing NM_001271589
 Red=Cloning site Blue=ORF Orange=Stop codon

TTTTGTAATACGACTCACTATAGGGCGGCCGGAATTCGTCGACTGGATCCGGTACCGAGGAGATCTGCC
 GCC**CGGATCGCC**

ATGATGGCAGCCCGGATCGACAGGGATGTGCAAACTCTAAACCATATTTTGGATGACATTGAATTTTTTA
 TCACAAACTCCAAAAGCCGCGAAGCGTTTTCTGAGCTTTCTAAAAGGAAGAAAAGTAAAGAAAAGTAA
 AAGGAAAAGGACCTGGAGAGGGCGTTTTAACACTGAGGGCAAACCCGCCACCTCCTGATGAATTTGTTGAC
 TGTTCAGAAAGTTAAACATGGATTCAACCTTCTGGCCAAGTTGAAGTCCCATATCCAGAACCCGAGTG
 CTTTCCAGATCTGGTTCAATTTTTTTTACTCCACTAAATATGGTGGTCCAGGCAACAGGTGGCCCCGAACT
 GGCCAGTTCGGTACTCAGCCACTGTTGACAAAAGACACAGTTGATTTCTTAAACTACACAGCCACTGCC
 GAGAACCGAAGCTGTGGATGTCCTGAGGAGATAGTTGGTGAAGTGAAGCAGAGTGGCCGAAAGAAC
 AGTTCATCCACCTTACGTCCCGAGGTTCCGCAACGGCTGGGAGCCCCGATGCTGAACTCATGGGCGC
 GCCACAGAGCAAGACATGTATCAACTGGCCGAGTCCGTGGCCAACGCAGAACACCAGCGCAAACAGGAC
 AGCAAGAGGCTGTCCACAGAGCATTCCAATGTGTCCGACTATCCTCCAGCCGACGGATATGCGTACAGTA
 GCAGCATGTACCACAGAGGACCACATGCAGACCACGGGGAGGCTGCCATGCCTTTCAAGTCAACTCTAA
 TCACCAAGTAGATAGGAATTATGACGCAGTCAAAACACAACCCAAAGAAATACGCCAAATCCAAGTACGAC
 TTTGTGGCGAGGAACAGCAGCGAGCTCTCGGTTATGAAAGATGATGTCTTAGAGATACTCGACGATCGAA
 GGCAGTGGTGGAAAGTCCGGAATGCCAGTGGAGACTCTGGGTTTGTGCCAAATAACATTCTGGATATCAT
 GAGAACTCCGGAATCTGGAGTGGGGCGCGCTGACCCCCATACACACATACCATACAGAAACAAAGGACG
 GAATACGGCCTGAGATCAGCTGACACTCCTTCTGCCCATCACCCCTCCAACGCCAGCACCCGTTCCGG
 TCCCTTCCACCTTCTGTACCAGCACCCGTTTCTGTGCCAAGGTCCCAGCCAATGTCACCCGCCAGAA
 CAGCAGCTCCAGTGACAGTGGGGCAGCATTGTGCGGGACAGCCAGAGATACAAACAACCTCCAGTGGAC
 CGAAGGAAGTCCCAGATGGAAGAGGTTCCAGGATGAGCTTCCAGAGGCTGACCATCGGGCGCAGTGCTG
 CACAGAGGAAGTCCACGTGCCACGGCAGAACGTTCCAGTATCAATATCACTTATGACTCCTCACCGGA
 AGAAGTAAAGACTTGGCTGCAGTCAAAGGGATTCAACCCCGTACTGTCAATAGCCTCGGGGTGTTGAAC
 GGAGCACAACTCTTTTCTCAACAAAGACGAACTGAGGTCTGTCTGCCGGAAGGTGCCAGAGTCTTTA
 ACCAAACTACTGTTCCAGAAAGCTGCTTTGGAGGACAGTAAAGGAGTCCGAGTTACAAGAGATCATGCC
 GAGACGGCAGGAGAAGATCAGCGCCGCTGCGAGCGACTCGGGAGTGGAGTCTTTCGATGAAGGGAGCAGC
 CACTGA

ACGCGTACGCGGCCGCTCGAGCAGAAACTCATCTCAGAAGAGGATCTGGCAGCAAATGATATCCTGGATT
 ACAAGGATGACGACGATAAGGTTTAA

- Restriction Sites:** SgfI-MluI
- ACCN:** NM_001271589
- Insert Size:** 1686 bp
- OTI Disclaimer:** Our molecular clone sequence data has been matched to the reference identifier above as a point of reference. Note that the complete sequence of our molecular clones may differ from the sequence published for this corresponding reference, e.g., by representing an alternative RNA splicing form or single nucleotide polymorphism (SNP).
- OTI Annotation:** Clone contains native stop codon, and expresses the complete ORF without any c-terminal tag.
- Components:** The ORF clone is ion-exchange column purified and shipped in a 2D barcoded Matrix tube containing 10ug of transfection-ready, dried plasmid DNA (reconstitute with 100 ul of water).

Reconstitution Method:

1. Centrifuge at 5,000xg for 5min.
2. Carefully open the tube and add 100ul of sterile water to dissolve the DNA.
3. Close the tube and incubate for 10 minutes at room temperature.
4. Briefly vortex the tube and then do a quick spin (less than 5000xg) to concentrate the liquid at the bottom.
5. Store the suspended plasmid at -20°C. The DNA is stable for at least one year from date of shipping when stored at -20°C.

RefSeq: [NM_001271589.1](#), [NP_001258518.1](#)

RefSeq Size: 3562 bp

RefSeq ORF: 1686 bp

Locus ID: 13860

UniProt ID: [Q08509](#)

Cytogenetics: 6 66.78 cM

Gene Summary: Signaling adapter that controls various cellular protrusions by regulating actin cytoskeleton dynamics and architecture. Depending on its association with other signal transducers, can regulate different processes. Together with SOS1 and ABI1, forms a trimeric complex that participates in transduction of signals from Ras to Rac by activating the Rac-specific guanine nucleotide exchange factor (GEF) activity. Acts as a direct regulator of actin dynamics by binding actin filaments and has both barbed-end actin filament capping and actin bundling activities depending on the context. Displays barbed-end actin capping activity when associated with ABI1, thereby regulating actin-based motility process: capping activity is auto-inhibited and inhibition is relieved upon ABI1 interaction. Also shows actin bundling activity when associated with BAIAP2, enhancing BAIAP2-dependent membrane extensions and promoting filopodial protrusions. Involved in the regulation of processes such as axonal filopodia growth, stereocilia length, dendritic cell migration and cancer cell migration and invasion. Acts as a regulator of axonal filopodia formation in neurons: in the absence of neurotrophic factors, negatively regulates axonal filopodia formation via actin-capping activity. In contrast, it is phosphorylated in the presence of BDNF leading to inhibition of its actin-capping activity and stimulation of filopodia formation. Component of a complex with WHRN and MYO15A that localizes at stereocilia tips and is required for elongation of the stereocilia actin core. Indirectly involved in cell cycle progression; its degradation following ubiquitination being required during G2 phase to promote cell shape changes.

[UniProtKB/Swiss-Prot Function]

Transcript Variant: This variant (5) contains a distinct 5' UTR, initiates translation at a downstream start codon and lacks an in-frame portion of the 5' coding region, compared to variant 1. The resulting isoform (3) has a shorter N-terminus, compared to isoform 1.

Sequence Note: This RefSeq record was created from transcript and genomic sequence data to make the sequence consistent with the reference genome assembly. The genomic coordinates used for the transcript record were based on transcript alignments.