

Product datasheet for **MC228387**

Enox2 (NM_001271450) Mouse Untagged Clone

Product data:

Product Type:	Expression Plasmids
Product Name:	Enox2 (NM_001271450) Mouse Untagged Clone
Tag:	Tag Free
Symbol:	Enox2
Synonyms:	APK1; Cova1; tNOX
Vector:	pCMV6-Entry (PS100001)
E. coli Selection:	Kanamycin (25 ug/mL)
Cell Selection:	Neomycin



[View online »](#)

Fully Sequenced ORF: >MC228387 representing NM_001271450
 Red=Cloning site Blue=ORF Orange=Stop codon

TTTTGTAATACGACTCACTATAGGGCGGCCGGAATTCGTCGACTGGATCCGGTACCGAGGAGATCTGCC
 GCC**CGGATCGCC**

ATGACGCTGCCTGTGTCTGATCCAGCTGCATGGGCCACAGCAATGAATAATCTTGAATGGCTCCACTGG
 GAATTGCTGGACAACCAATTTTACCTGACTTCGATCCTGCCCTTGGGATGATGACTGGAATACCACCAAT
 AACTCCCATGATGCCGGTTTGGGCATAGTCCCGCCACCGATTCTCCAGATATGCCGTAGCAAAGGAG
 ATCATACACTGCAAAAGCTGCACGCTCTTCCCTCCCAACCCAAATCTTCCACCACCTGCAACACGAGAAA
 GGCCACCAGGCTGTAAGACAGTGTGGTGGCTGCCTGAAAATGGGACAGAGCAGATCATTGTGGA
 AGTGTGTTGAACAGTGTGGAGAGATTATTGCTATCCGGAAGAGCAAAAAGAACTTCTGTACATTCGTTT
 GCTGAGGAATACATGGTGGACAAAGCCCTATCTGTCTGGTTACCGAATTCGTCTGGGCTAGTACTG
 ACAAGAAGGACACAGGCCGGCTCCATGTTGACTTTGCCAGGCTCGGGATGACTTATATGAGTGGGAGTG
 TAAACAGCGTATGCTAGCCAGAGAGGAGCGGCACCGTAGAAGGATGGAAGAAGAAAGAAATGCGTCCACCA
 TCCCCACCTCCAGTGGTCCACTATTCAGATCATGAATGCAGCATTGTTGCTGAAAAACTCAAAGATGATT
 CCAAATTCCTGGGAAGCTGTGCAGACCTTGTCACTGGATTGAGAGGGGGGAAGTGAACCGCCGAGTGC
 CAACCACTTCTACTCCATGATCCAGTCAGCCAACAGCCATGTCCGCCGCTGGTAAATGAGAAAAGCTACC
 CATGAGAAAAGAGATGGAAGAAGCAAAAGAGAAGTTCAAGCAGGCCCTTTCTGGAATTCTCATTCAATTTG
 AGCAGATAGTAGCTGTGTACCATTCCGCTTCCAAACAGAAGGCATGGGACCACCTCACAAAAGCACAACG
 TAAAAACATCAGTGTGGTGCAACAAGCTGAGGAAATTCGCAACATTCATAATGATGAATTAATGGGA
 ATCAGAAGAGAAGAAGAAATGAAATGTCTGATGATGAAATAGAAGAGACAACAGAAAACAAAAGAAACGG
 AGGAATCAGCCTTAGTGTACAGGCAGAAGCTCTGAAGGAGGAAAATGATAGCCTCCGCTGGCAGCTGGA
 CGCATACCGGAATGAGGTAGAAGTGTCTCAACAGGAACAAGGCAAGCCACAGGGAGGATGACCCAAAC
 AAGGAACAGCAGCTGAACTTCTGCAGCAAGCCCTACAAGGGATGCAACAGCATCTGCTCAAAGTCCAAG
 AGGAATACAAAAGAAAGAAAGCTGAGCTTGACCGAATCAAAGACGACAATTTACAGCAGGAAAAGTGTGG
 TTCTAGACTGTGTGCATCAAGCCAAGAAGGTGAGCAACCTCTTGAGAAGACCGCAGTCAGCAATCTGTGC
 AAATCTGAACGTGAGGCACTGCTAGTGGGGATCATCTCCACATTCCTTCCAGTCCACCCATTTGGAGCCA
 GCATTGAATACATCTGTTCTACTTGAATCGTCTCGATAATAAGGCAAGCTACCAATACCCAGCAAATT
 AACCACTTCTCCACTTACCGATCAGCACCAGTGTGGAATCTCTCATGAGTACTCCAGCATAACC
 TTCAGACAGGAAATGACTGGAGTTGGAGCCAGCCTGGAGAAGAGATGGAATTCGTGGTGGTGGAGGCT
 TGAAGCTGACCTGA

ACGCGTACGCGGCCGCTCGAGCAGAAACTCATCTCAGAAGAGGATCTGGCAGCAAATGATATCCTGGATT
 ACAAGGATGACGACGATAAGGTTTAA

- Restriction Sites:** SgfI-MluI
- ACCN:** NM_001271450
- Insert Size:** 1764 bp
- OTI Disclaimer:** Our molecular clone sequence data has been matched to the reference identifier above as a point of reference. Note that the complete sequence of our molecular clones may differ from the sequence published for this corresponding reference, e.g., by representing an alternative RNA splicing form or single nucleotide polymorphism (SNP).
- OTI Annotation:** Clone contains native stop codon, and expresses the complete ORF without any c-terminal tag.
- Components:** The ORF clone is ion-exchange column purified and shipped in a 2D barcoded Matrix tube containing 10ug of transfection-ready, dried plasmid DNA (reconstitute with 100 ul of water).

Reconstitution Method:

1. Centrifuge at 5,000xg for 5min.
2. Carefully open the tube and add 100ul of sterile water to dissolve the DNA.
3. Close the tube and incubate for 10 minutes at room temperature.
4. Briefly vortex the tube and then do a quick spin (less than 5000xg) to concentrate the liquid at the bottom.
5. Store the suspended plasmid at -20°C. The DNA is stable for at least one year from date of shipping when stored at -20°C.

RefSeq: [NM_001271450.1](#), [NP_001258379.1](#)

RefSeq Size: 3933 bp

RefSeq ORF: 1764 bp

Locus ID: 209224

UniProt ID: [Q8R0Z2](#)

Cytogenetics: X A5

Gene Summary: May be involved in cell growth. Probably acts as a terminal oxidase of plasma electron transport from cytosolic NAD(P)H via hydroquinones to acceptors at the cell surface. Hydroquinone oxidase activity alternates with a protein disulfide-thiol interchange/oxidoreductase activity which may control physical membrane displacements associated with vesicle budding or cell enlargement. The activities oscillate with a period length of 22 minutes and play a role in control of the ultradian cellular biological clock (By similarity).[UniProtKB/Swiss-Prot Function]

Transcript Variant: This variant (5) differs in the 5' UTR and uses an alternate in-frame splice site, compared to variant 1. The encoded isoform (b) is shorter than isoform a. Sequence Note: This RefSeq record was created from transcript and genomic sequence data to make the sequence consistent with the reference genome assembly. The genomic coordinates used for the transcript record were based on transcript alignments.