

Product datasheet for **MC228486**

Itk (NM_001281968) Mouse Untagged Clone

Product data:

Product Type:	Expression Plasmids
Product Name:	Itk (NM_001281968) Mouse Untagged Clone
Tag:	Tag Free
Symbol:	Itk
Synonyms:	Emt; Tcsk; Tsk
Vector:	pCMV6-Entry (PS100001)
E. coli Selection:	Kanamycin (25 ug/mL)
Cell Selection:	Neomycin



[View online »](#)

Fully Sequenced ORF: >MC228486 representing NM_001281968
 Red=Cloning site Blue=ORF Orange=Stop codon

TTTTGTAATACGACTCACTATAGGGCGGCCGGGAATTCGTCGACTGGATCCGGTACCGAGGAGATCTGCC
 GCC**CGCATCGCC**

ATGAACAACCTTCATCCTCCTGGAAGAACAGCTGATCAAGAAGTCCCAACAGAAGAGAAGGACTTCTCCCT
 CGAATTTTAAAGTTCGTTTCTTTGTCTTAACGAAAGCCAGCCTGGCCTACTTTGAGGACCGCCACGGGAA
 GCGCACGTTGAAGGGCTCCATTGAACTCTCCAGAATCAAGTGTGTGGAGATTGTCAAGAGTGACATTAGC
 ATCCCGTGCCACTATAAATACCCCTTTTCAGGTCGTGCATGACAACTATCTCCTGTATGTGTTTGCTCCAG
 ACTGTGAGAGTCGGCAGCGCTGGGTGCTGACCCTTAAAGAAGAAACGAGGAATAACAACAGCCTGGTATC
 CAAGTATCACCCTAATTTCTGGATGGATGGGCGGTGGAGGTGCTGCTCCCAGCTGGAGAAGCCTGCTGTA
 GGCTGTGCTCCCTACGACCCATCCAAGAATGCTTCAAAGAAGCCTCTTCTCCTACTCCTGAAGACAACA
 GCGGTCATTTCAGGAACCTGAAGAAACCTGGTCAATGCCTGTACGACTACCAAACCAACGACCCCTCA
 GGAGCTCGCACTCGGTGTGATGAAGAGTACTACCTGCTGGACAGCTCCGAGATCCACTGGTGGAGGGTT
 CAAGACAAAAATGGGCATGAAGGATATGCACCAAGCAGTTACCTGGTAGAAAAATCTCCAATAACCTTG
 AAACCTATGAGTGGTACAATAAAAGCATCAGCCGCGACAAGCTGAAAACTTCTTTTGGACACAGGTAA
 AGAAGGAGCTTTCATGGTCCGAGATTCAGGACACCCGGGACATACACAGTCTCTGTTTTTACCAAGGCC
 ATCATAAGTGAGAACCCCTGTATAAAACATTATCACATCAAAGAAACAAATGACAGCCCCAAGCGCTACT
 ACGTGGCTGAGAAGTATGTGTTTACTCCATCCCTCTCCTCATCCAGTATCACCAGTACAATGGAGGAGG
 TTTGGTCACTCGACTCCGCTATCCAGTTTGCTCCTGGAGACAAAAAGCCCTGTACAGCAGGGCTAAGA
 TATGGGAAGTGGGTGATCCAACCCTCAGAGCTAACGTTTCGTGCAGGAGATTGGCAGCGGGCAGTTTGGGC
 TGGTGCATCTCGGCTACTGGCTCAACAAGGACAAGGTGGCCATCAAGACCATTGAGGAAGGGGCGATGTC
 AGAAGAAGACTTTATCGAGGAGCGGAAGTCATGATGAAACTCTCTACCCCAAACCTGGTGCAGCTCTAT
 GGGGTGTGCCTGGAGCAAGCCCCATCTGCCTGGTGTGTTGAGTTCATGGAGCACGGCTGCCTGTCGGATT
 ACCTTCGAAGTCAGCGGGTCTCTTTGCTGCGGAGACCCTGCTGGGCATGTGCCTGGATGTGTGTGAGGG
 CATGGCTACCTGAAAAAGCTTGTGTCATCCACAGAGACCTGGCAGCCAGAACTGTTTGGTGGGAGAA
 AACAGGTCAAGGTGTCGACTTTGGGATGACAAGATTTGTCCTTGATGATCAATATACCAGCTCCA
 CGGGCACCAAATCCCAGTGAAGTGGGCATCCCCAGAGGTGTTCTCTTTAGTCGCTATAGCAGCAAGTC
 AGATGTGTGGTCTTTGGTGTACTGATGTGGGAAGTCTTCAGTGAAGGCAAAATCCCATACGAAAACCGA
 AGCAATCCGAGGTCGTGGAAGATATCAGCACTGGCTTTTCGGTTATACAAGCCCCGCTGGCCTCCTGTC
 ATGCTACCAGATCATGAATCATTGCTGGAAGAGAAACCAAGACCGCCACCCTTCTCCAGCTGTT
 AAGCCAGCTGGCTGAAATCGCAGAAGCTGGGCTTAG

ACGCGTACGCGGCCGCTCGAGCAGAACTCATCTCAGAAGAGGATCTGGCAGCAAATGATATCCTGGATT
 ACAAGGATGACGACGATAAGGTTTAA

- Restriction Sites:** SgfI-MluI
- ACCN:** NM_001281968
- Insert Size:** 1857 bp
- OTI Disclaimer:** Our molecular clone sequence data has been matched to the reference identifier above as a point of reference. Note that the complete sequence of our molecular clones may differ from the sequence published for this corresponding reference, e.g., by representing an alternative RNA splicing form or single nucleotide polymorphism (SNP).
- OTI Annotation:** Clone contains native stop codon, and expresses the complete ORF without any c-terminal tag.
- Components:** The ORF clone is ion-exchange column purified and shipped in a 2D barcoded Matrix tube containing 10ug of transfection-ready, dried plasmid DNA (reconstitute with 100 ul of water).

Reconstitution Method:

1. Centrifuge at 5,000xg for 5min.
2. Carefully open the tube and add 100ul of sterile water to dissolve the DNA.
3. Close the tube and incubate for 10 minutes at room temperature.
4. Briefly vortex the tube and then do a quick spin (less than 5000xg) to concentrate the liquid at the bottom.
5. Store the suspended plasmid at -20°C. The DNA is stable for at least one year from date of shipping when stored at -20°C.

RefSeq: [NM_001281968.1](#), [NP_001268897.1](#)

RefSeq Size: 4271 bp

RefSeq ORF: 1857 bp

Locus ID: 16428

UniProt ID: [Q03526](#)

Cytogenetics: 11 27.75 cM

Gene Summary: Tyrosine kinase that plays an essential role in regulation of the adaptive immune response. Regulates the development, function and differentiation of conventional T-cells and nonconventional NKT-cells. When antigen presenting cells (APC) activate T-cell receptor (TCR), a series of phosphorylation lead to the recruitment of ITK to the cell membrane, in the vicinity of the stimulated TCR receptor, where it is phosphorylated by LCK. Phosphorylation leads to ITK autophosphorylation and full activation. Once activated, phosphorylates PLCG1, leading to the activation of this lipase and subsequent cleavage of its substrates. In turn, the endoplasmic reticulum releases calcium in the cytoplasm and the nuclear activator of activated T-cells (NFAT) translocates into the nucleus to perform its transcriptional duty. Phosphorylates 2 essential adapter proteins: the linker for activation of T-cells/LAT protein and LCP2. Then, a large number of signaling molecules such as VAV1 are recruited and ultimately lead to lymphokine production, T-cell proliferation and differentiation. Phosphorylates TBX21 at 'Tyr-525' and mediates its interaction with GATA3 (PubMed:15662016).[UniProtKB/Swiss-Prot Function]

Transcript Variant: This variant (5) uses two alternate in-frame splice sites in the central coding region, compared to variant 1. The encoded isoform (5) is shorter, compared to isoform 1.