

Product datasheet for **MC228642**

Brsk2 (NM_001276763) Mouse Untagged Clone

Product data:

Product Type:	Expression Plasmids
Product Name:	Brsk2 (NM_001276763) Mouse Untagged Clone
Tag:	Tag Free
Symbol:	Brsk2
Synonyms:	4833424K13Rik; SAD-A; SADA
Vector:	pCMV6-Entry (PS100001)
E. coli Selection:	Kanamycin (25 ug/mL)
Cell Selection:	Neomycin



[View online »](#)

Fully Sequenced ORF: >MC228642 representing NM_001276763
 Red=Cloning site Blue=ORF Orange=Stop codon

TTTTGTAATACGACTCACTATAGGGCGGCCGGAATTCGTCGACTGGATCCGGTACCGAGGAGATCTGCC
 GCC**CGATCGCC**

ATGACATCGACGGGAAGGACGGCGGGCGCAGCACGCGCAGTATGTGGGGCCCTACCGGCTGGAGA
 AGACGCTGGGCAAGGGGACAGAGGCTTGGTGAAGCTGGGAATCCACTGTGTCACTTGCCAGAAGGTCGC
 CATCAAAATCGTGAACCGTGAGAAGCTCAGTGAGTCGGTCTGATGAAGGTGGAGCGAGAGATTGCCATC
 TTGAAGCTCATCGAGCATCCACATGTAAGCTGCATGACGTCTATGAAAAACAAAAATATTTATAACC
 TGGTGCTAGAACATGTGTCTGGGGGAGAGCTGTTGACTACCTGGTGAAGAAGGGCCGGCTGACCCCAA
 GGAGGCCCGCAAGTTCTCCGGCAGATCATCTGCACTGGACTTCTGTACAGCCACTCCATATGCCAT
 AGAGACTTGAAGCCAGAGAACCTGCTGCTAGATGAGAGGAACAACATCCGTATTGCAGACTTTGGCATGG
 CATCCCTGCAGGTGGGAGACAGCCTGCTGGAGACCAGCTGCGGATCTCCACACTATGCCTGTCCGGAAGT
 GATTCGGGGCGAGAAGTATGATGGCCGCAAGGCAGATGTGTGGAGCTGTGGTGTGATCCTGTTCCGCTTG
 CTGGTGGGGGCTCTGCCTTTTATGATGATGACAACCTGCGGCAGTTGCTGGAGAAGGTCAAGCGTGGTGTG
 TCCACATGCCACACTTTATCCCACCAGACTGCCAGAGTCTCCTGCGTGGCATGATTGAGGTGGATGCAGC
 TCGGCGCCTCACGCTAGAGCACATTCAGAAACACATATGGTATATAGGTGGCAAGAATGAGCCAGAGCCC
 GAACAGCCCATCCCACGCAAGGTGCAGATCCGCTCACTACCCAGCTTGGAAAGACATTGACCTGATGTGT
 TGGACAGCATGCACTCACTGGGCTGCTCCGAGACCGCAACAAGTGTGTCAGGATCTGCTATCTGAGGA
 GGAAATCAGGAAAAGATGATTTATTTCTCCTCCTGGATCGGAAAGAACGGTATCCAAGCCATGAGGAT
 GAGGACCTGCCCCCAGGAATGAGATAGACCCTCCCCGGAAGCGTGTGGATTCCCGATGCTGAACCGGC
 ATGGCAAGCGGGACCTGAGCGCAAGTCCATGGAAGTGTGCTCAGTGTGACAGATGGTGGCTCCCCAGTGCC
 TGCACGGAGAGCCATTGAGATGGCCAGCATGGCCAGAGTAAAGCAGTGTTCAGTAAAAGCCTGGATATC
 GCTGAAGCCCATCCCAATTCAGCAAAGAAGACAGATCTCGATCCATCAGTGGTGCCTCAGGCCCTT
 CTACAAGTCCACTCAGCAGTCTCGGGTACCCCTCACCCCTCACCAAGGGGTAGTCCCCTTCTACCCC
 CAAAGGGACGCCTGTCCACACGCCAAAGGAGAGCCAGCTGGCACACCCAACCCACACCACCATCCAGC
 CCTAGTGTGGAGGAGTGCCTGGCGGACAGACTGAACTCCATCAAGAACAGCTTCTGGGCTCACCTC
 GATTCCACCGCCGAAACTCCAAGTTCACCGCCAGAGGAGATGTCCAACCTGACCCAGAATCCTCTCC
 AGAGCTGGCCAAGAAATCGTGGTTCGGGAACCTCATCAACCTGGAGAAGGAGGAGCAGATCTTTGTGGT
 ATCAAGGACAAGCCCTGAGCTCCATCAAGGCTGACATCGTTCATGCCTCCTGTGATCCCCAGCCTCA
 GCCACAGCGTTATTTCCAGACAAGCTTCAAGGCTGAATACAAGGCCACAGGGGGCCAGCAGTGTCCA
 GAAGCCGGTCAAGTTCAGGTGGACATCACTTACTGAGGGCGGAGAGGCCAGAAAGGAGAATGGCATC
 TACTCAGTCACATTCACCTTACTCTCAGGCCCCAGTCGCCGCTTCAAGAGGGTGGTGGAGACCATCCAGG
 CCCAGCTGTTAAGCACCCATGACCAGCCATCAGCCAGCACCTGTGAGGAATTATCCCGAAAAGT**TAA**

ACGCGTACGCGGCCGCTCGAGCAGAAACTCATCTCAGAAGAGGATCTGGCAGCAAATGATATCCTGGATT
 ACAAGGATGACGACGATAAGGTTTAA

Restriction Sites: Sgfl-MluI

ACCN: NM_001276763

Insert Size: 2028 bp

OTI Disclaimer: Our molecular clone sequence data has been matched to the reference identifier above as a point of reference. Note that the complete sequence of our molecular clones may differ from the sequence published for this corresponding reference, e.g., by representing an alternative RNA splicing form or single nucleotide polymorphism (SNP).

OTI Annotation: Clone contains native stop codon, and expresses the complete ORF without any c-terminal tag.

Components:	The ORF clone is ion-exchange column purified and shipped in a 2D barcoded Matrix tube containing 10ug of transfection-ready, dried plasmid DNA (reconstitute with 100 ul of water).
Reconstitution Method:	<ol style="list-style-type: none">1. Centrifuge at 5,000xg for 5min.2. Carefully open the tube and add 100ul of sterile water to dissolve the DNA.3. Close the tube and incubate for 10 minutes at room temperature.4. Briefly vortex the tube and then do a quick spin (less than 5000xg) to concentrate the liquid at the bottom.5. Store the suspended plasmid at -20°C. The DNA is stable for at least one year from date of shipping when stored at -20°C.
RefSeq:	<u>NM_001276763.1, NP_001263692.1</u>
RefSeq Size:	4113 bp
RefSeq ORF:	2028 bp
Locus ID:	75770
Cytogenetics:	7 F5
Gene Summary:	<p>Serine/threonine-protein kinase that plays a key role in polarization of neurons and axonogenesis, cell cycle progress and insulin secretion. Phosphorylates CDK16, CDC25C, MAPT/TAU, PAK1 and WEE1. Following phosphorylation and activation by STK11/LKB1, acts as a key regulator of polarization of cortical neurons, probably by mediating phosphorylation of microtubule-associated proteins such as MAPT/TAU at 'Thr-504' and 'Ser-554'. Also regulates neuron polarization by mediating phosphorylation of WEE1 at 'Ser-642' in post-mitotic neurons, leading to down-regulate WEE1 activity in polarized neurons. Plays a role in the regulation of the mitotic cell cycle progress and the onset of mitosis. Plays a role in the regulation of insulin secretion in response to elevated glucose levels, probably via phosphorylation of CDK16 and PAK1. While BRSK2 phosphorylated at Thr-175 can inhibit insulin secretion (PubMed:22798068), BRSK2 phosphorylated at Thr-261 can promote insulin secretion (PubMed:22669945). Regulates reorganization of the actin cytoskeleton. May play a role in the apoptotic response triggered by endoplasmic reticulum (ER) stress.</p> <p>[UniProtKB/Swiss-Prot Function]</p> <p>Transcript Variant: This variant (4) has multiple differences in the coding region and 3' UTR, compared to variant 3. The resulting protein (isoform delta) is shorter than isoform gamma.</p> <p>Sequence Note: The RefSeq transcript and protein were derived from genomic sequence to make the sequence consistent with the reference genome assembly. The genomic coordinates used for the transcript record were based on alignments.</p>