

## Product datasheet for MC228756

### Cpeb3 (NM\_001290827) Mouse Untagged Clone

#### Product data:

Product Type:	Expression Plasmids
Product Name:	Cpeb3 (NM_001290827) Mouse Untagged Clone
Tag:	Tag Free
Symbol:	Cpeb3
Synonyms:	4831444O18Rik; CPE-BP3; mKIAA0940
Mammalian Cell Selection:	Neomycin
Vector:	pCMV6-Entry (PS100001)
E. coli Selection:	Kanamycin (25 ug/mL)

**OriGene Technologies, Inc.**  
9620 Medical Center Drive, Ste 200  
Rockville, MD 20850, US  
Phone: +1-888-267-4436  
<https://www.origene.com>  
[techsupport@origene.com](mailto:techsupport@origene.com)  
EU: [info-de@origene.com](mailto:info-de@origene.com)  
CN: [techsupport@origene.cn](mailto:techsupport@origene.cn)



[View online »](#)

This product is to be used for laboratory only. Not for diagnostic or therapeutic use.

©2024 OriGene Technologies, Inc., 9620 Medical Center Drive, Ste 200, Rockville, MD 20850, US

1 / 4

Fully Sequenced ORF: >MC228756 representing NM\_001290827  
Red=Cloning site Blue=ORF Orange=Stop codon

TTTTGTAATACGACTCACTATAGGGCGCCGGAAATTCTCGACTGGATCCGTACCGAGGAGATCTGCC  
GCCGCGATCGCC

ATGCAGGATGTTACTGATGGACAAAGCAAACCCAGCCCCAGTCTCAGCAGCAGCAGCGGCAGCAGC  
AGCAGCAGCAGCACAGCTCCAGCCCGAGCCCGCGCAGCTGAAGCCCCGTCCAGCCCCTCTCCTCAGA  
GATCCCAAGCCCGAAGACAGTAGCGCAGTGCCGGCCCTCAGCCCCCGCTCGGCCAGCCCCAAC  
GGCCCCGACAAGATGCAGATGGAGTCGCCGCTCCTGCCGGCTTGAGTTTCCATCAGCCCCCTCAGCAGC  
CGCCGCCGCCGAGGAGCCCACGGGCCGGAGCGCTGCTGCCGTCTCGGCCAGCACCTGGTCCAC  
AGGCACTACAAACCGGTGGAGGACAGCTTCCAGGGATCACCCAGTCACGGGACCATGCTCTTC  
CAAAACTCCCGCACCGTCAACCCGGTCTCGGAGGCACCTCTCGCCGAGATCGGCTGGCGCAGA  
CCCAGCACCATCAGCAGCCTCACCGCCCGCCGAGCCGCCGAGCCCCCGCAGGGCAG  
GCCCTCGCAGCAACGGCCTCGCCGCCAGCCCCAGCCAGGGCCCTATGCGCAGAGGAGCGCCGCTGCC  
TACGGCCACCAGCCATCATGACCAGCAAGCGCTCTCGCCGAGGAGTCAGCGTGAACGCCGCTGGAGTGTCCG  
CCGCTGCCTCTCGGCTCGTCAGCTGGAACACGCACAGAGCGTGAACGCCGCTGGAGTGTCCG  
CAACCCGTGGGTGGCTCGAGCGGGCCAGACCCCTGCCGCCGGTCGGCTGGCGTGGGTAGT  
GTGGGGTACCCCTCCCGTTAACCCATCTCGCCGCTAAAAGCCCTTCCAGCAATGTGATCGCTC  
CACCCAAGTTCCCTCGTCAGCCCGCTCACCTCAAGTCTGGATGGAGGATAACGTTCCGGACCGA  
TAATGGTAACAATCTGTTGCCCTTCAGGACGGAGTAGGCCCTATGACACTTTAACTTGACTCCTG  
GAGAACTCCTTAATGGATATGATAAGGACTGACCATGAGCCTCTGAAAGGTAACACTACCCCTCCAGTG  
GCCCAACATGAGTTCGTGTATATAATGAGGAAATCATTTGCAAGGACGATGGAAATAATTCCA  
CCATCCAGGAACAGATAACATTATGGACTTAACACCAGGAGCTATGGCGGAGACGAGGTCGGTCT  
CTCTTCCCTTGAAGACGCCCTCTGGATGACAGCCATGGTGTACGGCCTATCGTCTGGCTTAAGTT  
CTCCTACTCGATGTCAAAATGGGAACGAGTGGAACGCTACTCTAGAAAGGTGTTGGAGGCCCTCC  
TCCTGACATCGATGAAGATGAGATCACTGCCAGCTTCAGGTTGGACCCCTGTGGTGGACTGGCCT  
CATAAAGCCGAGAGCAAATCGTACTCCCTCTAAAGGCTATGCTTCTGCTGTTCCAGGAGGAGCT  
CAGTGAAGCTCTGATAGATGCCGCTGGAGGAGTAGGAAACTGTACCTGTGTGTCCAGCCAAC  
CATCAAGGATAAACGGTGCCTGGAAACCTAACGACTGACAGTGAACGTTGAAATGGATGGCTCT  
CAGCCTTGGACCCAGAAAACATCTTGGGGAGTTCCAGCACCCCTGAGCTGTTGAACCTGG  
CAATGATAATGGACCGTTGTACGGTGTGTTGCTATGCTGGCATTGACACAGACCCAGAGCTGAAGTA  
CCCCAAAGGTGCTGGCGTGTGCTCCAAATCAGCAGAGCTACATTGCAAGCCATCAGGCCGCTT  
GTGCAACTTCAACAACGACATTGACAAACGGGTGAAAGTGAAGCCGTACGTGCTGGATGATCAGATGT  
GTGACGAGTGTGAGGGCACACGCTGCCGGGGAGTTGCCCTTGTGCCAACGTACCTGTCT  
GCAGTACTACTGTGAATACTGCTGCCGCCAGCATCCACTCCGAGCCGCCGTGAGTCCACAAACCGCTG  
GTGAGGAGGGAGGCACGCCCGGCCAGTCCGCTGGAGGCTGA

ACCGTACGCGCCGCTCGAGCAGAAACTCATCTCAGAAGAGGATCTGGCAGCAAATGATATCCTGGATT  
ACAAGGATGACGACGATAAGTTAA

Restriction Sites: Sgfl-Mlul

ACCN: NM\_001290827

Insert Size: 2151 bp

OTI Disclaimer: Our molecular clone sequence data has been matched to the reference identifier above as a point of reference. Note that the complete sequence of our molecular clones may differ from the sequence published for this corresponding reference, e.g., by representing an alternative RNA splicing form or single nucleotide polymorphism (SNP).

<b>OTI Annotation:</b>	Clone contains native stop codon, and expresses the complete ORF without any c-terminal tag.
<b>Components:</b>	The ORF clone is ion-exchange column purified and shipped in a 2D barcoded Matrix tube containing 10ug of transfection-ready, dried plasmid DNA (reconstitute with 100 ul of water).
<b>Reconstitution Method:</b>	<ol style="list-style-type: none"><li>1. Centrifuge at 5,000xg for 5min.</li><li>2. Carefully open the tube and add 100ul of sterile water to dissolve the DNA.</li><li>3. Close the tube and incubate for 10 minutes at room temperature.</li><li>4. Briefly vortex the tube and then do a quick spin (less than 5000xg) to concentrate the liquid at the bottom.</li><li>5. Store the suspended plasmid at -20°C. The DNA is stable for at least one year from date of shipping when stored at -20°C.</li></ol>
<b>Note:</b>	Plasmids are not sterile. For experiments where strict sterility is required, filtration with 0.22um filter is required.
<b>RefSeq:</b>	<u><a href="#">NM_001290827.1</a></u> , <u><a href="#">NP_001277756.1</a></u>
<b>RefSeq Size:</b>	5908 bp
<b>RefSeq ORF:</b>	2151 bp
<b>Locus ID:</b>	208922
<b>UniProt ID:</b>	<u><a href="#">Q7TN99</a></u>
<b>Cytogenetics:</b>	19 C2

Gene Summary:	Sequence-specific RNA-binding protein which acts as a translational repressor in the basal unstimulated state but, following neuronal stimulation, acts as a translational activator (PubMed:17024188, PubMed:26074072). In contrast to CPEB1, does not bind to the cytoplasmic polyadenylation element (CPE), a uridine-rich sequence element within the mRNA 3' UTR, but binds to a U-rich loop within a stem-loop structure (PubMed:17024188). Required for the consolidation and maintenance of hippocampal-based long term memory (PubMed:26074003). In the basal state, binds to the mRNA 3' UTR of the glutamate receptors GRIA1 and GRIA2 and negatively regulates their translation (PubMed:17024188, PubMed:22153079). Also represses the translation of DLG4, GRIN1, GRIN2A and GRIN2B (PubMed:24155305). When activated, acts as a translational activator of GRIA1 and GRIA2 (PubMed:22153079, PubMed:26074003). In the basal state, suppresses SUMO2 translation but activates it following neuronal stimulation (PubMed:26074071). Binds to the 3' UTR of TRPV1 mRNA and represses TRPV1 translation which is required to maintain normal thermoception (PubMed:26915043). Binds actin mRNA, leading to actin translational repression in the basal state and to translational activation following neuronal stimulation (PubMed:26074072). Negatively regulates target mRNA levels by binding to TOB1 which recruits CNOT7/CAF1 to a ternary complex and this leads to target mRNA deadenylation and decay (By similarity). In addition to its role in translation, binds to and inhibits the transcriptional activation activity of STAT5B without affecting its dimerization or DNA-binding activity. This, in turn, represses transcription of the STAT5B target gene EGFR which has been shown to play a role in enhancing learning and memory performance (By similarity). In contrast to CPEB1, CPEB2 and CPEB4, not required for cell cycle progression (By similarity). [UniProtKB/Swiss-Prot Function] Transcript Variant: This variant (3) differs in the 5' UTR, compared to variant 1. Variants 1, 2, and 3 encode the same isoform (a). Sequence Note: This RefSeq record was created from transcript and genomic sequence data to make the sequence consistent with the reference genome assembly. The genomic coordinates used for the transcript record were based on transcript alignments.
---------------	---