

Product datasheet for **MG200193**

Fcer1g (BC034163) Mouse Tagged ORF Clone

Product data:

Product Type:	Expression Plasmids
Product Name:	Fcer1g (BC034163) Mouse Tagged ORF Clone
Tag:	TurboGFP
Symbol:	Fcer1g
Synonyms:	FcR[g], FcR-gamma, CD23
Mammalian Cell Selection:	Neomycin
Vector:	pCMV6-AC-GFP (PS100010)
E. coli Selection:	Ampicillin (100 ug/mL)
ORF Nucleotide Sequence:	<p>>MG200193 representing BC034163 Red=Cloning site Blue=ORF Green=Tags(s)</p> <p>TTTGTGTAATACGACTCACTATAGGGCGGCCGGAATTCGTCGACTGGATCCGGTACCGAGGAGATCTGCCGCCCGATCGCC</p> <p>ATGATCTCAGCCGTGATCTTGTCTTGCTCCTTTTGGTGAACAAGCAGCCGCCCTGGGAGAGCCGCAGCTCTGTATATCCTGGATGCTGCTCTGTTTTGTATGGTATTGTCCTTACCCTACTCTACTGTCGACTCAAGATCCAGGTCCGAAAGGCAGCTATAGCCAGCCGTGAGAAAGCAGATGCTGTCTACACGGGCTGAACACCCGAGCCAGGAGACATATGAGACTCTGAAGCATGAGAAACCACCCAG</p> <p>ACGCGTACGCGGCCGCTCGAG - GFP Tag - GTTTAA</p>
Protein Sequence:	<p>>MG200193 representing BC034163 Red=Cloning site Green=Tags(s)</p> <p>MISAVILFLLLLVEQAAALGEPQLCYILDAVLFLYGIVLTLLYCRLKIQVRKAAIASREKADAVYTGLNTRSQETYETLKHEKPPQ</p> <p>TRTRPLE - GFP Tag - V</p>
Restriction Sites:	SgfI-MluI



[View online »](#)

Cloning Scheme:



ACCN:	BC034163
ORF Size:	258 bp
OTI Disclaimer:	The molecular sequence of this clone aligns with the gene accession number as a point of reference only. However, individual transcript sequences of the same gene can differ through naturally occurring variations (e.g. polymorphisms), each with its own valid existence. This clone is substantially in agreement with the reference, but a complete review of all prevailing variants is recommended prior to use. More info
OTI Annotation:	This clone was engineered to express the complete ORF with an expression tag. Expression varies depending on the nature of the gene.
Components:	The ORF clone is ion-exchange column purified and shipped in a 2D barcoded Matrix tube containing 10ug of transfection-ready, dried plasmid DNA (reconstitute with 100 ul of water).
Reconstitution Method:	<ol style="list-style-type: none"> 1. Centrifuge at 5,000xg for 5min. 2. Carefully open the tube and add 100ul of sterile water to dissolve the DNA. 3. Close the tube and incubate for 10 minutes at room temperature. 4. Briefly vortex the tube and then do a quick spin (less than 5000xg) to concentrate the liquid at the bottom. 5. Store the suspended plasmid at -20°C. The DNA is stable for at least one year from date of shipping when stored at -20°C.
Note:	Plasmids are not sterile. For experiments where strict sterility is required, filtration with 0.22um filter is required.

RefSeq: [BC034163](#), [AAH34163](#)

RefSeq Size: 800 bp


RefSeq ORF: 260 bp

Locus ID: 14127

Cytogenetics: 1 79.23 cM

Gene Summary: Adapter protein containing an immunoreceptor tyrosine-based activation motif (ITAM) that transduces activation signals from various immunoreceptors. As a component of the high-affinity immunoglobulin E (IgE) receptor, mediates allergic inflammatory signaling in mast cells (PubMed:14764707). As a constitutive component of interleukin-3 receptor complex, selectively mediates interleukin 4/IL4 production by basophils, priming T-cells toward effector T-helper 2 subset (PubMed:19098920). Associates with pattern recognition receptors CLEC4D and CLEC4E to form a functional signaling complex in myeloid cells. Binding of mycobacterial trehalose 6,6'-dimycolate (TDM) to this receptor complex leads to phosphorylation of ITAM, triggering activation of SYK, CARD9 and NF-kappa-B, consequently driving maturation of antigen-presenting cells and shaping antigen-specific priming of T-cells toward effector T-helper 1 and T-helper 17 cell subtypes (PubMed:23602766) (Probable). May function cooperatively with other activating receptors. Functionally linked to integrin beta-2/ITGB2-mediated neutrophil activation (PubMed:17086186). Also involved in integrin alpha-2/ITGA2-mediated platelet activation (PubMed:9171347).[UniProtKB/Swiss-Prot Function]

Product images:



Circular map for MG200193