

Product datasheet for **MG200241**

Hmgn2 (BC083085) Mouse Tagged ORF Clone

Product data:

Product Type: Expression Plasmids
Product Name: Hmgn2 (BC083085) Mouse Tagged ORF Clone
Tag: TurboGFP
Symbol: Hmgn2
Synonyms: HMG-17; Hmg17
Mammalian Cell Selection: Neomycin
Vector: pCMV6-AC-GFP (PS100010)
E. coli Selection: Ampicillin (100 ug/mL)
ORF Nucleotide Sequence: >MG200241 representing BC083085
 Red=Cloning site Blue=ORF Green=Tags(s)

TTTTGTAATACGACTCACTATAGGGCGGCCGGAATTCGTCGACTGGATCCGGTACCGAGGAGATCTGCC
 GCC**CGATCGCC**

ATGCCAAAAGAAAGGCTGAAGGGGATGCTAAAGGAGACAAAACCAAGGTGAAGGACGAGCCACAGAGAA
 GATCTGCAAGGTTGTCTGCTGTAAGTAAACCTGCTCCTCAAAGCCAGAGCCCAAACCTAAAAGGCCCC
 TGCGAAGAAGGGAGAGAAGGTACCAAGGGGAAGAAGGGAAAGCTGACGCCGGGAAGGATGCGAATAAT
 CCTGCAGAAAATGGAGATGCCAAAACAGACCAGGCACAGAAAGCTGAAGGTGCTGGAGATGCCAAG

ACGCGTACGCGGCCGCTCGAG - GFP Tag - GTTTAA

Protein Sequence: >MG200241 representing BC083085
 Red=Cloning site Green=Tags(s)

MPKRKAEGDAKGDKTKVKDEPQRRSARLSAVSKPAPPKPEPKPKKAPAKKGEKVPKGGKADAGKDANN
 PAENGDAKTDQAQKAEGAGDAK

TRTRPLE - GFP Tag - V

Restriction Sites: Sgfl-MluI



[View online »](#)

OTI Annotation:	This clone was engineered to express the complete ORF with an expression tag. Expression varies depending on the nature of the gene.
Components:	The ORF clone is ion-exchange column purified and shipped in a 2D barcoded Matrix tube containing 10ug of transfection-ready, dried plasmid DNA (reconstitute with 100 ul of water).
Reconstitution Method:	<ol style="list-style-type: none">1. Centrifuge at 5,000xg for 5min.2. Carefully open the tube and add 100ul of sterile water to dissolve the DNA.3. Close the tube and incubate for 10 minutes at room temperature.4. Briefly vortex the tube and then do a quick spin (less than 5000xg) to concentrate the liquid at the bottom.5. Store the suspended plasmid at -20°C. The DNA is stable for at least one year from date of shipping when stored at -20°C.
RefSeq:	BC083085 , AAH83085
RefSeq Size:	1349 bp
RefSeq ORF:	278 bp
Locus ID:	15331
Cytogenetics:	4 66.47 cM
Gene Summary:	Binds to the inner side of the nucleosomal DNA thus altering the interaction between the DNA and the histone octamer. May be involved in the process which maintains transcribable genes in a unique chromatin conformation (By similarity).[UniProtKB/Swiss-Prot Function]