

# **Product datasheet for MG200291**

## Cxcl10 (BC030067) Mouse Tagged ORF Clone

**Product data:** 

**Product Type:** Expression Plasmids

**Product Name:** Cxcl10 (BC030067) Mouse Tagged ORF Clone

Tag: TurboGFP

Symbol: Cxcl10

**Synonyms:** CRG-2, IP-10, C7, INP10, mob-1, gIP-10, IP10

Mammalian Cell

Selection:

Neomycin

Vector: pCMV6-AC-GFP (PS100010)

E. coli Selection: Ampicillin (100 ug/mL)

ORF Nucleotide >MG200291 representing BC030067

Sequence: Red=Cloning site Blue=ORF Green=Tags(s)

TTTTGTAATACGACTCACTATAGGGCGGCCGGGAATTCGTCGACTGGATCCGGTACCGAGGAGATCTGCC

GCCGCGATCGCC

ATGAACCCAAGTGCTGCCGTCATTTTCTGCCTCATCCTGCTGGGTCTGAGTGGGACTCAAGGGATCCCTC
TCGCAAGGACGGTCCGCTGCAACTGCATCCATATCGATGACGGGCCAGTGAGAATGAGGGCCATAGGGAA
GCTTGAAATCATCCCTGCGAGCCTATCCTGCCCACGTGTTGAGATCATTGCCACGATGAAAAAGAATGAT
GAGCAGAGATGTCTGAATCCGGAATCTAAGACCATCAAGAATTTAATGAAAGCGTTTTAGCCAAAAAAGGT

CTAAAAGGGCTCCT

ACGCGTACGCGGCCGCTCGAG - GFP Tag - GTTTAA

**Protein Sequence:** >MG200291 representing BC030067

Red=Cloning site Green=Tags(s)

MNPSAAVIFCLILLGLSGTQGIPLARTVRCNCIHIDDGPVRMRAIGKLEIIPASLSCPRVEIIATMKKND

EQRCLNPESKTIKNLMKAFSQKRSKRAP

TRTRPLE - GFP Tag - V

**Restriction Sites:** Sgfl-Mlul




**OriGene Technologies, Inc.** 9620 Medical Center Drive, Ste 200


CN: techsupport@origene.cn

Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com




### **Cloning Scheme:**





### Plasmid Map:



**ACCN:** BC030067 **ORF Size:** 296 bp



**OTI Disclaimer:** 

Due to the inherent nature of this plasmid, standard methods to replicate additional amounts of DNA in E. coli are highly likely to result in mutations and/or rearrangements. Therefore, OriGene does not guarantee the capability to replicate this plasmid DNA. Additional amounts of DNA can be purchased from OriGene with batch-specific, full-sequence verification at a reduced cost. Please contact our customer care team at <a href="mailto:customport@origene.com">customport@origene.com</a> or by calling 301.340.3188 option 3 for pricing and delivery.

The molecular sequence of this clone aligns with the gene accession number as a point of reference only. However, individual transcript sequences of the same gene can differ through naturally occurring variations (e.g. polymorphisms), each with its own valid existence. This clone is substantially in agreement with the reference, but a complete review of all prevailing variants is recommended prior to use. <u>More info</u>

**OTI Annotation:** 

This clone was engineered to express the complete ORF with an expression tag. Expression varies depending on the nature of the gene.

Components:

The ORF clone is ion-exchange column purified and shipped in a 2D barcoded Matrix tube containing 10ug of transfection-ready, dried plasmid DNA (reconstitute with 100 ul of water).

**Reconstitution Method:** 

- 1. Centrifuge at 5,000xg for 5min.
- 2. Carefully open the tube and add 100ul of sterile water to dissolve the DNA.
- 3. Close the tube and incubate for 10 minutes at room temperature.
- 4. Briefly vortex the tube and then do a quick spin (less than 5000xg) to concentrate the liquid at the bottom.
- 5. Store the suspended plasmid at -20°C. The DNA is stable for at least one year from date of shipping when stored at -20°C.

**RefSeq:** <u>BC030067</u>, <u>AAH30067</u>

RefSeq Size:1087 bpRefSeq ORF:296 bpLocus ID:15945

**Cytogenetics:** 5 46.57 cM



#### **Gene Summary:**

Pro-inflammatory cytokine that is involved in a wide variety of processes such as chemotaxis, differentiation, and activation of peripheral immune cells, regulation of cell growth, apoptosis and modulation of angiostatic effects (By similarity) (PubMed:28623423). Plays thereby an important role during viral infections by stimulating the activation and migration of immune cells to the infected sites (PubMed:18624292, PubMed:19017990, PubMed:28468883). Mechanistically, binding of CXCL10 to the CXCR3 receptor activates G protein-mediated signaling and results in downstream activation of phospholipase C-dependent pathway, an increase in intracellular calcium production and actin reorganization. In turn, recruitment of activated Th1 lymphocytes occurs at sites of inflammation (By similarity). Activation of the CXCL10/CXCR3 axis plays also an important role in neurons in response to brain injury for activating microglia, the resident macrophage population of the central nervous system, and directing them to the lesion site. This recruitment is an essential element for neuronal reorganization (PubMed:15456824).[UniProtKB/Swiss-Prot Function]