

Product datasheet for **MG200390**

Rbx1 (NM_019712) Mouse Tagged ORF Clone

Product data:

Product Type: Expression Plasmids
Product Name: Rbx1 (NM_019712) Mouse Tagged ORF Clone
Tag: TurboGFP
Symbol: Rbx1
Synonyms: 1500002P15Rik; AA517855; ROC1
Mammalian Cell Selection: Neomycin
Vector: pCMV6-AC-GFP (PS100010)
E. coli Selection: Ampicillin (100 ug/mL)
ORF Nucleotide Sequence: >MG200390 representing NM_019712
Red=Cloning site Blue=ORF Green=Tags(s)

TTTTGTAATACGACTCACTATAGGGCGGCCGGAATTCGTCGACTGGATCCGGTACCGAGGAGATCTGCC
GCC**CGATCGCC**

ATGGCGGCGCGATGGATGTGGATACCCCAAGCGGCACCAACAGCGCGCGGGCAAGAAGCGCTTTGAAG
TTAAAAAGTGAATGCAGTGGCCCTCTGGGCTGGGACATTGTGGTTGATAACTGTGCCATCTGCAGGAA
CCACATTATGGATCTTTGTATCGAATGTCAGGCCAACCAGGCGTCAGCTACTCCGAAGAGTGTACGGTT
GCATGGGGAGTCTGCAACCATGCTTTTCATTTCCACTGCATCTCTCGATGGCTCAAACGAGGCAGGTGT
GTCCGTTGGACAACAGAGAGTGGGAGTCCAGAAGTATGGGCAT

ACGCGTACGCGGCCGCTCGAG - GFP Tag - GTTTAA

Protein Sequence: >MG200390 representing NM_019712
Red=Cloning site Green=Tags(s)
MAAAMDVDTPSGTNSGAGKKRFEVKKWNAVALWAWDIVVDNCAICRNHIMDLCEIQANQASATSEECTV
AWGVCNHAFHFHCISRWLKTRQVCPLDNREWEFQKYGH

TRTRPLE - GFP Tag - V

Restriction Sites: Sgfl-MluI



[View online »](#)

OTI Annotation:	This clone was engineered to express the complete ORF with an expression tag. Expression varies depending on the nature of the gene.
Components:	The ORF clone is ion-exchange column purified and shipped in a 2D barcoded Matrix tube containing 10ug of transfection-ready, dried plasmid DNA (reconstitute with 100 ul of water).
Reconstitution Method:	<ol style="list-style-type: none">1. Centrifuge at 5,000xg for 5min.2. Carefully open the tube and add 100ul of sterile water to dissolve the DNA.3. Close the tube and incubate for 10 minutes at room temperature.4. Briefly vortex the tube and then do a quick spin (less than 5000xg) to concentrate the liquid at the bottom.5. Store the suspended plasmid at -20°C. The DNA is stable for at least one year from date of shipping when stored at -20°C.
RefSeq:	NM_019712.3 , NP_062686.1
RefSeq Size:	1655 bp
RefSeq ORF:	327 bp
Locus ID:	56438
UniProt ID:	P62878
Cytogenetics:	15 E1
Gene Summary:	<p>E3 ubiquitin ligase component of multiple cullin-RING-based E3 ubiquitin-protein ligase (CRLs) complexes which mediate the ubiquitination and subsequent proteasomal degradation of target proteins, including proteins involved in cell cycle progression, signal transduction, transcription and transcription-coupled nucleotide excision repair (PubMed:22118460). CRLs complexes and ARIH1 collaborate in tandem to mediate ubiquitination of target proteins, ARIH1 mediating addition of the first ubiquitin on CRLs targets (By similarity). The functional specificity of the E3 ubiquitin-protein ligase complexes depends on the variable substrate recognition components (By similarity). As a component of the CSA complex promotes the ubiquitination of ERCC6 resulting in proteasomal degradation (By similarity). Through the RING-type zinc finger, seems to recruit the E2 ubiquitination enzyme, like CDC34, to the complex and brings it into close proximity to the substrate (By similarity). Probably also stimulates CDC34 autoubiquitination (By similarity). May be required for histone H3 and histone H4 ubiquitination in response to ultraviolet and for subsequent DNA repair (By similarity). Promotes the neddylation of CUL1, CUL2, CUL4 and CUL4 via its interaction with UBE2M (By similarity). Involved in the ubiquitination of KEAP1, ENC1 and KLHL41 (By similarity). In concert with ATF2 and CUL3, promotes degradation of KAT5 thereby attenuating its ability to acetylate and activate ATM (By similarity).[UniProtKB/Swiss-Prot Function]</p>