

## **Product datasheet for MG200817**

## Med10 (BC048638) Mouse Tagged ORF Clone

**Product data:** 

**Product Type:** Expression Plasmids

**Product Name:** Med10 (BC048638) Mouse Tagged ORF Clone

Tag: TurboGFP Symbol: Med10

**Synonyms:** AA959813; AI385605; C78613; D13Wsu50e

Mammalian Cell Neomycin

Selection:

**Vector:** pCMV6-AC-GFP (PS100010)

E. coli Selection: Ampicillin (100 ug/mL)

ORF Nucleotide >MG200817 representing BC048638

Sequence: Red=Cloning site Blue=ORF Green=Tags(s)

TTTTGTAATACGACTCACTATAGGGCGGCCGGGAATTCGTCGACTGGATCCGGTACCGAGGAGATCTGCC

GCCGCGATCGCC

ACGCGTACGCGGCCGCTCGAG - GFP Tag - GTTTAA

**Protein Sequence:** >MG200817 representing BC048638

Red=Cloning site Green=Tags(s)

MAEKFDHLEEHLEKFVENIRQLGIIVSDFQPSSQAGLSQKLNFIVTGLQDIDKCRQQLHDITVPLEVFEY IDQGRNPQLYTKECLERALAKNEQVKGKIDTMKKFKSLLIQELSKVFPEDMAKYRSIRGEDHPPS

TRTRPLE - GFP Tag - V

Restriction Sites: Sgfl-Mlul




**OriGene Technologies, Inc.** 9620 Medical Center Drive, Ste 200


CN: techsupport@origene.cn

Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com



## **Cloning Scheme:**





**ACCN:** BC048638 **ORF Size:** 407 bp

**OTI Disclaimer:** The molecular sequence of this clone aligns with the gene accession number as a point of

reference only. However, individual transcript sequences of the same gene can differ through naturally occurring variations (e.g. polymorphisms), each with its own valid existence. This clone is substantially in agreement with the reference, but a complete review of all prevailing

variants is recommended prior to use. More info

**OTI Annotation:** This clone was engineered to express the complete ORF with an expression tag. Expression

varies depending on the nature of the gene.

**Components:** The ORF clone is ion-exchange column purified and shipped in a 2D barcoded Matrix tube

containing 10ug of transfection-ready, dried plasmid DNA (reconstitute with 100 ul of water).

**Reconstitution Method:** 1. Centrifuge at 5,000xg for 5min.

2. Carefully open the tube and add 100ul of sterile water to dissolve the DNA.

3. Close the tube and incubate for 10 minutes at room temperature.

4. Briefly vortex the tube and then do a quick spin (less than 5000xg) to concentrate the liquid

at the bottom.

5. Store the suspended plasmid at -20°C. The DNA is stable for at least one year from date of

shipping when stored at -20°C.

**RefSeq:** <u>BC048638</u>, <u>AAH48638</u>

RefSeq Size: 888 bp RefSeq ORF: 407 bp



**Locus ID:** 28077


Cytogenetics: 13 35.55 cM

**Gene Summary:** Component of the Mediator complex, a coactivator involved in the regulated transcription of

nearly all RNA polymerase II-dependent genes. Mediator functions as a bridge to convey information from gene-specific regulatory proteins to the basal RNA polymerase II transcription machinery. Mediator is recruited to promoters by direct interactions with regulatory proteins and serves as a scaffold for the assembly of a functional preinitiation complex with RNA polymerase II and the general transcription factors (By similarity).

[UniProtKB/Swiss-Prot Function]

## **Product images:**



Circular map for MG200817