

## Product datasheet for **MG201267**

### Ubd (NM\_023137) Mouse Tagged ORF Clone

#### Product data:

**Product Type:** Expression Plasmids  
**Product Name:** Ubd (NM\_023137) Mouse Tagged ORF Clone  
**Tag:** TurboGFP  
**Symbol:** Ubd  
**Synonyms:** FAT10  
**Mammalian Cell Selection:** Neomycin  
**Vector:** pCMV6-AC-GFP (PS100010)  
**E. coli Selection:** Ampicillin (100 ug/mL)  
**ORF Nucleotide Sequence:** >MG201267 representing NM\_023137  
Red=Cloning site Blue=ORF Green=Tags(s)

TTTTGTAATACGACTCACTATAGGGCGGCCGGAATTCGTCGACTGGATCCGGTACCGAGGAGATCTGCC  
GCC**CGATCGCC**

ATGGCTTCTGTCCGCACCTGTGTTGTCGGTTCAGACCAATGGCGGTTAATGACCTTTGAGACCACTGAGA  
ATGACAAAGTGAAGAAGATAAATGAACATATTAGGTCCCAACCAAGGTCTCTGTACAGGACCAGATCCT  
TCTGCTAGACTCCAAAATCCTCAAGCCCCATCGAAAATTGTCATCCTATGGGATTGACAAGGAAACCACT  
ATCCACCTTACCCTGAAGGTGGTGAAGCCAGTGAAGAGCTGCCCTGTTTCTGGTGGAGTCCAAA  
ACGAGGGGCAAAGGCACCTCCTCCGAGTTCGAAGATCCAGCTCAGTGGCCAGGTGAAAGAGATGATCGA  
GAGTGTGACCTCTGTGATCCCTAAGAAGCAGGTTGTGAATTGCAACGGAAAGAAGCTGGAAGATGAAAAG  
ATCATGGCTGACTACAACATCAAGAGTGGCAGTTTCTCTTTCTGACAACACACTGCACTGGGGGA

**ACGCGT**ACGCGGCCGCTCGAG - GFP Tag - GTTTAA

**Protein Sequence:** >MG201267 representing NM\_023137  
Red=Cloning site Green=Tags(s)

MASVRTCVVRSDQWRLMTFETTENDKVKKINEHIRSQTKVSVQDQILLLDSKILKPHRKLSSYGIDKETT  
IHLTLKVVKPSDEELPLFLVESKNEGQRHLLRVRSSSSVAQVKEMIESVTSVIPKKQVVCNCGKLEDGK  
IMADYNIKSGSLLFLTHCTGG

**TRTRPLE** - GFP Tag - V

**Restriction Sites:** SgfI-MluI




[View online »](#)

Cloning Scheme:



Plasmid Map:



ACCN: NM\_023137

ORF Size: 486 bp

OTI Disclaimer: The molecular sequence of this clone aligns with the gene accession number as a point of reference only. However, individual transcript sequences of the same gene can differ through naturally occurring variations (e.g. polymorphisms), each with its own valid existence. This clone is substantially in agreement with the reference, but a complete review of all prevailing variants is recommended prior to use. [More info](#)

OTI Annotation: This clone was engineered to express the complete ORF with an expression tag. Expression varies depending on the nature of the gene.

<b>Components:</b>	The ORF clone is ion-exchange column purified and shipped in a 2D barcoded Matrix tube containing 10ug of transfection-ready, dried plasmid DNA (reconstitute with 100 ul of water).
<b>Reconstitution Method:</b>	<ol style="list-style-type: none"><li>1. Centrifuge at 5,000xg for 5min.</li><li>2. Carefully open the tube and add 100ul of sterile water to dissolve the DNA.</li><li>3. Close the tube and incubate for 10 minutes at room temperature.</li><li>4. Briefly vortex the tube and then do a quick spin (less than 5000xg) to concentrate the liquid at the bottom.</li><li>5. Store the suspended plasmid at -20°C. The DNA is stable for at least one year from date of shipping when stored at -20°C.</li></ol>
<b>RefSeq:</b>	<u>NM_023137.3, NP_075626.1</u>
<b>RefSeq Size:</b>	1006 bp
<b>RefSeq ORF:</b>	489 bp
<b>Locus ID:</b>	24108
<b>UniProt ID:</b>	<u>P63072</u>
<b>Cytogenetics:</b>	17 B1
<b>Gene Summary:</b>	Ubiquitin-like protein modifier which can be covalently attached to target protein and subsequently leads to their degradation by the 26S proteasome, in a NUB1-dependent manner. Probably functions as a survival factor. Promotes the expression of the proteasome subunit beta type-9 (PSMB9/LMP2). Regulates TNF-alpha-induced and LPS-mediated activation of the central mediator of innate immunity NF-kappa-B by promoting TNF-alpha-mediated proteasomal degradation of ubiquitinated-I-kappa-B-alpha. Required for TNF-alpha-induced p65 nuclear translocation in renal tubular epithelial cells (RTECs). May be involved in dendritic cell (DC) maturation, the process by which immature dendritic cells differentiate into fully competent antigen-presenting cells that initiate T-cell responses. Mediates mitotic non-disjunction and chromosome instability, in long-term in vitro culture and cancers, by abbreviating mitotic phase and impairing the kinetochore localization of MAD2L1 during the prometaphase stage of the cell cycle. May be involved in the formation of aggresomes when proteasome is saturated or impaired. Mediates apoptosis in a caspase-dependent manner, especially in renal epithelium and tubular cells during renal diseases.[UniProtKB/Swiss-Prot Function]