

## **Product datasheet for MG201303**

## Selenot (NM 001040396) Mouse Tagged ORF Clone

**Product data:** 

**Product Type:** Expression Plasmids

**Product Name:** Selenot (NM\_001040396) Mouse Tagged ORF Clone

Symbol: Selenot

**Synonyms:** 2810407C02Rik; 5730408P04Rik; Se; Selt

Mammalian Cell

Selection:

Neomycin

**Vector:** pCMV6-AC-GFP (PS100010)

E. coli Selection: Ampicillin (100 ug/mL)

ORF Nucleotide >MG201303 representing NM\_001040396
Sequence: Red=Cloning site Blue=ORF Green=Tags(s)

TTTTGTAATACGACTCACTATAGGGCGGCCGGGAATTCGTCGACTGGATCCGGTACCGAGAGATCTGCC

GCCGCGATCGCC

GGATTCAATCCCACATCATCGATCA

ACGCGTACGCGGCCGCTCGAG - GFP Tag - GTTTAA

**Protein Sequence:** >MG201303 representing NM\_001040396

Red=Cloning site Green=Tags(s)

MRLLLLLVAASAVVRSEASANLGGVPSKRLKMQYATGPLLKFQICVS\*GYRRVFEEYMRVISQRYPDIR IEGENYLPOPIYRHIASFLSVFKLVLIGLIIVGKDPFAFFGMOAPSIWOWGQENKVYACMMVFFLSNMIE

NQCMSTGAFEITLNDVPVWSKLESGHLPSMQQLVQILDNEMKLNVHMDSIPHHRS

TRTRPLE - GFP Tag - V

Restriction Sites: Sgfl-Mlul




**OriGene Technologies, Inc.** 9620 Medical Center Drive, Ste 200


CN: techsupport@origene.cn

Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com



## **Cloning Scheme:**





<sup>\*</sup> The last codon before the Stop codon of the ORF

**ACCN:** NM 001040396

OTI Disclaimer: The molecular sequence of this clone aligns with the gene accession number as a point of

reference only. However, individual transcript sequences of the same gene can differ through naturally occurring variations (e.g. polymorphisms), each with its own valid existence. This clone is substantially in agreement with the reference, but a complete review of all prevailing

variants is recommended prior to use. More info The expression of this clone is not

guaranteed due to the nature of selenoproteins.

**OTI Annotation:** This clone encodes a selenoprotein containing the rare amino acid selenocysteine (Sec). Sec is

encoded by UGA codon, which normally signals translational termination. Expression of this

clone is not guaranteed due to the nature of selenoproteins.

**Components:** The ORF clone is ion-exchange column purified and shipped in a 2D barcoded Matrix tube

containing 10ug of transfection-ready, dried plasmid DNA (reconstitute with 100 ul of water).

**Reconstitution Method:** 1. Centrifuge at 5,000xg for 5min.

2. Carefully open the tube and add 100ul of sterile water to dissolve the DNA.

3. Close the tube and incubate for 10 minutes at room temperature.

4. Briefly vortex the tube and then do a quick spin (less than 5000xg) to concentrate the liquid

at the bottom.

5. Store the suspended plasmid at -20°C. The DNA is stable for at least one year from date of

shipping when stored at -20°C.

**RefSeq:** <u>NM 001040396.3</u>

 RefSeq Size:
 3575 bp

 RefSeq ORF:
 588 bp

 Locus ID:
 69227




UniProt ID: P62342

Cytogenetics: 3 D

**Gene Summary:** This gene encodes a selenoprotein, containing a selenocysteine (Sec) residue at the active

site. Sec is encoded by the UGA codon that normally signals translation termination. The 3' UTRs of selenoprotein mRNAs contain a conserved stem-loop structure, the Sec insertion sequence (SECIS) element, that is necessary for the recognition of UGA as a Sec codon rather than as a stop signal. This protein is localized in the endoplasmic reticulum. It belongs to the SelWTH family that possesses a thioredoxin-like fold and a conserved CxxU (C is cysteine, U is Sec) motif found in several redox active proteins. Studies in mice indicate a crucial role for this gene in the protection of dopaminergic neurons against oxidative stress in Parkinson's disease, and in the control of glucose homeostasis in pancreatic beta-cells. A pseudogene of this locus has been identified on chromosome 8. [provided by RefSeq, Aug 2017]

## **Product images:**



Circular map for MG201303