

## **Product datasheet for MG201511**

## Phca (BC023924) Mouse Tagged ORF Clone

**Product data:** 

**Product Type:** Expression Plasmids

**Product Name:** Phca (BC023924) Mouse Tagged ORF Clone

Tag: TurboGFP

Symbol: Phca

**Synonyms:** 1110057L18Rik; 5430429L08Rik; AV015045; Phca

Mammalian Cell Neomycin

Selection:

Vector: pCMV6-AC-GFP (PS100010)

E. coli Selection: Ampicillin (100 ug/mL)

ORF Nucleotide >MG201511 representing BC023924

Sequence: Red=Cloning site Blue=ORF Green=Tags(s)

TTTTGTAATACGACTCACTATAGGGCGGCCGGGAATTCGTCGACTGGATCCGGTACCGAGGAGATCTGCC

GCCGCGATCGCC

TTACATGTGTATCTCCAGAGTCTTGTCTGTAC

ACGCGTACGCGGCCGCTCGAG - GFP Tag - GTTTAA

**Protein Sequence:** >MG201511 representing BC023924

Red=Cloning site Green=Tags(s)

MAPAVDRKGYWGPTTSTLDWCEENYVVTLFVAEFWNTVSNLIMIIPPIFGAIQGIRDRLEKRYIAAYLAL TVVGMGSWCFHMTLKYEMQLLDELPMIYSCCIFVYCMFECFKTKSSINYHLLFTLFLYSLTVTTIYLKVK

**EPIFHQVMYGMLVFTLVLRSIYIVTCVSPESCLY** 

TRTRPLE - GFP Tag - V

**Restriction Sites:** Sgfl-Mlul




**OriGene Technologies, Inc.** 9620 Medical Center Drive, Ste 200


CN: techsupport@origene.cn

Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com




## **Cloning Scheme:**





## Plasmid Map:



**ACCN:** BC023924 **ORF Size:** 524 bp

**OTI Disclaimer:** The molecular sequence of this clone aligns with the gene accession number as a point of

reference only. However, individual transcript sequences of the same gene can differ through naturally occurring variations (e.g. polymorphisms), each with its own valid existence. This clone is substantially in agreement with the reference, but a complete review of all prevailing

variants is recommended prior to use. More info

OTI Annotation: This clone was engineered to express the complete ORF with an expression tag. Expression

varies depending on the nature of the gene.



**Components:** The ORF clone is ion-exchange column purified and shipped in a 2D barcoded Matrix tube

containing 10ug of transfection-ready, dried plasmid DNA (reconstitute with 100 ul of water).

**Reconstitution Method:** 1. Centrifuge at 5,000xg for 5min.

2. Carefully open the tube and add 100ul of sterile water to dissolve the DNA.

3. Close the tube and incubate for 10 minutes at room temperature.

4. Briefly vortex the tube and then do a quick spin (less than 5000xg) to concentrate the liquid

at the bottom.

5. Store the suspended plasmid at -20°C. The DNA is stable for at least one year from date of

shipping when stored at -20°C.

**RefSeq:** <u>BC023924</u>, <u>AAH23924</u>

RefSeq Size: 1460 bp
RefSeq ORF: 524 bp
Locus ID: 66190
Cytogenetics: 7 E1

**Gene Summary:** Endoplasmic reticulum and Golgi ceramidase that catalyzes the hydrolysis of unsaturated

long-chain C18:1-, C20:1- and C20:4-ceramides, dihydroceramides and phytoceramides into sphingoid bases like sphingosine and free fatty acids at alkaline pH (PubMed:26474409). Ceramides, sphingosine, and its phosphorylated form sphingosine-1-phosphate are bioactive

lipids that mediate cellular signaling pathways regulating several biological processes including cell proliferation, apoptosis and differentiation (PubMed:26474409). Controls the generation of sphingosine in erythrocytes, and thereby sphingosine-1-phosphate in plasma

(By similarity). Through the regulation of ceramides and sphingosine-1-phosphate  $\,$ 

homeostasis in the brain may play a role in neurons survival and function

(PubMed:26474409). By regulating the levels of proinflammatory ceramides in immune cells and tissues, may modulate the inflammatory response (PubMed:26938296).[UniProtKB/Swiss-

Prot Function]