

Product datasheet for MG201707

Cd160 (NM_018767) Mouse Tagged ORF Clone

Product data:

OriGene Technologies, Inc.


9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

Product Type:	Expression Plasmids
Product Name:	Cd160 (NM_018767) Mouse Tagged ORF Clone
Tag:	TurboGFP
Symbol:	Cd160
Synonyms:	AU045688; By55
Mammalian Cell Selection:	Neomycin
Vector:	pCMV6-AC-GFP (PS100010)
E. coli Selection:	Ampicillin (100 ug/mL)
ORF Nucleotide Sequence:	>MG201707 representing NM_018767 Red=Cloning site Blue=ORF Green=Tags(s)
	TTTTGTAATACGACTCACTATAGGGCGGCCGGGAATTCGTCGACTGGATCCGGTACCGAGGAGATCTGCC GCC <mark>GCGATCGC</mark> C
	ATGCAAAGAATCCTGATGGCCCCTGGCCAAAGCTGCTGTGCCCTGGCCATCCTGCTGGCAATTGTGAACT TCCAACATGGTGGATGTATTCATGTCACCAGCTCAGCATCCCAGAAAGGAGGGCGACTGGACCTCACCTG TACTTTGTGGCACAAGAAAGACGAAGCTGAGGGGGCTAATACTCTTCTGGTGCAAAGACAATCCTTGGAAC TGTTCCCCTGAGACCAGCTTAGAACAGCTTAGGGTTAAAAGGGATCCTGAGACAGATGGCATCACTGAAA AGTCATCTCAGTTGGTGTTCACCATAGAACAAGCTACACCATCAGACAGTGGGACCTACCAGTGCTGTGC CAGAAGCCAGAAACCAGAAATCTACATTCATGGCCACTTTCTCTCCGTTCTAGTCACAGGGAACCACACA GAGATAAGACAGAGACAAAGGTCACACCCTGACTTCAGCCATATCAACGGCACTCTCAGTTCAGGCTTCC TGCAAGTAAAGGCTTGGGGGATGTTGGTCACCAGCCTGGTGGCCCTTCAAGCTCTATATACCTTG ACGCCTACGCGGCCGCTCCAG - GFP Tag - GTTTAA
Protein Sequence:	<pre>>MG201707 representing NM_018767 Red=Cloning site Green=Tags(s)</pre>
	MQRILMAPGQSCCALAILLAIVNFQHGGCIHVTSSASQKGGRLDLTCTLWHKKDEAEGLILFWCKDNPWN CSPETSLEQLRVKRDPETDGITEKSSQLVFTIEQATPSDSGTYQCCARSQKPEIYIHGHFLSVLVTGNHT EIRQRQRSHPDFSHINGTLSSGFLQVKAWGMLVTSLVALQALYTL
	TRTRPLE - GFP Tag - V
Restriction Sites:	Sgfl-Mlul




This product is to be used for laboratory only. Not for diagnostic or therapeutic use. ©2022 OriGene Technologies, Inc., 9620 Medical Center Drive, Ste 200, Rockville, MD 20850, US

Cloning Scheme:



Plasmid Map:



ACCN:	NM_018767
ORF Size:	2510 bp
OTI Disclaimer:	The molecular sequence of this clone aligns with the gene accession number as a point of reference only. However, individual transcript sequences of the same gene can differ through naturally occurring variations (e.g. polymorphisms), each with its own valid existence. This clone is substantially in agreement with the reference, but a complete review of all prevailing variants is recommended prior to use. <u>More info</u>
OTI Annotation:	This clone was engineered to express the complete ORF with an expression tag. Expression varies depending on the nature of the gene.

This product is to be used for laboratory only. Not for diagnostic or therapeutic use. ©2022 OriGene Technologies, Inc., 9620 Medical Center Drive, Ste 200, Rockville, MD 20850, US

Cd160 (NM_018767) Mouse Tagged ORF Clone – MG201707		
Components:	The ORF clone is ion-exchange column purified and shipped in a 2D barcoded Matrix tube containing 10ug of transfection-ready, dried plasmid DNA (reconstitute with 100 ul of water).	
Reconstitution Method:	 Centrifuge at 5,000xg for 5min. Carefully open the tube and add 100ul of sterile water to dissolve the DNA. Close the tube and incubate for 10 minutes at room temperature. Briefly vortex the tube and then do a quick spin (less than 5000xg) to concentrate the liquid at the bottom. Store the suspended plasmid at -20°C. The DNA is stable for at least one year from date of shipping when stored at -20°C. 	
RefSeq:	<u>NM 018767.2</u>	
RefSeq Size:	1106 bp	
RefSeq ORF:	558 bp	
Locus ID:	54215	
UniProt ID:	<u>088875</u>	
Cytogenetics:	3 F2.1	
Gene Summary:	CD160 antigen: Receptor on immune cells capable to deliver stimulatory or inhibitory signals that regulate cell activation and differentiation. Exists as a GPI-anchored and as a transmembrane form, each likely initiating distinct signaling pathways via phosphoinositol 3-kinase in activated NK cells and via LCK and CD247/CD3 zeta chain in activated T cells (By similarity). Receptor for both classical and non-classical MHC class I molecules (PubMed:16177084). Receptor or ligand for TNF superfamily member TNFRSF14, participating in bidirectional cell-cell contact signaling between antigen presenting cells and lymphocytes. Upon ligation of TNFRSF14, provides stimulatory signal to NK cells enhancing IFNG production	

~ \$1/-

Upon ligation of TNFRSF14, provides stimulatory signal to NK cells enhancing IFNG production and anti-tumor immune response (PubMed:25711213). On activated CD4+ T cells, interacts with TNFRSF14 and downregulates CD28 costimulatory signaling, restricting memory and alloantigen-specific immune response (By similarity). In the context of bacterial infection, acts as a ligand for TNFRSF14 on epithelial cells, triggering the production of antimicrobial proteins and proinflammatory cytokines (PubMed:22801499).[UniProtKB/Swiss-Prot Function]

This product is to be used for laboratory only. Not for diagnostic or therapeutic use. ©2022 OriGene Technologies, Inc., 9620 Medical Center Drive, Ste 200, Rockville, MD 20850, US