


## Product datasheet for MG202058

## Gpx1 (NM\_008160) Mouse Tagged ORF Clone

## **Product data:**

Product Type:	Expression Plasmids
Product Name:	Gpx1 (NM_008160) Mouse Tagged ORF Clone
Symbol:	Gpx1
Synonyms:	Al195024; AL033363; CGP; CGPx; Gp; Gpx; GPx-; GPx-1; GSHPx; GSHPx-1
Mammalian Cell Selection:	Neomycin
Vector:	pCMV6-AC-GFP (PS100010)
E. coli Selection:	Ampicillin (100 ug/mL)
<b>Restriction Sites:</b>	Sgfl-Mlul
Cloning Scheme:	Cloning sites used for ORF Shuttling:



\* The last codon before the Stop codon of the ORF




9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn



This product is to be used for laboratory only. Not for diagnostic or therapeutic use. ©2022 OriGene Technologies, Inc., 9620 Medical Center Drive, Ste 200, Rockville, MD 20850, US

```
Gpx1 (NM_008160) Mouse Tagged ORF Clone – MG202058
```

## Plasmid Map:



ACCN:	NM_008160
OTI Disclaimer:	The molecular sequence of this clone aligns with the gene accession number as a point of reference only. However, individual transcript sequences of the same gene can differ through naturally occurring variations (e.g. polymorphisms), each with its own valid existence. This clone is substantially in agreement with the reference, but a complete review of all prevailing variants is recommended prior to use. <u>More info</u> The expression of this clone is not guaranteed due to the nature of selenoproteins.
OTI Annotation:	This clone encodes a selenoprotein containing the rare amino acid selenocysteine (Sec). Sec is encoded by UGA codon, which normally signals translational termination. Expression of this clone is not guaranteed due to the nature of selenoproteins.
Components:	The ORF clone is ion-exchange column purified and shipped in a 2D barcoded Matrix tube containing 10ug of transfection-ready, dried plasmid DNA (reconstitute with 100 ul of water).
Reconstitution Method:	<ol> <li>Centrifuge at 5,000xg for 5min.</li> <li>Carefully open the tube and add 100ul of sterile water to dissolve the DNA.</li> <li>Close the tube and incubate for 10 minutes at room temperature.</li> <li>Briefly vortex the tube and then do a quick spin (less than 5000xg) to concentrate the liquid at the bottom.</li> <li>Store the suspended plasmid at -20°C. The DNA is stable for at least one year from date of shipping when stored at -20°C.</li> </ol>

This product is to be used for laboratory only. Not for diagnostic or therapeutic use. ©2022 OriGene Technologies, Inc., 9620 Medical Center Drive, Ste 200, Rockville, MD 20850, US

	Gpx1 (NM_008160) Mouse Tagged ORF Clone – MG202058
RefSeq:	<u>NM 008160.6, NP 032186.2</u>
RefSeq Size:	848 bp
RefSeq ORF:	606 bp
Locus ID:	14775
UniProt ID:	<u>P11352</u>
Cytogenetics:	9 59.24 cM
Gene Summary:	The protein encoded by this gene belongs to the glutathione peroxidase fa which catalyze the reduction of organic hydroperoxides and hydrogen per glutathione, and thereby protect cells against oxidative damage. Knockout

family, members of eroxide (H2O2) by ut mice lacking this gene are highly sensitive to oxidants, and develop mature cataracts due to damage to the eye lens nucleus. Other studies indicate that H2O2 is also essential for growth-factor mediated signal transduction, mitochondrial function, and maintenance of thiol redox-balance; therefore, by limiting H2O2 accumulation, glutathione peroxidases are also involved in modulating these processes. Several isozymes of this gene family exist in vertebrates, which vary in cellular location and substrate specificity. This isozyme is the most abundant, is ubiquitously expressed and localized in the cytoplasm, and whose preferred substrate is hydrogen peroxide. It is also a selenoprotein, containing the rare amino acid selenocysteine (Sec) at its active site. Sec is encoded by the UGA codon, which normally signals translation termination. The 3' UTRs of selenoprotein mRNAs contain a conserved stem-loop structure, designated the Sec insertion sequence (SECIS) element, that is necessary for the recognition of UGA as a Sec codon, rather than as a stop signal. Alternatively spliced transcript variants have been found for this gene. [provided by RefSeq, Jul 2016]

This product is to be used for laboratory only. Not for diagnostic or therapeutic use. ©2022 OriGene Technologies, Inc., 9620 Medical Center Drive, Ste 200, Rockville, MD 20850, US