

## Product datasheet for **MG202087**

### **Psmb6 (NM\_008946) Mouse Tagged ORF Clone**

#### Product data:

**Product Type:** Expression Plasmids  
**Product Name:** Psmb6 (NM\_008946) Mouse Tagged ORF Clone  
**Tag:** TurboGFP  
**Symbol:** Psmb6  
**Synonyms:** Lmp19; Mpnd  
**Mammalian Cell Selection:** Neomycin  
**Vector:** pCMV6-AC-GFP (PS100010)  
**E. coli Selection:** Ampicillin (100 ug/mL)  
**ORF Nucleotide Sequence:** >MG202087 representing NM\_008946  
 Red=Cloning site Blue=ORF Green=Tags(s)

TTTTGTAATACGACTCACTATAGGGCGGCCGGAATTCGTCGACTGGATCCGGTACCGAGGAGATCTGCC  
 GCC**CGATCGCC**

ATGGCTGTGCAGTTTAAATGGGGCGTGGTTCTAGGAGCGGACTCCAGGACAACCACTGGGTCTACATCG  
 CCAATCGAGTGACTGACAAGCTGACCCCTATCCACGATCACATCTTCTGTCGCCGCTCAGGCTCAGCCGC  
 TGATACCAAGCAGTGGCAGACGCTGTCACTTACCAGCTTGGTTCCACAGTATTGAACTGAACGAGCCT  
 CCACTAGTCCACACAGCCGCCAGTCTCTTAAGGAGATGTGTACCGGTACAGAGAAGATCTGATGGCAG  
 GAATCATCATTGCAGGCTGGGACCTCAAGAAGGAGGGCAGGTGTACTCTGTTCCATGGGGGTATGAT  
 GGTAAAGACAGTCCTTTGCCATCGGAGGCTCCCGGAGCTCGTACATCTATGGCTATGTTGATGCTAGGTAT  
 CGGGAAGGCATGACCAAGGACGAATGTCTGCAGTTCCTGCAATGCTCTCGCTTTGGCCATGGAACGCG  
 ACGGCTCCAGTGGAGGGGTATCCGCTTGGCAGCCATTCAGGAGTCAGGGGTAGAGCGGCAGGTGCTTTT  
 GGGAGACCAAATCCCAAGTTCACCATTGCCACGTTGCCACCTCCC

**ACGCGTACGCGGCCGCTCGAG** - GFP Tag - GTTTAA

**Protein Sequence:** >MG202087 representing NM\_008946  
 Red=Cloning site Green=Tags(s)

MAVQFNGGVVLGADSRRTTGSYIANRVTDKLTPIHDHIFCCRSRSAADTQAVADAVTYQLGFHSIELNEP  
 PLVHTAASLFKEMCYRYREDLMAGIIAGWDPQEGGQVYVPMGMMVVRQSFAIGGSRSSYIYGVDATY  
 REGMTKDECLQFTANALALAMERDGSSSGVIRLAAIQESGVERQVLLGDQIPKFTIATLPPP

**TRTRPLE** - GFP Tag - V

**Restriction Sites:** Sgfl-MluI




[View online »](#)

**Cloning Scheme:**



**Plasmid Map:**



**ACCN:** NM\_008946

**ORF Size:** 606 bp

**OTI Disclaimer:** The molecular sequence of this clone aligns with the gene accession number as a point of reference only. However, individual transcript sequences of the same gene can differ through naturally occurring variations (e.g. polymorphisms), each with its own valid existence. This clone is substantially in agreement with the reference, but a complete review of all prevailing variants is recommended prior to use. [More info](#)

**OTI Annotation:** This clone was engineered to express the complete ORF with an expression tag. Expression varies depending on the nature of the gene.

**Components:** The ORF clone is ion-exchange column purified and shipped in a 2D barcoded Matrix tube containing 10ug of transfection-ready, dried plasmid DNA (reconstitute with 100 ul of water).

**Reconstitution Method:**

1. Centrifuge at 5,000xg for 5min.
2. Carefully open the tube and add 100ul of sterile water to dissolve the DNA.
3. Close the tube and incubate for 10 minutes at room temperature.
4. Briefly vortex the tube and then do a quick spin (less than 5000xg) to concentrate the liquid at the bottom.
5. Store the suspended plasmid at -20°C. The DNA is stable for at least one year from date of shipping when stored at -20°C.

**RefSeq:** [NM\\_008946.3](#)

**RefSeq Size:** 805 bp

**RefSeq ORF:** 717 bp

**Locus ID:** 19175

**UniProt ID:** [Q60692](#)

**Cytogenetics:** 11 42.99 cM

**Gene Summary:** Component of the 20S core proteasome complex involved in the proteolytic degradation of most intracellular proteins. This complex plays numerous essential roles within the cell by associating with different regulatory particles. Associated with two 19S regulatory particles, forms the 26S proteasome and thus participates in the ATP-dependent degradation of ubiquitinated proteins. The 26S proteasome plays a key role in the maintenance of protein homeostasis by removing misfolded or damaged proteins that could impair cellular functions, and by removing proteins whose functions are no longer required. Associated with the PA200 or PA28, the 20S proteasome mediates ubiquitin-independent protein degradation. This type of proteolysis is required in several pathways including spermatogenesis (20S-PA200 complex) or generation of a subset of MHC class I-presented antigenic peptides (20S-PA28 complex). Within the 20S core complex, PSMB6 displays a peptidylglutamyl-hydrolyzing activity also termed postacidic or caspase-like activity, meaning that the peptides bond hydrolysis occurs directly after acidic residues.[UniProtKB/Swiss-Prot Function]