

# Product datasheet for MG202096

### Arl6ip1 (NM\_019419) Mouse Tagged ORF Clone

### **Product data:**

#### OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn


Product Type:	Expression Plasmids
Product Name:	Arl6ip1 (NM_019419) Mouse Tagged ORF Clone
Tag:	TurboGFP
Symbol:	Arl6ip1
Synonyms:	Aip-1; AIP-6; AL022945; Arl6ip; ARMER; AU042858; C85138; mKIAA0069
Mammalian Cell Selection:	Neomycin
Vector:	pCMV6-AC-GFP (PS100010)
E. coli Selection:	Ampicillin (100 ug/mL)
ORF Nucleotide Sequence:	>MG202096 representing NM_019419 Red=Cloning site Blue=ORF Green=Tags(s)
	TTTTGTAATACGACTCACTATAGGGCGGCCGGGAATTCGTCGACTGGATCCGGTACCGAGGAGATCTGCC GCC <mark>GCGATCGC</mark> C
	ATGGCGGAGGGGATAACCGCAGCAGCAACCTGCTGGCCGTGGAGACTGCAAGTCTTGAAGAGCAGCTGC AAGGCTGGGGAGAGGTGATGCTGATGGCTGACAAAGTCCTTCGATGGGAAAGAGCCTGGTTTCCACCTGC CATCATGGGTGTGGTTTCCCTGCTGTTCCTGATTATCTATTATCTCGATCCATCTGTGCTGTCTGGTGTT TCCTGCTTTGTTATGTTTTTGTGCCTGGCTGACTACCTTGTTCCCATTCTAGCACCAAGAATTTTTGGCT CTAATAAATGGACCACTGAACAACAGCAAAGATTTCATGAAATCTGCAGTAATCTAGTAAAAACTCGACG CAGAGCTGTGGGCTGGTGGAAACGCCTCTTTTCCCTAAAGGAAGAAAAGCCTAAAATGTACTTCATGACC ATGATCATTTCTCTTGCTGCGGTGGCTTGGGTGGGACAGCAAGTTCACAACCTGCTTCTACCTAC
Protein Sequence:	ACGCGTACGCGGCCGCTCGAG - GFP Tag - GTTTAA >MG202096 representing NM_019419 Red=Cloning site Green=Tags(s)
	MAEGDNRSSNLLAVETASLEEQLQGWGEVMLMADKVLRWERAWFPPAIMGVVSLLFLIIYYLDPSVLSGV SCFVMFLCLADYLVPILAPRIFGSNKWTTEQQQRFHEICSNLVKTRRRAVGWWKRLFSLKEEKPKMYFMT MIISLAAVAWVGQQVHNLLLTYLIVTFVLLLPGLNQHGIILKYIGMAKREINKLLKQKEKKNE
	TRTRPLE - GFP Tag - V
Restriction Sites:	Sgfl-Mlul




This product is to be used for laboratory only. Not for diagnostic or therapeutic use. ©2022 OriGene Technologies, Inc., 9620 Medical Center Drive, Ste 200, Rockville, MD 20850, US

### **Cloning Scheme:**





#### Plasmid Map:



ACCN:	NM_019419
ORF Size:	609 bp
OTI Disclaimer:	The molecular sequence of this clone aligns with the gene accession number as a point of reference only. However, individual transcript sequences of the same gene can differ through naturally occurring variations (e.g. polymorphisms), each with its own valid existence. This clone is substantially in agreement with the reference, but a complete review of all prevailing variants is recommended prior to use. <u>More info</u>
OTI Annotation:	This clone was engineered to express the complete ORF with an expression tag. Expression varies depending on the nature of the gene.
Components:	The ORF clone is ion-exchange column purified and shipped in a 2D barcoded Matrix tube containing 10ug of transfection-ready, dried plasmid DNA (reconstitute with 100 ul of water).

This product is to be used for laboratory only. Not for diagnostic or therapeutic use. ©2022 OriGene Technologies, Inc., 9620 Medical Center Drive, Ste 200, Rockville, MD 20850, US

## **PORIGENE** Arl6ip1 (NM\_019419) Mouse Tagged ORF Clone – MG202096

Reconstitution Method:	<ol> <li>Centrifuge at 5,000xg for 5min.</li> <li>Carefully open the tube and add 100ul of sterile water to dissolve the DNA.</li> <li>Close the tube and incubate for 10 minutes at room temperature.</li> <li>Briefly vortex the tube and then do a quick spin (less than 5000xg) to concentrate the liquid at the bottom.</li> <li>Store the suspended plasmid at -20°C. The DNA is stable for at least one year from date of shipping when stored at -20°C.</li> </ol>
RefSeq:	<u>NM 019419.2</u>
RefSeq Size:	2080 bp
RefSeq ORF:	612 bp
Locus ID:	54208
UniProt ID:	Q9JKW0
Cytogenetics:	7 F1
Gene Summary:	Positively regulates SLC1A1/EAAC1-mediated glutamate transport by increasing its affinity for glutamate in a PKC activity-dependent manner. Promotes the catalytic efficiency of SLC1A1/EAAC1 probably by reducing its interaction with ARL6IP5, a negative regulator of SLC1A1/EAAC1-mediated glutamate transport (PubMed:18684713). Plays a role in the formation and stabilization of endoplasmic reticulum tubules. Negatively regulates apoptosis, possibly by modulating the activity of caspase-9 (CASP9). Inhibits cleavage of CASP9-dependent substrates and downstream markers of apoptosis but not CASP9 itself. May be involved in protein transport, membrane trafficking, or cell signaling during hematopoietic maturation (By similarity).[UniProtKB/Swiss-Prot Function]