

Product datasheet for MG202330

Chmp2b (NM_026879) Mouse Tagged ORF Clone

Product data:

Product Type: Expression Plasmids

Product Name: Chmp2b (NM_026879) Mouse Tagged ORF Clone

Tag: TurboGFP Symbol: Chmp2b

Synonyms: 1190006E07Rik

Mammalian Cell

Selection:

Neomycin

Vector: pCMV6-AC-GFP (PS100010)

E. coli Selection: Ampicillin (100 ug/mL)

ORF Nucleotide >MG202330 representing NM_026879

Sequence: Red=Cloning site Blue=ORF Green=Tags(s)

TTTTGTAATACGACTCACTATAGGGCGGCCGGGAATTCGTCGACTGGATCCGGTACCGAGGAGATCTGCC

GCCGCGATCGCC

ATGGCCTCCCTCTTCAAGAAGAAGACTGTGGACGATGTCATAAAGGAACAGAATCGAGAGCTGCGTGGCA
CCCAGAGGGCCATCATCAGGGACCGAGCAGCCTTAGAGAAACAGGAGAAACAGCTGGAGTTAGAAATTAA
GAAGATGGCCAAGATTGGTAATAAGGAAGCGTGCAGGGTTTTAGCTAAGCAGCTTGTCCACCTACGGAAA
CAGAAGACAAGAACGTTCGCTGTAAGCTCAAAAGTCACCTCTATGTCCACACAAACGAAAGTGAATT
CCCAGATGAAGATGGCCGGTGCCATGTCTACCACTGCAAAGACAATGCAAGCAGTCAACAAGAAAATGGA
TCCACAGAAGACACTACAAACAATGCAGAAATTTCCAGAAGGAAAACATGAAAATGGAAATGACACCCTGGATGACATCTTTGATGGCTCTGATGATGAAGAAAGCCAGGACATCGTGA
ATCAAGTTCTTGATGAAATTGGGATTGAAATCTCTGGAAAGATGGCCAAAGCTCCATCGGCTGCAAGAAG
CTTACCGTCTGCCTCCACTTCAAAGGCTACAATTTCTGATGAAGAGAAGAATCGAACTCAAACTCAAAGCTTTA

GGGGTAGAT

ACGCGTACGCGGCCGCTCGAG - GFP Tag - GTTTAA



OriGene Technologies, Inc. 9620 Medical Center Drive, Ste 200

CN: techsupport@origene.cn

Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com



Protein Sequence: >MG202330 representing NM_026879

Red=Cloning site Green=Tags(s)


MASLFKKKTVDDVIKEQNRELRGTQRAIIRDRAALEKQEKQLELEIKKMAKIGNKEACRVLAKQLVHLRK QKTRTFAVSSKVTSMSTQTKVMNSQMKMAGAMSTTAKTMQAVNKKMDPQKTLQTMQNFQKENMKMEMTEE MINDTLDDIFDGSDDEEESQDIVNQVLDEIGIEISGKMAKAPSAARSLPSASTSKATISDEEIERQLKAL GVD


TRTRPLE - GFP Tag - V

Restriction Sites:


Sgfl-Mlul

Cloning Scheme:





Plasmid Map:



ACCN: NM_026879

ORF Size: 639 bp

ORIGENE

OTI Disclaimer: The molecular sequence of this clone aligns with the gene accession number as a point of

reference only. However, individual transcript sequences of the same gene can differ through naturally occurring variations (e.g. polymorphisms), each with its own valid existence. This clone is substantially in agreement with the reference, but a complete review of all prevailing

variants is recommended prior to use. More info

OTI Annotation: This clone was engineered to express the complete ORF with an expression tag. Expression

varies depending on the nature of the gene.

Components: The ORF clone is ion-exchange column purified and shipped in a 2D barcoded Matrix tube

containing 10ug of transfection-ready, dried plasmid DNA (reconstitute with 100 ul of water).

Reconstitution Method: 1. Centrifuge at 5,000xg for 5min.

16 C1.3

2. Carefully open the tube and add 100ul of sterile water to dissolve the DNA.

3. Close the tube and incubate for 10 minutes at room temperature.

4. Briefly vortex the tube and then do a quick spin (less than 5000xg) to concentrate the liquid

at the bottom.

5. Store the suspended plasmid at -20°C. The DNA is stable for at least one year from date of

shipping when stored at -20°C.

RefSeq: NM 026879.3

 RefSeq Size:
 1838 bp

 RefSeq ORF:
 642 bp

 Locus ID:
 68942

 UniProt ID:
 Q8BJF9

Cytogenetics:

Gene Summary: Probable core component of the endosomal sorting required for transport complex III

(ESCRT-III) which is involved in multivesicular bodies (MVBs) formation and sorting of endosomal cargo proteins into MVBs. MVBs contain intraluminal vesicles (ILVs) that are generated by invagination and scission from the limiting membrane of the endosome and mostly are delivered to lysosomes enabling degradation of membrane proteins, such as stimulated growth factor receptors, lysosomal enzymes and lipids. The MVB pathway appears to require the sequential function of ESCRT-O, -I,-II and -III complexes. ESCRT-III proteins mostly dissociate from the invaginating membrane before the ILV is released. The ESCRT machinery also functions in topologically equivalent membrane fission events, such as the terminal stages of cytokinesis. ESCRT-III proteins are believed to mediate the necessary vesicle extrusion and/or membrane fission activities, possibly in conjunction with the AAA

ATPase VPS4 (By similarity).[UniProtKB/Swiss-Prot Function]