

Product datasheet for MG202441

Nudt5 (NM_016918) Mouse Tagged ORF Clone

Product data:

Product Type: Expression Plasmids

Product Name: Nudt5 (NM_016918) Mouse Tagged ORF Clone

Tag: TurboGFP

Symbol: Nudt5

Mammalian Cell Neomycin

Selection:

Vector: pCMV6-AC-GFP (PS100010)

E. coli Selection: Ampicillin (100 ug/mL)

ORF Nucleotide >MG202441 representing NM_016918

Sequence: Red=Cloning site Blue=ORF Green=Tags(s)

TTTTGTAATACGACTCACTATAGGGCGGCCGGGAATTCGTCGACTGGATCCGGTACCGAGGAGATCTGCC

GCCGCGATCGCC

ATGGAGACCCGAGAATCCACAGAGTCTTCTCCAGGCAAGCACCTTGTTACCTCAGAGGAGTTGATCTCAG
AAGGAAAATGGGTCAAATTTGAAAAAACAACTTATATGGATCCCACTGGTAAAACCAGAACTTGGGAAAC
AGTGAAACTTACAACCAGGAAGGGAAAATCTGCTGATGCCGTGTCGGTCATACCTGTGCTGCAAAGAACC
CTGCACCATGAGTGCGTCATCCTGGTGAAGCAGTTCCGGCCCCGATGGGCAGCTACTGCCTGGAGTTTC
CAGCAGGGTTCATCGAAGACGGAGAAAACCCAGAGGCGGCTGCTCTTCGGGAGCTGGAGGAAAACTGG
CTACAAAGGTGAAGTTGCGGAATGCTCTCCAGCTGTGGCATGGATCCAGGCTTGTCAAACTGCACCACA
CATGTTGTGACAGTGACCATCAATGGAGATGATGCAGGAAATGTAAGGCCAAAACCCAAACCAGGGGATG
GAGAATTTATGGAAGTGATTTCTTTACCAAAGAATGATCTGCTGACAAGACTTGACGCTTTTGGGAGCAGA
ACAACACCTTACAGTGGATGCCAAGGTCTACGCCTACGGTCTGGACCAACCCCCAACTCGAAGCCA

TTCGAAGTGCCCTTCCTCAAATTT

ACGCGTACGCGGCCGCTCGAG - GFP Tag - GTTTAA

Protein Sequence: >MG202441 representing NM_016918

Red=Cloning site Green=Tags(s)

METRESTESSPGKHLVTSEELISEGKWVKFEKTTYMDPTGKTRTWETVKLTTRKGKSADAVSVIPVLQRT LHHECVILVKQFRPPMGSYCLEFPAGFIEDGENPEAAALRELEEETGYKGEVAECSPAVCMDPGLSNCTT HVVTVTINGDDAGNVRPKPKPGDGEFMEVISLPKNDLLTRLDALGAEQHLTVDAKVYAYGLALKHANSKP

FEVPFLKF

TRTRPLE - GFP Tag - V



OriGene Technologies, Inc. 9620 Medical Center Drive, Ste 200

Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com


CN: techsupport@origene.cn




Restriction Sites:


Sgfl-Mlul

Cloning Scheme:





Plasmid Map:



ACCN: NM 016918

ORF Size: 654 bp

OTI Disclaimer: The molecular sequence of this clone aligns with the gene accession number as a point of

reference only. However, individual transcript sequences of the same gene can differ through naturally occurring variations (e.g. polymorphisms), each with its own valid existence. This clone is substantially in agreement with the reference, but a complete review of all prevailing

variants is recommended prior to use. More info

OTI Annotation: This clone was engineered to express the complete ORF with an expression tag. Expression

varies depending on the nature of the gene.



Components:

The ORF clone is ion-exchange column purified and shipped in a 2D barcoded Matrix tube containing 10ug of transfection-ready, dried plasmid DNA (reconstitute with 100 ul of water).

Reconstitution Method:

- 1. Centrifuge at 5,000xg for 5min.
- 2. Carefully open the tube and add 100ul of sterile water to dissolve the DNA.
- 3. Close the tube and incubate for 10 minutes at room temperature.
- 4. Briefly vortex the tube and then do a quick spin (less than 5000xg) to concentrate the liquid at the bottom.
- 5. Store the suspended plasmid at -20°C. The DNA is stable for at least one year from date of shipping when stored at -20°C.

RefSeq: <u>NM 016918.2</u>, <u>NP 058614.1</u>

RefSeq Size: 1592 bp
RefSeq ORF: 657 bp
Locus ID: 53893
UniProt ID: Q9JKX6
Cytogenetics: 2 A1

Gene Summary: Enzyme that can either act as an ADP-sugar pyrophosphatase in absence of diphosphate or

catalyze the synthesis of ATP in presence of diphosphate (By similarity). In absence of diphosphate, hydrolyzes with similar activities various modified nucleoside diphosphates

such as ADP-ribose, ADP-mannose, ADP-glucose, 8-oxo-GDP and 8-oxo-dGDP (PubMed:10722730). Can also hydrolyze other nucleotide sugars with low activity (PubMed:10722730). In presence of diphosphate, mediates the synthesis of ATP in the nucleus by catalyzing the conversion of ADP-ribose to ATP and ribose 5-phosphate (By

similarity). Nuclear ATP synthesis takes place when dephosphorylated at Thr-44 (By similarity). Nuclear ATP generation is required for extensive chromatin remodeling events that are energy-consuming (By similarity). Does not play a role in U8 snoRNA decapping activity

(PubMed:21070968). Binds U8 snoRNA (PubMed:21070968).[UniProtKB/Swiss-Prot Function]