

Product datasheet for MG202493

Chmp5 (NM_029814) Mouse Tagged ORF Clone

Product data:

OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn


Product Type:	Expression Plasmids
Product Name:	Chmp5 (NM_029814) Mouse Tagged ORF Clone
Tag:	TurboGFP
Symbol:	Chmp5
Synonyms:	2210412K09Rik; AW545668
Mammalian Cell Selection:	Neomycin
Vector:	pCMV6-AC-GFP (PS100010)
E. coli Selection:	Ampicillin (100 ug/mL)
ORF Nucleotide Sequence:	>MG202493 representing NM_029814 Red=Cloning site Blue=ORF Green=Tags(s)
	TTTTGTAATACGACTCACTATAGGGCGGCCGGGAATTCGTCGACTGGATCCGGTACCGAGGAGATCTGCC GCC <mark>GCGATCGC</mark> C
	ATGAACCGATTCTTCGGAAAAGCGAAACCCAAGGCTCCGCCACCTAGCTTGACGGACTGCATTGGGACGG

ATGAALCGATTCTTCGGAAAAGCGAAACCCAAGGCTCCGCCACCTAGCTTGACGGACTGCATTGGGACGG TGGATAGCAGGGCAGAATCCATTGACAAAAAGATTTCCCGGCTGGATGCTGAACTAGTGAAATATAAGGA TCAAATCAAGAAGATGAGAGAGAGGGTCCTGCTAAGAACATGGTCAAACAGAAAGCCCTGAGAGTTTTAAAG CAAAAGCGGATGTATGAGCAACAGCGAGACAACCTGGCCCAACAGTCCTTTAACATGGAGCAAGCTAATT ACACCATCCAGTCACTAAAGGACACCCAAGACCACGGTTGATGCCATGAAGTTGGGAGTAAAGGAAATGAA GAAGGCATATAAGGAAGTAAAAATTGACCAAGATTGAGGACTTACAAGACCAGCTGGAGGATATGATGGAA GATGCAAATGAGATCCAGGAAGCCCTGGGCCGCAGCTACGGCACCCCAGAGTTAGATGAGGACGACCTGG AAGCAGAGTTAGAGTCCAGGAAGCCCTGGGCCGCAGCTACGGCACCCCAGAGTTAGATGAGGGACGACCTGG AAGCAGAGTTAGATGCGCTGGGCGATGAGCTTCTGGCTGATGAAGATAGCTCCTACTTGGATGAGGCAGC TTCCGCTCCTGCAATTCCGGAAGGTGTTCCCACTGACACAAAAACAAGGATGGCGTGCTGGTGGATGAA TTTGGACTGCCGCAGATTCCGCTTCG


ACGCGTACGCGGCCGCTCGAG - GFP Tag - GTTTAA



This product is to be used for laboratory only. Not for diagnostic or therapeutic use. ©2022 OriGene Technologies, Inc., 9620 Medical Center Drive, Ste 200, Rockville, MD 20850, US







ACCN: ORF Size: NM_029814 657 bp

This product is to be used for laboratory only. Not for diagnostic or therapeutic use. ©2022 OriGene Technologies, Inc., 9620 Medical Center Drive, Ste 200, Rockville, MD 20850, US

Chmp5 (NM_029814) Mouse Tagged ORF Clone – MG202493	
OTI Disclaimer:	The molecular sequence of this clone aligns with the gene accession number as a point of reference only. However, individual transcript sequences of the same gene can differ through naturally occurring variations (e.g. polymorphisms), each with its own valid existence. This clone is substantially in agreement with the reference, but a complete review of all prevailing variants is recommended prior to use. <u>More info</u>
OTI Annotation:	This clone was engineered to express the complete ORF with an expression tag. Expression varies depending on the nature of the gene.
Components:	The ORF clone is ion-exchange column purified and shipped in a 2D barcoded Matrix tube containing 10ug of transfection-ready, dried plasmid DNA (reconstitute with 100 ul of water).
Reconstitution Method	 Centrifuge at 5,000xg for 5min. Carefully open the tube and add 100ul of sterile water to dissolve the DNA. Close the tube and incubate for 10 minutes at room temperature. Briefly vortex the tube and then do a quick spin (less than 5000xg) to concentrate the liquid at the bottom. Store the suspended plasmid at -20°C. The DNA is stable for at least one year from date of shipping when stored at -20°C.
RefSeq:	<u>NM 029814.1, NP 084090.1</u>
RefSeq Size:	1372 bp
RefSeq ORF:	660 bp
Locus ID:	76959
UniProt ID:	<u>Q9D7S9</u>
Cytogenetics:	4 A5
Gene Summary:	Probable peripherally associated component of the endosomal sorting required for transport complex III (ESCRT-III) which is involved in multivesicular bodies (MVBs) formation and sorting of endosomal cargo proteins into MVBs. MVBs contain intraluminal vesicles (ILVs) that are generated by invagination and scission from the limiting membrane of the endosome and mostly are delivered to lysosomes enabling degradation of membrane proteins, such as stimulated growth factor receptors, lysosomal enzymes and lipids. The MVB pathway appears to require the sequential function of ESCRT-O, -I,-II and -III complexes. ESCRT-III proteins mostly dissociate from the invaginating membrane before the ILV is released. The ESCRT machinery also functions in topologically equivalent membrane fission events, such as the

vesicle extrusion and/or membrane fission activities, possibly in conjunction with the AAA ATPase VPS4 (By similarity).[UniProtKB/Swiss-Prot Function]

terminal stages of cytokinesis. ESCRT-III proteins are believed to mediate the necessary

This product is to be used for laboratory only. Not for diagnostic or therapeutic use. ©2022 OriGene Technologies, Inc., 9620 Medical Center Drive, Ste 200, Rockville, MD 20850, US