

Product datasheet for **MG203079**

Mdfi (BC006018) Mouse Tagged ORF Clone

Product data:

Product Type:	Expression Plasmids
Product Name:	Mdfi (BC006018) Mouse Tagged ORF Clone
Tag:	TurboGFP
Symbol:	Mdfi
Synonyms:	I-mf; I-mfa
Mammalian Cell Selection:	Neomycin
Vector:	pCMV6-AC-GFP (PS100010)
E. coli Selection:	Ampicillin (100 ug/mL)
ORF Nucleotide Sequence:	>MG203079 representing BC006018 Red=Cloning site Blue=ORF Green=Tags(s)

TTTTGTAATACGACTCACTATAGGGCGGCCGGAATTCGTCGACTGGATCCGGTACCGAGGAGATCTGCC
GCC**CGATCGCC**

ATGTCCCAGGTGAGCGGTGAGTCCCTTCTCGCTGCGACGCGCCTCATGGAGTCCCAGCGCTGCCCTGG
ACCCAGCCAGACCATGTCCCTCCTCCCTGGGCTGGAGGTAGCAAGATCCACTCACCTGTAGAGGCATC
TTCTGAAGAGGGCTTCCCGGAGGAGGCGGCACCCTCCATGCCCATGACAGTGGTCTCCGGGCTCAGCAG
GCTCTGAACAGCATTGACCTCGATGTCCCACAGAAGCTGTGACGTGCCAGCCTCAAGGGAACCCCAAG
GCTGCACCCCACTACTGCCAAATGGCTCCAGCCACGACCACCTCTCAGAACCAGGAGTGCAGGGCATGC
GGGGAACGGTGTCTGGGCGGGTCCAAGGCCACCGGAAGTTGCAGACGCATCCATCTCTGGGCAGCCAG
GCTGGAAGGAAAAGCAGAGGCAGCGCCCGTCCAGCTCACAGGTCCCTCTCCAGGCACAGGAAGATTGCT
GCGTCCACTGCATACTGTCTTATTCTGTGAGTTCCTGACGCTCTGTAACATCCTCCTGGACTGCGC
CACCTGTGGTCTCTGCAGCTCTGAGGACTCCTGCCTCTGCTGCTGCTGTGGTCCGGCAGTGCAGC
GACTGTGACCTGCCCTGCGACCTGGACTGCGGCATCGTGGATGCCTGCTGCGAGTCCGCAGACTGCTTGG
AGATATGCATGGAGTGTGTGGACTCTGTTTCTCCTCC

ACGCGTACGCGGCCGCTCGAG - GFP Tag - GTTTAA



[View online »](#)

Protein Sequence: >MG203079 representing BC006018
 Red=Cloning site Green=Tags(s)

MSQVSGQCPSRCDAPHGVPSAALDPAQTMSELLPGLEVARSTHPVEASSEEGFPPEEAAPSMPHDSGLRAQQ
 ALNSIDLDPVTEAVTCQPQGNPQCTPLLPNGSSHDHLSEPGSAGHAGNGALGGSKAHRKLQTHPSLGSQ
 AGRKSRGSARSASQVPLQAQEDCCVHCILSCLFCEFLTLCNILLDCATCGSCSSEDSCLCCCCGSGECA
 DCDLPCDLDCGIVDACCESADCLEICMECCGLCFSS

TRTRPLE - GFP Tag - V

Restriction Sites: SgfI-MluI

Cloning Scheme:



ACCN: BC006018

ORF Size: 740 bp


OTI Disclaimer: Due to the inherent nature of this plasmid, standard methods to replicate additional amounts of DNA in E. coli are highly likely to result in mutations and/or rearrangements. Therefore, OriGene does not guarantee the capability to replicate this plasmid DNA. Additional amounts of DNA can be purchased from OriGene with batch-specific, full-sequence verification at a reduced cost. Please contact our customer care team at custsupport@origene.com or by calling 301.340.3188 option 3 for pricing and delivery.

The molecular sequence of this clone aligns with the gene accession number as a point of reference only. However, individual transcript sequences of the same gene can differ through naturally occurring variations (e.g. polymorphisms), each with its own valid existence. This clone is substantially in agreement with the reference, but a complete review of all prevailing variants is recommended prior to use. [More info](#)

OTI Annotation: This clone was engineered to express the complete ORF with an expression tag. Expression varies depending on the nature of the gene.

Components:	The ORF clone is ion-exchange column purified and shipped in a 2D barcoded Matrix tube containing 10ug of transfection-ready, dried plasmid DNA (reconstitute with 100 ul of water).
Reconstitution Method:	<ol style="list-style-type: none"> 1. Centrifuge at 5,000xg for 5min. 2. Carefully open the tube and add 100ul of sterile water to dissolve the DNA. 3. Close the tube and incubate for 10 minutes at room temperature. 4. Briefly vortex the tube and then do a quick spin (less than 5000xg) to concentrate the liquid at the bottom. 5. Store the suspended plasmid at -20°C. The DNA is stable for at least one year from date of shipping when stored at -20°C.
RefSeq:	<u>BC006018</u> , <u>AAH06018</u>
RefSeq Size:	1462 bp
RefSeq ORF:	740 bp
Locus ID:	17240
Cytogenetics:	17 23.99 cM
Gene Summary:	Inhibits the transactivation activity of the Myod family of myogenic factors and represses myogenesis. Acts by associating with Myod family members and retaining them in the cytoplasm by masking their nuclear localization signals. Can also interfere with the DNA-binding activity of Myod family members. Plays an important role in trophoblast and chondrogenic differentiation. Regulates the transcriptional activity of TCF7L1/TCF3 by interacting directly with TCF7L1/TCF3 and preventing it from binding DNA. Binds to the axin complex, resulting in an increase in the level of free beta-catenin. Affects axin regulation of the WNT and JNK signaling pathways.[UniProtKB/Swiss-Prot Function]

Product images:



Circular map for MG203079