

Product datasheet for MG203396

Ptpmt1 (NM_025576) Mouse Tagged ORF Clone

TurboGFP Ptpmt1

Neomycin

Expression Plasmids

Product data:

Product Type:

Product Name:

Tag:

Symbol:

Synonyms:

Selection: Vector:

Mammalian Cell

E. coli Selection:

Restriction Sites:

Cloning Scheme:

Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com


CN: techsupport@origene.cn


OriGene Technologies, Inc. 9620 Medical Center Drive, Ste 200

pCMV6-AC-GFP (PS100010) Ampicillin (100 ug/mL) Ascl-Mlul Cloning sites used for ORF Shuttling: Asc I ORF Mlu I

Ptpmt1 (NM 025576) Mouse Tagged ORF Clone

1110001D10Rik; 2810004N20Rik; PLIP





ACCN: **ORF Size:** NM_025576 783 bp



This product is to be used for laboratory only. Not for diagnostic or therapeutic use. ©2024 OriGene Technologies, Inc., 9620 Medical Center Drive, Ste 200, Rockville, MD 20850, US

	1 (NM_025576) Mouse Tagged ORF Clone – MG203396
OTI Disclaimer:	The molecular sequence of this clone aligns with the gene accession number as a point of reference only. However, individual transcript sequences of the same gene can differ through naturally occurring variations (e.g. polymorphisms), each with its own valid existence. This clone is substantially in agreement with the reference, but a complete review of all prevailing variants is recommended prior to use. <u>More info</u>
OTI Annotation:	This clone was engineered to express the complete ORF with an expression tag. Expression varies depending on the nature of the gene.
Components:	The ORF clone is ion-exchange column purified and shipped in a 2D barcoded Matrix tube containing 10ug of transfection-ready, dried plasmid DNA (reconstitute with 100 ul of water).
Reconstitution Method:	 Centrifuge at 5,000xg for 5min. Carefully open the tube and add 100ul of sterile water to dissolve the DNA. Close the tube and incubate for 10 minutes at room temperature. Briefly vortex the tube and then do a quick spin (less than 5000xg) to concentrate the liquid at the bottom. Store the suspended plasmid at -20°C. The DNA is stable for at least one year from date of shipping when stored at -20°C.
RefSeq:	<u>NM 025576.2, NP 079852.1</u>
RefSeq Size:	1322 bp
RefSeq ORF:	786 bp
Locus ID:	66461
UniProt ID:	<u>Q66GT5</u>
Cytogenetics:	2 E1
Gene Summary:	Lipid phosphatase which dephosphorylates phosphatidylglycerophosphate (PGP) to phosphatidylglycerol (PG) (PubMed:21641550, PubMed:21730175). PGP is an essential intermediate in the biosynthetic pathway of cardiolipin, a mitochondrial-specific phospholipid regulating the membrane integrity and activities of the organelle (PubMed:21641550). Has also been shown to display phosphatase activity toward phosphoprotein substrates, specifically mediates dephosphorylation of mitochondrial proteins, thereby playing an essential role in ATP production (By similarity). Has probably a preference for proteins phosphorylated on Ser and/or Thr residues compared to proteins phosphorylated on Tyr residues (By similarity). Probably involved in regulation of insulin secretion in pancreatic beta


cells (By similarity). May prevent intrinsic apoptosis, probably by regulating mitochondrial

membrane integrity (By similarity).[UniProtKB/Swiss-Prot Function]

This product is to be used for laboratory only. Not for diagnostic or therapeutic use. ©2024 OriGene Technologies, Inc., 9620 Medical Center Drive, Ste 200, Rockville, MD 20850, US



Product images:



Circular map for MG203396

This product is to be used for laboratory only. Not for diagnostic or therapeutic use. ©2024 OriGene Technologies, Inc., 9620 Medical Center Drive, Ste 200, Rockville, MD 20850, US