

## **Product datasheet for MG204659**

## Ybx1 (BC031472) Mouse Tagged ORF Clone

**Product data:** 

**Product Type:** Expression Plasmids

**Product Name:** Ybx1 (BC031472) Mouse Tagged ORF Clone

Tag: TurboGFP

Symbol: Ybx1

Synonyms: 1700102N10Rik, MSY1, mYB-1a, EF1A, YB-1, dbpB

Mammalian Cell

Selection:

Neomycin

Vector: pCMV6-AC-GFP (PS100010)

E. coli Selection: Ampicillin (100 ug/mL)

ORF Nucleotide >MG204659 representing BC031472

Sequence: Red=Cloning site Blue=ORF Green=Tags(s)

TTTTGTAATACGACTCACTATAGGGCGGCCGGGAATTCGTCGACTGGATCCGGTACCGAGGAGATCTGCC

GCCGCGATCGCC

ACGCGTACGCGGCCGCTCGAG - GFP Tag - GTTTAA



**OriGene Technologies, Inc.** 9620 Medical Center Drive, Ste 200

CN: techsupport@origene.cn

Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com



**Protein Sequence:** 

>MG204659 representing BC031472 Red=Cloning site Green=Tags(s)

MSSEAETQQPPAAPAAALSAADTKPGSTGSGAGSGGPGGLTSAAPAGGDKKVIATKVLGTVKWFNVRNGY GFINRNDTKEDVFVHQTAIKKNNPRKYLRSVGDGETVEFDVVEGEKGAEAANVTGPGGVPVQGSKYAADR NHYRRYPRRRGPPRNYQQNYQNSESGEKNEGSESAPEGQAQQRRPYRRRRFPPYYMRRPYARRPQYSNPP VQGEVMEGADNQGAGEQGRPVRQNMYRGYRPRFRRGPPRQRQPREDGNEEDKENQGDETQGQQPPQRRYR RNFNYRRRPENPKPQDGKETKAADPPAENSSAPEAEQGGAE


TRTRPLE - GFP Tag - V

**Restriction Sites:** 

Sgfl-Mlul

Cloning Scheme:





**ACCN:** BC031472 **ORF Size:** 968 bp

OTI Disclaimer: Due to the inhere

Due to the inherent nature of this plasmid, standard methods to replicate additional amounts of DNA in E. coli are highly likely to result in mutations and/or rearrangements. Therefore, OriGene does not guarantee the capability to replicate this plasmid DNA. Additional amounts of DNA can be purchased from OriGene with batch-specific, full-sequence verification at a reduced cost. Please contact our customer care team at <a href="mailto:custsupport@origene.com">custsupport@origene.com</a> or by calling 301.340.3188 option 3 for pricing and delivery.

The molecular sequence of this clone aligns with the gene accession number as a point of reference only. However, individual transcript sequences of the same gene can differ through naturally occurring variations (e.g. polymorphisms), each with its own valid existence. This clone is substantially in agreement with the reference, but a complete review of all prevailing variants is recommended prior to use. More info

**OTI Annotation:** 

This clone was engineered to express the complete ORF with an expression tag. Expression varies depending on the nature of the gene.



Components:

The ORF clone is ion-exchange column purified and shipped in a 2D barcoded Matrix tube containing 10ug of transfection-ready, dried plasmid DNA (reconstitute with 100 ul of water).

**Reconstitution Method:** 

- 1. Centrifuge at 5,000xg for 5min.
- 2. Carefully open the tube and add 100ul of sterile water to dissolve the DNA.
- 3. Close the tube and incubate for 10 minutes at room temperature.
- 4. Briefly vortex the tube and then do a quick spin (less than 5000xg) to concentrate the liquid at the bottom.
- 5. Store the suspended plasmid at -20°C. The DNA is stable for at least one year from date of shipping when stored at -20°C.

**RefSeq:** <u>BC031472</u>, <u>AAH31472</u>


RefSeq Size: 1562 bp
RefSeq ORF: 968 bp
Locus ID: 22608
Cytogenetics: 4 D2.1

**Gene Summary:** Mediates pre-mRNA alternative splicing regulation. Component of the CRD-mediated complex that promotes MYC mRNA stability. Binds to splice sites in pre-mRNA and re-

complex that promotes MYC mRNA stability. Binds to splice sites in pre-mRNA and regulates splice site selection. Binds and stabilizes cytoplasmic mRNA. Contributes to the regulation of translation by modulating the interaction between the mRNA and eukaryotic initiation factors. Binds to promoters that contain a Y-box (5'-CTGATTGGCCAA-3'), such as HLA class II genes. Regulates the transcription of numerous genes. Promotes separation of DNA strands that contain mismatches or are modified by cisplatin. Has endonucleolytic activity and can introduce nicks or breaks into double-stranded DNA (in vitro). May play a role in DNA repair. Its transcriptional activity on the multidrug resistance gene MDR1 is enhanced in presence of the APEX1 acetylated form at 'Lys-6' and 'Lys-7'. Binds preferentially to 5'-[CU]CUGCG-3' motified with a similarity of the interaction.

in vitro (By similarity).[UniProtKB/Swiss-Prot Function]

## **Product images:**



Circular map for MG204659