

Product datasheet for **MG204763**

Mlst8 (NM_019988) Mouse Tagged ORF Clone

Product data:

Product Type:	Expression Plasmids
Product Name:	Mlst8 (NM_019988) Mouse Tagged ORF Clone
Tag:	TurboGFP
Symbol:	Mlst8
Synonyms:	0610033N12Rik; AA409454; AI505104; AI851821; Gbl
Mammalian Cell Selection:	Neomycin
Vector:	pCMV6-AC-GFP (PS100010)
E. coli Selection:	Ampicillin (100 ug/mL)
ORF Nucleotide Sequence:	>MG204763 representing NM_019988 Red=Cloning site Blue=ORF Green=Tags(s)

TTTTGTAATACGACTCACTATAGGGCGGCCGGAATTCGTCGACTGGATCCGGTACCGAGGAGATCTGCC
GCC**CGATCGC**C

ATGAATACCACCCAGGCACAGTGGGCAGTGACCCTGTCATCTTAGCAACTGCAGGCTATGACCACCGG
TGCGCTTTTGGCAGGCTCACAGTGAATCTGCACTCGAACAGTGCAGCATCAGGACTCTCAGGTAATGC
ATTGGAGATTACTCCGGACCGAAGCATGATTGCTGCTGCAGGTTACCAACATATCCGCATGTATGATCTC
AACTCCAATAACCCCAACCCCATCATCAGTTATGACGGAGTCAGTAAGAATTGCATCAGTGGGCTTTC
ACGAGGATGGTCGCTGGATGTATACAGGTGGCGAGGACTGCACAGCTCGCATCTGGGACCTCAGGTCCCG
GAATTGCAGTGTGAGCGTATCTTCCAGGTGAACGCACCCATAATTGCGTGTGTCTGCATCCCAACCAG
GCAGAACTCATTGTGGGTGATCAGAGCGGTGCTATCCACATCTGGGACCTGAAGACAGACCACAACGAGC
AGCTGATCCCGAGCCTGAGTCTTCCATCACGTCTGCTCACATCGACCCAGATGCTAGCTACATGGCAGC
CGTCAATAGTGCCGAAACTGCTATGTCTGGAACCTGACAGGGGGCATTGGTGACGATGTGACTCAGCTC
ATCCCTAAGACCAAGATCCCAGCCCATACACGCTATGCCCTGCAATGCCGCTTCAGCCCTGATTCCACGC
TTCTTGCCACCTGTTGAGCTGACCAGACATGTAATCTGGAGGACATCCAATTCTCCCTGATGACAGA
GCTTAGCATCAAGAGTAGTAACCCTGGAGAGTCATCCCGTGGCTGGATGTGGGGCTGTGCCTTCTCAGGG
GATTCCAGTACATTGTCACAGTCTTCTGACAACCTAGCCCGCTCTGGTGTGTAGAGACTGGAGAAA
TCAAGAGAGAGTATGGTGCCATCAGAAAGCTGTCGTCTGCTTGGCCTCAATGACAGTGTGCTGGT

ACGCGTACGCGGCCGCTCGAG - GFP Tag - GTTTAA



[View online »](#)

Protein Sequence: >MG204763 representing NM_019988
 Red=Cloning site Green=Tags(s)

MNTTPGTVGSDPVILATAGYDHTVRFWQAHSGICTRTVQHQSQVNALEITPDRSMIAAAGYQHIRMVYDL
 NSNNPNPIISYDGVSKNIASVGFHEDGRWMTGGEDCTARIWDLRSRNLQCQRIFQVNPAPINCVCLHPNQ
 AELIVGDQSGAIHIWDLKTDHNEQLIPEPESSITSAHIDPDASYMAAVNSAGNCYVWNL TGGIGDDVTQL
 IPKTKIPAHTRYALQCRFSPDSTLLATCSADQTKIWRTSNFSLMTELSIKSSNPGESSRGWMMWGCASFSG
 DSQYIVTASSDNLARLWCVETGEIKREYGGHQKAVVCLAFNDSVLG

TRTRPLE - GFP Tag - V


Restriction Sites: SgfI-MluI

Cloning Scheme:

Cloning sites used for ORF Shuttling:



Plasmid Map:



ACCN: NM_019988

ORF Size: 978 bp

OTI Disclaimer:	The molecular sequence of this clone aligns with the gene accession number as a point of reference only. However, individual transcript sequences of the same gene can differ through naturally occurring variations (e.g. polymorphisms), each with its own valid existence. This clone is substantially in agreement with the reference, but a complete review of all prevailing variants is recommended prior to use. More info
OTI Annotation:	This clone was engineered to express the complete ORF with an expression tag. Expression varies depending on the nature of the gene.
Components:	The ORF clone is ion-exchange column purified and shipped in a 2D barcoded Matrix tube containing 10ug of transfection-ready, dried plasmid DNA (reconstitute with 100 ul of water).
Reconstitution Method:	<ol style="list-style-type: none">1. Centrifuge at 5,000xg for 5min.2. Carefully open the tube and add 100ul of sterile water to dissolve the DNA.3. Close the tube and incubate for 10 minutes at room temperature.4. Briefly vortex the tube and then do a quick spin (less than 5000xg) to concentrate the liquid at the bottom.5. Store the suspended plasmid at -20°C. The DNA is stable for at least one year from date of shipping when stored at -20°C.
RefSeq:	NM_019988.2 , NP_064372.2
RefSeq Size:	3361 bp
RefSeq ORF:	981 bp
Locus ID:	56716
UniProt ID:	Q9DCJ1
Cytogenetics:	17 A3.3

Gene Summary:

Subunit of both mTORC1 and mTORC2, which regulates cell growth and survival in response to nutrient and hormonal signals. mTORC1 is activated in response to growth factors or amino acids. Growth factor-stimulated mTORC1 activation involves a AKT1-mediated phosphorylation of TSC1-TSC2, which leads to the activation of the RHEB GTPase that potentially activates the protein kinase activity of mTORC1. Amino acid-signaling to mTORC1 requires its relocalization to the lysosomes mediated by the Ragulator complex and the Rag GTPases. Activated mTORC1 up-regulates protein synthesis by phosphorylating key regulators of mRNA translation and ribosome synthesis. mTORC1 phosphorylates EIF4EBP1 and releases it from inhibiting the elongation initiation factor 4E (eIF4E). mTORC1 phosphorylates and activates S6K1 at 'Thr-389', which then promotes protein synthesis by phosphorylating PDCD4 and targeting it for degradation. Within mTORC1, LST8 interacts directly with MTOR and enhances its kinase activity. In nutrient-poor conditions, stabilizes the MTOR-RPTOR interaction and favors RPTOR-mediated inhibition of MTOR activity. mTORC2 is also activated by growth factors, but seems to be nutrient-insensitive. mTORC2 seems to function upstream of Rho GTPases to regulate the actin cytoskeleton, probably by activating one or more Rho-type guanine nucleotide exchange factors. mTORC2 promotes the serum-induced formation of stress-fibers or F-actin. mTORC2 plays a critical role in AKT1 'Ser-473' phosphorylation, which may facilitate the phosphorylation of the activation loop of AKT1 on 'Thr-308' by PDK1 which is a prerequisite for full activation. mTORC2 regulates the phosphorylation of SGK1 at 'Ser-422'. mTORC2 also modulates the phosphorylation of PRKCA on 'Ser-657'. [UniProtKB/Swiss-Prot Function]