

Product datasheet for **MG204972**

B3gat3 (NM_024256) Mouse Tagged ORF Clone

Product data:

Product Type: Expression Plasmids
Product Name: B3gat3 (NM_024256) Mouse Tagged ORF Clone
Tag: TurboGFP
Symbol: B3gat3
Synonyms: 2810405M13Rik
Mammalian Cell Selection: Neomycin
Vector: pCMV6-AC-GFP (PS100010)
E. coli Selection: Ampicillin (100 ug/mL)
ORF Nucleotide Sequence: >MG204972 representing NM_024256
Red=Cloning site Blue=ORF Green=Tags(s)

TTTTGTAATACGACTCACTATAGGGCGGCCGGAATTCGTCGACTGGATCCGGTACCGAGGAGATCTGCC
GCC**CGATCGCC**

ATGAAGCTGAAGCTGAAGAACGTGTTTCTTGCCTACTTCCTGGTGTGATCGCCGGCCTCCTCTACGCTC
TGGTGCAGCTCGGCCAGCCTTGCAGCTGCCTCCCTCCGTTTCGAGCTGCAGCTGAGCAGCTTCGGCAGAA
GGACCTGAGGATATCCCAGTTGCAAGCTGATCTCCGTCGCCACCCCTGTCCAGCCCAGCCCCCTGAA
CCTGAGGCCCTGCCTACTATCTATGTATTACCCACCTACGCCAGGCTGGTACAAAAGGCAGAGCTGG
TTCGGCTGTCCCAGACCCTGAGCTTGGTGCCCGTCTACACTGGCTGCTAGTGGAGGACGCTGAGAGCCC
TACCCCGCTGGTCTCGGGGCTGTTGGCCGCTCTGGTCTCCTCTTTACACACCTGGCTGTCTTACCCCC
AAGGCTCAACGGCTTAGGGAAGGTGAGCCAGGCTGGGTCCGGCCCGAGGAGTGGAAACAGCGCAATAAGG
CCCTCGACTGGCTCCGAGGAAAAGGGGGTGTGTTGGGGGGGAGAAGGATCCACCGCCACCAGGGACCCA
AGGAGTCGTGATTTTGTGACGATGACAACACCTACAGCCGGGAGCTCTTTAAGGAGATGCGTTGGACT
CGCGGTGTCTCAGTGTGGCCTGTGGGCTGGTGGGTGGCTGCGATTTGAAGTCTCAGGTACAGGATG
GCCGCTTGTGGGTTCCACACAGCATGGGAACCAACAGGCCCTTCCCTTGGACATGGCGGGATTTGC
GGTTGCCCTGCCCTTGTATTGGCTAAGCCCAATGCCAGTTTGTGCTACTGCACCCCGGGGCCACCTG
GAAAGTAGTCTCCTGAGCCACCTTGTAGATCCCAAGGACCTGGAGCCACGGGCTGCCAATTGTAACAGG
TACTGGTATGGCACACCCGGACAGAGAAACCTAAGATGAAGCAGGAGGAGCAGCTACAACGGCAGGGCCA
GGGCTCAGACCCAGCCATTGAGGTG

ACGCGTACGCGGCCGCTCGAG - GFP Tag - GTTTAA



[View online »](#)

Protein Sequence: >MG204972 representing NM_024256
 Red=Cloning site Green=Tags(s)

MKLKLNKVFLLAYFLVSIAGLLYALVQLGQPCDCLPPLRAAAEQLRQKDLRISQLQADLRRPPVPVPAQPPE
 PEALPTIYVITPTYARLVQKAELVRLSQTLSLVPRLHWLLVEDAESPTPLVSGLLAASGLLFTHLAVLTP
 KAQRLREGEPGWVRPRGVEQRNKALDWLRGKGGAVGGEKDPPTGQGVVYFADDNTYSRELFKEMRWT
 RGVSVWPVGLVGGLRFEGPQVQDGRVVGFTAWEPNRPFLDMAGFAVALPLLLAKPNAQFDATAPRGHL
 ESSLLSHLVDPKDLPEPRAANCTQVLVWHTRTEKPKMKQEEQLQRQGGSDPAIEV

TRTRPLE - GFP Tag - V


Restriction Sites: SgfI-MluI

Cloning Scheme:

Cloning sites used for ORF Shuttling:



Plasmid Map:



ACCN: NM_024256

ORF Size: 1005 bp

OTI Disclaimer:	The molecular sequence of this clone aligns with the gene accession number as a point of reference only. However, individual transcript sequences of the same gene can differ through naturally occurring variations (e.g. polymorphisms), each with its own valid existence. This clone is substantially in agreement with the reference, but a complete review of all prevailing variants is recommended prior to use. More info
OTI Annotation:	This clone was engineered to express the complete ORF with an expression tag. Expression varies depending on the nature of the gene.
Components:	The ORF clone is ion-exchange column purified and shipped in a 2D barcoded Matrix tube containing 10ug of transfection-ready, dried plasmid DNA (reconstitute with 100 ul of water).
Reconstitution Method:	<ol style="list-style-type: none">1. Centrifuge at 5,000xg for 5min.2. Carefully open the tube and add 100ul of sterile water to dissolve the DNA.3. Close the tube and incubate for 10 minutes at room temperature.4. Briefly vortex the tube and then do a quick spin (less than 5000xg) to concentrate the liquid at the bottom.5. Store the suspended plasmid at -20°C. The DNA is stable for at least one year from date of shipping when stored at -20°C.
RefSeq:	NM_024256.2 , NP_077218.1
RefSeq Size:	1599 bp
RefSeq ORF:	1008 bp
Locus ID:	72727
UniProt ID:	P58158
Cytogenetics:	19 A
Gene Summary:	Glycosaminoglycans biosynthesis. Involved in forming the linkage tetrasaccharide present in heparan sulfate and chondroitin sulfate. Transfers a glucuronic acid moiety from the uridine diphosphate-glucuronic acid (UDP-GlcUA) to the common linkage region trisaccharide Gal-beta-1,3-Gal-beta-1,4-Xyl covalently bound to a Ser residue at the glycosaminylglycan attachment site of proteoglycans. Can also play a role in the biosynthesis of I2/HNK-1 carbohydrate epitope on glycoproteins. Stimulates 2-phosphoxylose phosphatase activity of PXYLP1 in presence of uridine diphosphate-glucuronic acid (UDP-GlcUA) during completion of linkage region formation.[UniProtKB/Swiss-Prot Function]