

Product datasheet for **MG205259**

Asna1 (NM_019652) Mouse Tagged ORF Clone

Product data:

Product Type: Expression Plasmids
Product Name: Asna1 (NM_019652) Mouse Tagged ORF Clone
Tag: TurboGFP
Symbol: Asna1
Synonyms: 1810048H22Rik; ArsA
Mammalian Cell Selection: Neomycin
Vector: pCMV6-AC-GFP (PS100010)
E. coli Selection: Ampicillin (100 ug/mL)
ORF Nucleotide Sequence: >MG205259 representing NM_019652
 Red=Cloning site Blue=ORF Green=Tags(s)

TTTTGTAATACGACTCACTATAGGGCGGCCGGAATTCGTCGACTGGATCCGGTACCGAGGAGATCTGCC
 GCC**CGATCGCC**

ATGGCGCGGGGGTGGCCGGGTGGGGGTTGAAGCAGAAGAGTTCGAGGATGCACCTGATGTGGAGCCGC
 TGGAAACCACGCTTAGCAATATCATCGAGCAGCGTAGCCTTAAGTGGATCTTTGTCGGGGCAAGGGTGG
 CGTTGGTAAAGACCACCTGCAGCTGCAGCCTGGCAGTCCAGCTGTCGAAGGGACGTGAGAGTGTCTGATC
 ATTTCCACAGACCAGCTCACAACATCTCAGATGCTTTTGACCAGAAGTCTCCAAGGTGCCTACCAAGG
 TCAAAGGCTATGACAACCTCTTTGCTATGGAGATAGACCCAGCCTGGGGTTCAGAGCTCCCCGACGA
 GTTCTTCGAGGAAGACAACATGCTGAGCATGGGCAAGAAGATGATGCAGGAGGCCATGAGCGCCTCCCT
 GGCATCGATGAAGCCATGAGTTATGCTGAGGTCATGAGGCTGGTAAAAGGCATGAACTTCTCAGTGGTAG
 TGTTTCGACACAGCACCCACCGGCCATACACTCAGGCTCCTGAACTTCCCGACCATCGTGGAGAGGGGCT
 GGGCCGCTGATGCAGATCAAGAACCAGATCAGCCCTTCATCTCACAGATGTGCAACATGCTGGGCCTG
 GGGGACATGAATGCTGACCAACTGGCCTCAAGCTAGAAGAGACGTTGCCTGTATCCGATCTGTTAGCG
 AACAGTTCAAGGACCCTGAACAGACAACGTTCTCTGTGTGTGCATCGCCGAATCTTGTCTTGTACGA
 GACGGAGCGGCTGATCCAGGAGCTGGCTAAGTGAAGATCGACACCCACAACATCATCGTCAACCAGCTT
 GTCTTCCCTGACCCTGAGAAACCCTGCAAGATGTGTGAGGCCCGACACAAGATCCAGGCCAAGTACCTGG
 ACCAGATGGAAGACCTATATGAAGACTTTCACATTGTAAGCTGCCACTGTTACCTCACGAGGTTCCGGG
 AGCCGACAAAGTCAACACCTTCTCTGCCCTCCTCTGGAGCCCTACAAGCCCCCAGCACCCAG

ACGCGTACGCGGCCGCTCGAG - GFP Tag - GTTTAA



[View online »](#)

Protein Sequence: >MG205259 representing NM_019652
 Red=Cloning site Green=Tags(s)


MAAGVAGWGVEAEFEADPDVEPLEPTLSNIEQRSLKWIFVGGKGGVGTTCSCSLAVQLSKGRESVLI
 ISTDPAHNISDAFDQKF SKVPTKVKGYDNL FAMEIDPSLGV AELPDEFF EEDNMLSMGKMMQEAMSAFP
 GIDEAMSYAEVMRLVKGMNFSVVVFD TAPTGH LRLNFP TIVERGLGRLMQIKNQISPFISQMCNMLGL
 GDMNADQLASKLEETLPVIRSVSEQFKDPEQTTFCVCI AEFLSLYETERLIQELAKCKIDTHNIIVNQL
 VFPDPEKPCMKCEARHKIQAKYLDQMEDLYEDFHIVKLP LLPHEVRGADKVNTFSALLLEPYKPPSTQ

TRTRPLE - GFP Tag - V


Restriction Sites: SgfI-MluI

Cloning Scheme:

Cloning sites used for ORF Shuttling:



Plasmid Map:



ACCN: NM_019652

ORF Size: 1044 bp

OTI Disclaimer:	The molecular sequence of this clone aligns with the gene accession number as a point of reference only. However, individual transcript sequences of the same gene can differ through naturally occurring variations (e.g. polymorphisms), each with its own valid existence. This clone is substantially in agreement with the reference, but a complete review of all prevailing variants is recommended prior to use. More info
OTI Annotation:	This clone was engineered to express the complete ORF with an expression tag. Expression varies depending on the nature of the gene.
Components:	The ORF clone is ion-exchange column purified and shipped in a 2D barcoded Matrix tube containing 10ug of transfection-ready, dried plasmid DNA (reconstitute with 100 ul of water).
Reconstitution Method:	<ol style="list-style-type: none">1. Centrifuge at 5,000xg for 5min.2. Carefully open the tube and add 100ul of sterile water to dissolve the DNA.3. Close the tube and incubate for 10 minutes at room temperature.4. Briefly vortex the tube and then do a quick spin (less than 5000xg) to concentrate the liquid at the bottom.5. Store the suspended plasmid at -20°C. The DNA is stable for at least one year from date of shipping when stored at -20°C.
RefSeq:	NM_019652.1 , NP_062626.1
RefSeq Size:	1280 bp
RefSeq ORF:	1047 bp
Locus ID:	56495
UniProt ID:	O54984
Cytogenetics:	8 C3
Gene Summary:	ATPase required for the post-translational delivery of tail-anchored (TA) proteins to the endoplasmic reticulum. Recognizes and selectively binds the transmembrane domain of TA proteins in the cytosol. This complex then targets to the endoplasmic reticulum by membrane-bound receptors, where the tail-anchored protein is released for insertion. This process is regulated by ATP binding and hydrolysis. ATP binding drives the homodimer towards the closed dimer state, facilitating recognition of newly synthesized TA membrane proteins. ATP hydrolysis is required for insertion. Subsequently, the homodimer reverts towards the open dimer state, lowering its affinity for the membrane-bound receptor, and returning it to the cytosol to initiate a new round of targeting.[UniProtKB/Swiss-Prot Function]