

Product datasheet for **MG205350**

Ada (NM_007398) Mouse Tagged ORF Clone

Product data:

Product Type:	Expression Plasmids
Product Name:	Ada (NM_007398) Mouse Tagged ORF Clone
Tag:	TurboGFP
Symbol:	Ada
Mammalian Cell Selection:	Neomycin
Vector:	pCMV6-AC-GFP (PS100010)
E. coli Selection:	Ampicillin (100 ug/mL)
Restriction Sites:	SgfI-MluI
Cloning Scheme:	

Cloning sites used for ORF Shuttling:



EcoRI BamHI KpnI RBS Kozac Consensus SgfI AscI
 CTATAGGGCGGCCGGGAATTCGTCGACTGGATCCGGTACCGAGGAGATCTGCCGCCGCGATCGCCGCGCGCCAGATCT
 HindIII NheI RsrII MluI NotI XhoI GFP Tag
 CAAGCTTAAGTACTAGTAGCGGACCG ACG CGT ACG CGG CCG CTC GAG ATG GAG AGC GAC --- --- ---
 T R T R P L E M E S D - - -
 PmeI FseI
 --- GAA GAA AGA GTT TAA ACGGCCGCCGCGGAGCT
 - - E E R V stop

ACCN: NM_007398


ORF Size: 1698 bp



[View online »](#)

OTI Disclaimer:	The molecular sequence of this clone aligns with the gene accession number as a point of reference only. However, individual transcript sequences of the same gene can differ through naturally occurring variations (e.g. polymorphisms), each with its own valid existence. This clone is substantially in agreement with the reference, but a complete review of all prevailing variants is recommended prior to use. More info
OTI Annotation:	This clone was engineered to express the complete ORF with an expression tag. Expression varies depending on the nature of the gene.
Components:	The ORF clone is ion-exchange column purified and shipped in a 2D barcoded Matrix tube containing 10ug of transfection-ready, dried plasmid DNA (reconstitute with 100 ul of water).
Reconstitution Method:	<ol style="list-style-type: none"> 1. Centrifuge at 5,000xg for 5min. 2. Carefully open the tube and add 100ul of sterile water to dissolve the DNA. 3. Close the tube and incubate for 10 minutes at room temperature. 4. Briefly vortex the tube and then do a quick spin (less than 5000xg) to concentrate the liquid at the bottom. 5. Store the suspended plasmid at -20°C. The DNA is stable for at least one year from date of shipping when stored at -20°C.
Note:	Plasmids are not sterile. For experiments where strict sterility is required, filtration with 0.22um filter is required.
RefSeq:	NM_007398.2
RefSeq Size:	1627 bp
RefSeq ORF:	1059 bp
Locus ID:	11486
UniProt ID:	P03958
Cytogenetics:	2 H3
Gene Summary:	Catalyzes the hydrolytic deamination of adenosine and 2-deoxyadenosine (PubMed:9272950). Plays an important role in purine metabolism and in adenosine homeostasis (PubMed:9272950, PubMed:10720488). Modulates signaling by extracellular adenosine, and so contributes indirectly to cellular signaling events (PubMed:11435465). Acts as a positive regulator of T-cell coactivation, by binding DPP4. Its interaction with DPP4 regulates lymphocyte-epithelial cell adhesion (By similarity). Enhances dendritic cell immunogenicity by affecting dendritic cell costimulatory molecule expression and cytokines and chemokines secretion (By similarity). Enhances CD4+ T-cell differentiation and proliferation (By similarity). Acts as a positive modulator of adenosine receptors ADORA1 and ADORA2A, by enhancing their ligand affinity via conformational change (By similarity). Stimulates plasminogen activation (By similarity). Plays a role in male fertility (By similarity). Plays a protective role in early postimplantation embryonic development (PubMed:9272950).[UniProtKB/Swiss-Prot Function]

Product images:



Circular map for MG205350