

Product datasheet for **MG20555**

Morf4l1 (NM_001039147) Mouse Tagged ORF Clone

Product data:

Product Type: Expression Plasmids
Product Name: Morf4l1 (NM_001039147) Mouse Tagged ORF Clone
Tag: TurboGFP
Symbol: Morf4l1
Synonyms: mKIAA4002; MORFRG15; MRG15; TEG-189; Tex189
Mammalian Cell Selection: Neomycin
Vector: pCMV6-AC-GFP (PS100010)
E. coli Selection: Ampicillin (100 ug/mL)
ORF Nucleotide Sequence: >MG20555 representing NM_001039147
 Red=Cloning site Blue=ORF Green=Tags(s)

TTTTGTAATACGACTCACTATAGGGCGCCGGGAATTCGTCGACTGGATCCGGTACCGAGGAGATCTGCC
 GCC**CGATCGCC**

ATGGCGCCCAAGCAGGACCCTAAGCCGAAATTCAGGAGGGCGAGCGAGTGCTGTGCTTTCATGGGCCTC
 TTCTCTATGAAGCAAAGTGTGTAAGGTTGCCATAAAGGACAAACAAGTAAATACTTCATCCATTACAG
 TGGCTGGAATAAAAAAGTGTGTGAGGCCAGGCGCTCTGAAAAATCTTGAAGACACGTGAGGATATT
 GTAGCCCTTTTCTGTTCCTGAAGGAGCTCCCTCAGTACACCACCCCTCTGACCTCTAGTTGGGATG
 AATGGGTGCCAGAAAGCAGAGTACTCAAATACGTGGACACCAATTTGCAGAAACACGGAACCTCAAAA
 GGCCAATCAGGAACAATATGCAGAGGGCAAGATGAGAGGGGCTGCTCCGGGGAAGAAGACATCCGGCCTG
 CAACAGAAAAATGTCGAAGTAAAAAAGAAACAAGCAGAAAAACACCTGGAAATGGAGATGGTGGCA
 GTACCAGTGAACACCTCAGCCTCTAGGAAGAAGAGAGCCCGGTAGATCCTACCGTTGAAAATGAAGA
 AACCTTCATGAACAGAGTTGAAGTTAAAGTGAAGATCCCTGAAGAGCTGAAACCTGGCTTGTGGATGAC
 TGGGACTTGATCACCAGACAGAAGCAGCTTTTTATCTTCTGCCAAGAAGAATGTGGATTCCATTTGG
 AGGATTATGCAAATTATAAGAAGTCTCGAGGAAATACAGATAATAAGAGTATGCTGTTAATGAGGTGGT
 GGCAGGCATAAAGGAGTACTTCAATGTGATGTTGGGCACTCAGCTCCTCTACAAGTTTGAGAGACCACAG
 TACGCTGAAATCCTCGCCGACCACCCGGATGCGCCCATGTCACAGGTGTACGGAGCGCCACATTTGCTGA
 GATTATTTGTGCGAATTGGAGCGATGTTGGCCTACACGCCTCTAGATGAGAAAAGCCTTGCTTTATTACT
 GAACTATCTACATGATTTTCTCAAGTACCTGGCGAAGAATTCTGCAACCTTGTTCACTGCCAGTGATTAT
 GAAGTGGCCCTCTGAGTACCACCGAAAGCCGTG

ACCGTACGCGGCCGCTCGAG - GFP Tag - GTTTAA



[View online »](#)

Protein Sequence: >MG205555 representing NM_001039147
 Red=Cloning site Green=Tags(s)

MAPKQDPKPKFQEGERVLCFHGPLL YEAKCVKVAIKDKQVKYFIHYSGWNKKS AVRPRRSEKSLKTREDI
 VALFPVPEGAPSVHHPLL TSSWDEWVPE SRVLKYVDNLQKQRELQKANQEYAEKMRGAAPGKKTSGL
 QQKNVEVKTKKNKQKTPGNGDGGSTSETPQPPRKKRVRDPTVENEETFMRNVEVKVKIPEELKPWLVD
 WDLITRQKQLFYLPAAKNVDSILEDYANYKKS RGN TDNKEYAVNEVVAGIKEYFNVM LGTQLLYKFERPQ
 YAEILADHPDAPMSQVYGAPHLRL FVRIGAMLAYTPLDEKSLALLLNLYLHDFLKYLAKNSATLFSASD
 YVAPPEYHRKAV

TRTRPLE - GFP Tag - V


Restriction Sites:

SgfI-MluI

Cloning Scheme:



Plasmid Map:



ACCN:

NM_001039147

ORF Size:	1086 bp
OTI Disclaimer:	The molecular sequence of this clone aligns with the gene accession number as a point of reference only. However, individual transcript sequences of the same gene can differ through naturally occurring variations (e.g. polymorphisms), each with its own valid existence. This clone is substantially in agreement with the reference, but a complete review of all prevailing variants is recommended prior to use. More info
OTI Annotation:	This clone was engineered to express the complete ORF with an expression tag. Expression varies depending on the nature of the gene.
Components:	The ORF clone is ion-exchange column purified and shipped in a 2D barcoded Matrix tube containing 10ug of transfection-ready, dried plasmid DNA (reconstitute with 100 ul of water).
Reconstitution Method:	<ol style="list-style-type: none">1. Centrifuge at 5,000xg for 5min.2. Carefully open the tube and add 100ul of sterile water to dissolve the DNA.3. Close the tube and incubate for 10 minutes at room temperature.4. Briefly vortex the tube and then do a quick spin (less than 5000xg) to concentrate the liquid at the bottom.5. Store the suspended plasmid at -20°C. The DNA is stable for at least one year from date of shipping when stored at -20°C.
RefSeq:	NM_001039147.2 , NP_001034236.1
RefSeq Size:	1927 bp
RefSeq ORF:	1089 bp
Locus ID:	21761
UniProt ID:	P60762
Cytogenetics:	9 E3.1
Gene Summary:	Component of the NuA4 histone acetyltransferase complex which is involved in transcriptional activation of select genes principally by acetylation of nucleosomal histones H4 and H2A. This modification may both alter nucleosome - DNA interactions and promote interaction of the modified histones with other proteins which positively regulate transcription. This complex may be required for the activation of transcriptional programs associated with oncogene and proto-oncogene mediated growth induction, tumor suppressor mediated growth arrest and replicative senescence, apoptosis, and DNA repair. The NuA4 complex ATPase and helicase activities seem to be, at least in part, contributed by the association of RUVBL1 and RUVBL2 with EP400. NuA4 may also play a direct role in DNA repair when directly recruited to sites of DNA damage. Also component of the mSin3A complex which acts to repress transcription by deacetylation of nucleosomal histones. Required for homologous recombination repair (HRR) and resistance to mitomycin C (MMC). Involved in the localization of PALB2, BRCA2 and RAD51, but not BRCA1, to DNA-damage foci (By similarity).[UniProtKB/Swiss-Prot Function]