OriGene Technologies, Inc.
9620 Medical Center Drive, Ste 200
Rockville, MD 20850, US
Phone: +1-888-267-4436
https://www.origene.com
techsupport@origene.com
EU: info-de@origene.com
CN: techsupport@origene.cn

Product datasheet for MG206229

Ercc8 (NM_028042) Mouse Tagged ORF Clone

Product data:

Product Type:	Expression Plasmids
Product Name:	Ercc8 (NM_028042) Mouse Tagged ORF Clone
Tag:	TurboGFP
Symbol:	Ercc8
Synonyms:	2410022P04Rik; 2810431L23Rik; 4631412O06Rik; B130065P18Rik; Ckn1; Csa
Mammalian Cell Selection:	Neomycin
Vector:	pCMV6-AC-GFP (PS100010)
E. coli Selection:	Ampicillin (100 ug/mL)

ORF Nucleotide Sequence:

>MG206229 representing NM_028042
Red=Cloning site Blue=ORF Green=Tags(s)

TTTTGTAATACGACTCACTATAGGGCGGCCGGGAATTCGTCGACTGGATCCGGTACCGAGGAGATCTGCC
GCCGCGATCGCC

ATGTTGAGCTTTTTGTCCGCTCGACAGTCGGGTTTGGAGGACCCTCTTCGCTTGCGTAGAGCGCAGTCCA
CACGGAGGGTCCTGGGGCTGGAGTTAAACAAAGACAGGGATGTGGAAAGAATCCATGGCAGTGGGGTTAA
CACCCTCGACATTGAGCCCGTTGAAGGAAGATACATGTTGTCAGGTGGCTCCGATGGCGTGGTTGTACTC
TATGACCTTGAGAACGCCAGCAGACAGCCCCATTACACATGTAAAGCAGTGTGTTCCATTGGCAGAAGCC
ATCCCGATGTTCACAAATACAGCGTGGAGACTGTTCAGTGGTATCCTCATGACACTGGCATGTTCACATC
CAGCTCATTTGATAAAACTCTGAAAGTGTGGGATACAAACACATTGCAGGCTGCAGATGTGTTTAATTTT
GAGGAAACAGTTTACAGTCATCACATGTCCCCAGCAGCCACCAAGCACTGTCTGGTAGCAGTTGGAACAA
GAGGACCCAAGGTACAACTTTGTGACTTAAAGTCTGGATCCTGTTCTCACATTCTACAGGGTCACAGACA
GGAAATCTTGGCAGTTTCCTGGTCACCACGCCATGACTATATCTTGGCAACAGCAAGTGCTGACAGTAGA
GTAAAATTATGGGATGTGAGAAGAGCGTCAGGATGCTTGCTTACTCTTGACCAGCATAATGGGAAAAAGT
CACAAGCTGCTGAATCAGCAAACACTGCTCACAATGGGAAAGTTAACGGCTTATGTTTTACAAGTGACGG
CCTTCACCTGCTCACCATTGGCACAGACAACCGAATGCGCCTCTGGAATAGCTCCAGCGGGGACAACACC
CTGGTGAACTATGGGAAGGTGTGTAATGACAGCAGGAAAGGGCTGCAGTTCGCTGTCTCCTGTGGCTGCA
GCTCAGAGTTTGTGTTTGTTCCCCACGGCAGCACCATCGCTGTGTACGCAGTCCACTCGGGAGAGCGGCT
CGCCATGCTCAAGGGACATTACAAAAGCGTCGACTGCTGTGTGTTCCAGCCTAATTTCCAGGAACTTTAC
AGTGGAAGCAGGGACTGCAATATTCTTGCTTGGGTACCACCTTCCTATGAGCCAGTTCCTGATGATGACG
ATGAAGCTCCAGCCAAATCCCAGTTAAACCCAGCATTCGCGGACGCCTGGAGCAGCAGTGACGAGGATGG
G

ACGCGTACGCGGCCGCTCGAG - GFP Tag - GTTTAA

View online »

Protein Sequence:

>MG206229 representing NM_028042
Red=Cloning site Green=Tags(s)

MLSFLSARQSGLEDPLRLRRAQSTRRVLGLELNKDRDVERIHGSGVNTLDIEPVEGRYMLSGGSDGVVVL
YDLENASRQPHYTCKAVCSIGRSHPDVHKYSVETVQWYPHDTGMFTSSSFDKTLKVWDTNTLQAADVFNF
EETVYSHHMSPAATKHCLVAVGTRGPKVQLCDLKSGSCSHILQGHRQEILAVSWSPRHDYILATASADSR
VKLWDVRRASGCLLTLDQHNGKKSQAAESANTAHNGKVNGLCFTSDGLHLLTIGTDNRMRLWNSSSGDNT
LVNYGKVCNDSRKGLQFAVSCGCSSEFVFVPHGSTIAVYAVHSGERLAMLKGHYKSVDCCVFQPNFQELY
SGSRDCNILAWVPPSYEPVPDDDDEAPAKSQLNPAFADAWSSSDEDG

TRTRPLE – GFP Tag – V

Restriction Sites: SgfI-MluI

Cloning Scheme:



Plasmid Map:



ACCN: NM_028042

ORF Size:	1191 bp
OTI Disclaimer:	The molecular sequence of this clone aligns with the gene accession number as a point of reference only. However, individual transcript sequences of the same gene can differ through naturally occurring variations (e.g. polymorphisms), each with its own valid existence. This clone is substantially in agreement with the reference, but a complete review of all prevailing variants is recommended prior to use. More info
OTI Annotation:	This clone was engineered to express the complete ORF with an expression tag. Expression varies depending on the nature of the gene.
Components:	The ORF clone is ion-exchange column purified and shipped in a 2D barcoded Matrix tube containing 10ug of transfection-ready, dried plasmid DNA (reconstitute with 100 ul of water).
Reconstitution Method:	1. Centrifuge at 5,000xg for 5min. 2. Carefully open the tube and add 100ul of sterile water to dissolve the DNA. 3. Close the tube and incubate for 10 minutes at room temperature. 4. Briefly vortex the tube and then do a quick spin (less than 5000xg) to concentrate the liquid at the bottom. 5. Store the suspended plasmid at -20°C. The DNA is stable for at least one year from date of shipping when stored at -20°C.
RefSeq:	NM_028042.2, NP_082318.2
RefSeq Size:	2108 bp
RefSeq ORF:	1194 bp
Locus ID:	71991
UniProt ID:	Q8CFD5
Cytogenetics:	13 D2.1
Gene Summary:	Involved in transcription-coupled nucleotide excision repair. It is required for the recruitment of XAB2, HMGN1 and TCEA1/TFIIS to a transcription-coupled repair complex which removes RNA polymerase II-blocking lesions from the transcribed strand of active genes. It is the substrate-recognition component of the CSA complex (DCX(ERCC8) complex) which promotes the ubiquitination and subsequent proteasomal degradation of ERCC6 in a UV-dependent manner; ERCC6 degradation is essential for the recovery of RNA synthesis after transcription-coupled repair.[UniProtKB/Swiss-Prot Function]