

Product datasheet for **MG206260**

F2r11 (NM_007974) Mouse Tagged ORF Clone

Product data:

Product Type: Expression Plasmids
Product Name: F2r11 (NM_007974) Mouse Tagged ORF Clone
Tag: TurboGFP
Symbol: F2r11
Synonyms: Gpcr11; PAR-2; Par2
Mammalian Cell Selection: Neomycin
Vector: pCMV6-AC-GFP (PS100010)
E. coli Selection: Ampicillin (100 ug/mL)
ORF Nucleotide Sequence: >MG206260 representing NM_007974
 Red=Cloning site Blue=ORF Green=Tags(s)

TTTTGTAATACGACTCACTATAGGGCGGCCGGAATTCGTCGACTGGATCCGGTACCGAGGAGATCTGCC
 GCC**CGATCGCC**

ATGCGAAGTCTCAGCCTGGCGTGGCTGCTGGGAGGTATCACCTTCTGGCGGCCTCGGTCTCCTGCAGCC
 GGACCGAGAACCTTGACCCGGGACGCAACAACAGTAAAGGAAGAAGTCTTATTGGCAGATTAGAAACCCA
 GCCTCCAATCACTGGGAAAGGGTTCCGGTAGAACAGGCTTTCCATCGATGAGTTCTCTCGTCCATC
 CTCACCGGAAGCTGACCACGGTCTTCTCCGGTCGTCTACATTATTGTGTTTGTATTGGTTTGGCCA
 GTAATGGCATGGCCCTCTGGATTTCTTTCCGAACGAAGAAGAACACCCCGCGTGATTTACATGGC
 CAACCTGGCCTTGCCGACCTCCTCTGTGCATCTGGTTCCCCCTGAAGATCTCCTACCACCTACATGGC
 AACAACTGGGTCTACGGGAGGCCCTGTGCAAGGTGCTCATTGGCTTTTTCTATGGTAACATGTATTGCT
 CCATCCTTTCATGACCTGCCTCAGCGTGCAGAGGTAAGTGGTGCATCGTGAACCCATGGGACACCCAG
 GAAGAAGGCAAACATCGCCGTTGGCGTCTCCTTGGCAATCTGGCTCCTGATTTTTCTGGTCACCATCCCT
 TTGTATGTCATGAAGCAGACCATCTACATTCCAGCATTGAACATCACCACCTGTCACGATGTGCTGCCTG
 AGGAGGTATTGGTGGGGACATGTTCAATTACTTCTCTACTGGCCATTGGAGTCTTCTGTTCCTGGC
 CCTCCTTACTGCATCTGCCTACGTGCTCATGATCAAGACGCTCCGCTCTTCTGCTATGGATGAACACTCA
 GAGAAGAAAAGGCAGAGGGCTATCCGACTCATCACCGTGTGGCCATGTACTTCATCTGCTTTGCTC
 CTAGCAACCTTCTGCTCGTAGTGCAATTATTTCTAATCAAACCCAGAGGCAGAGCCACGCTACGCCCT
 CTACCTTGTGCGCCTCTGCCTGTGACCCCTCAACAGCTGCATAGACCCCTTTGTCTATTACTTTGTCTCA
 AAAGATTTAGGGATCACGCCAGAAACGCGCTCCTCTGCCAAGTGTCCGCACTGTGAATCGCATGCAAA
 TCTCGCTCAGCTCCAACAAGTTCTCCAGGAAGTCCGGCTCCTACTCTTCAAGCTCAACCAGTGTTAAAC
 CTCCTAC

ACGCGTACGCGGCCGCTCGAG - GFP Tag - GTTTAA



[View online >](#)

Protein Sequence: >MG206260 representing NM_007974
 Red=Cloning site Green=Tags(s)

MRSLSLAWLLGGITLLAASVSCSRTENLAPGRNNSKGRSLIGRLETQPPITGKGVPEVPGFSIDFSASIL
 LTGKLTTFVFLPVVYIIVFVIGLPSNGMALWIFLFRTKKKHPAVIYMANLALADLLSVIWFPLKISYHLHG
 NNWVYGEALCKVLIGFFYGNMYCSILFMTCLSVQRYWVI VNPMGHPRKKANIAVGVSLAIWLLIFLVTIP
 LYVMKQTIYIPALNITTTCHDVLPEEVLVGDMFNFLSLAIGVFLFPALLTASAYVLMIKTLRSSAMDEHS
 EKRRQRAIRLIITVLMYFICFAPSNNLLVVHYFLIKTQRQSHVYALYLVALCLSTLNSCIDPFVYVYFVS
 KDFRDHARNALLCRSVRTVNRMQISLSSNKFSRKSGSYSSSSTS VKTSY

TRTRPLE - GFP Tag - V

Restriction Sites: SgfI-MluI

Cloning Scheme:



ACCN: NM_007974

ORF Size: 1197 bp

OTI Disclaimer: Due to the inherent nature of this plasmid, standard methods to replicate additional amounts of DNA in E. coli are highly likely to result in mutations and/or rearrangements. Therefore, OriGene does not guarantee the capability to replicate this plasmid DNA. Additional amounts of DNA can be purchased from OriGene with batch-specific, full-sequence verification at a reduced cost. Please contact our customer care team at custsupport@origene.com or by calling 301.340.3188 option 3 for pricing and delivery.

The molecular sequence of this clone aligns with the gene accession number as a point of reference only. However, individual transcript sequences of the same gene can differ through naturally occurring variations (e.g. polymorphisms), each with its own valid existence. This clone is substantially in agreement with the reference, but a complete review of all prevailing variants is recommended prior to use. [More info](#)


OTI Annotation: This clone was engineered to express the complete ORF with an expression tag. Expression varies depending on the nature of the gene.

Components:	The ORF clone is ion-exchange column purified and shipped in a 2D barcoded Matrix tube containing 10ug of transfection-ready, dried plasmid DNA (reconstitute with 100 ul of water).
Reconstitution Method:	<ol style="list-style-type: none">1. Centrifuge at 5,000xg for 5min.2. Carefully open the tube and add 100ul of sterile water to dissolve the DNA.3. Close the tube and incubate for 10 minutes at room temperature.4. Briefly vortex the tube and then do a quick spin (less than 5000xg) to concentrate the liquid at the bottom.5. Store the suspended plasmid at -20°C. The DNA is stable for at least one year from date of shipping when stored at -20°C.
RefSeq:	<u>NM_007974.4, NP_032000.3</u>
RefSeq Size:	2773 bp
RefSeq ORF:	1200 bp
Locus ID:	14063
UniProt ID:	<u>P55086</u>
Cytogenetics:	13 49.53 cM

Gene Summary:

Receptor for trypsin and trypsin-like enzymes coupled to G proteins. Its function is mediated through the activation of several signaling pathways including phospholipase C (PLC), intracellular calcium, mitogen-activated protein kinase (MAPK), I-kappaB kinase/NF-kappaB and Rho. Can also be transactivated by cleaved F2r/Par1. Involved in modulation of inflammatory responses and regulation of innate and adaptive immunity, and acts as a sensor for proteolytic enzymes generated during infection. Generally is promoting inflammation. Can signal synergistically with Tlr4 and probably Tlr2 in inflammatory responses and modulates Tlr3 signaling. Has a protective role in establishing the endothelial barrier; the activity involves coagulation factor X. Regulates endothelial cell barrier integrity during neutrophil extravasation, probably following proteolytic cleavage by PRTN3 (By similarity). Proposed to have a bronchoprotective role in airway epithelium, but also shown to compromise the airway epithelial barrier by interrupting E-cadherin adhesion. Involved in the regulation of vascular tone; activation results in hypotension presumably mediated by vasodilation. Associates with a subset of G proteins alpha subunits such as GNAQ, GNA11, GNA14, GNA12 and GNA13, but probably not with G(o) alpha, G(i) subunit alpha-1 and G(i) subunit alpha-2. Believed to be a class B receptor which internalizes as a complex with arrestin and traffic with it to endosomal vesicles, presumably as desensitized receptor, for extended periods of time. Mediates inhibition of TNF-alpha stimulated JNK phosphorylation via coupling to GNAQ and GNA11; the function involves dissociation of Ripk1 and Tradd from Tnfr1. Mediates phosphorylation of nuclear factor NF-kappa-B RELA subunit at 'Ser-536'; the function involves Ikbkb and is predominantly independent of G proteins. Involved in cellular migration. Involved in cytoskeletal rearrangement and chemotaxis through beta-arrestin-promoted scaffolds; the function is independent of GNAQ and GNA11 and involves promotion of cofilin dephosphorylation and actin filament severing. Induces redistribution of Cops5 from the plasma membrane to the cytosol and activation of the JNK cascade is mediated by Cops5. Involved in the recruitment of leukocytes to the sites of inflammation and is the major PAR receptor capable of modulating eosinophil function such as proinflammatory cytokine secretion, superoxide production and degranulation. During inflammation promotes dendritic cell maturation, trafficking to the lymph nodes and subsequent T-cell activation. Involved in antimicrobial response of innate immune cells; activation enhances phagocytosis of Gram-positive and killing of Gram-negative bacteria. Acts synergistically with interferon-gamma in enhancing antiviral responses (By similarity). [UniProtKB/Swiss-Prot Function]

Product images:



Circular map for MG206260