

Product datasheet for **MG206306**

Ager (NM_007425) Mouse Tagged ORF Clone

Product data:

Product Type: Expression Plasmids
Product Name: Ager (NM_007425) Mouse Tagged ORF Clone
Tag: TurboGFP
Symbol: Ager
Synonyms: RAGE
Mammalian Cell Selection: Neomycin
Vector: pCMV6-AC-GFP (PS100010)
E. coli Selection: Ampicillin (100 ug/mL)
ORF Nucleotide Sequence: >MG206306 representing NM_007425
 Red=Cloning site Blue=ORF Green=Tags(s)

TTTTGTAATACGACTCACTATAGGGCGGCCGGAATTCGTCGACTGGATCCGGTACCGAGGAGATCTGCC
 GCC**CGATCGCC**

ATGCCAGCGGGACAGCAGCTAGAGCCTGGGTGCTGGTTCTTGCTCTATGGGGAGCTGTAGCTGGTGGTC
 AGAACATCACAGCCCGATTGGAGAGCCACTTGTGCTAAGCTGTAAGGGGCCCCAAGAAGCCGCCCA
 GCAGCTAGAATGGAACTGAACACAGGAAGAAGTGAAGCTTGAAGGTCTCTCTCCCCAGGGAGGCC
 TGGGACAGCGTGGCTCGAATCTCCCAATGGTCCCTCCTCCTCCAGCCACTGGAATTGTCGATGAGG
 GGACTTTCGGGTGTCGGGCAACTAACAGGCGAGGGAAGGAGTCAAGTCCAACCTACCGAGTCCGAGTCTA
 CCAGATTCCTGGGAAGCCAGAAATTGTGGATCCTGCCTCTGAACCTCACAGCCAGTGTCCCTAATAAGGTG
 GGGACATGTGTCTGAGGGAAGCTACCTGCAGGGACCCTTAGCTGGCACTTAGATGGGAACTTCTGA
 TTCCCCGATGGCAAAGAACTCGTGAAGGAAGAGACCAGGAGACACCCTGAGACGGGACTCTTTACACT
 GCGGTCAGAGCTGACAGTATCCCCACCAAGGAGGAACCCATCCTACCTTCTCCTGCAGTTTCAGCCTG
 GGCCTTCCCCGGCGCAGACCCCTGAACACAGCCCCATCCAACCTCCGAGTCAGGGAGCCTGGGCCTCCAG
 AGGGCATTAGCTGTTGGTTGAGCCTGAAGGTGGAATAGTCGCTCCTGGTGGGACTGTGACCTTGACCTG
 TGCCATCTTGCCAGCCCCCTCCTCAGGTCCACTGGATAAAGGATGGTGCACCCCTGCCCTGGCTCCC
 AGCCCTGTGCTCCTCCTGAGGTGGGACGAGGATGAGGGCACCTATAGCTGCGTGGCCACCACCC
 CTAGCCACGGACCTCAGGAAAGCCCTCCTGTGAGCATCAGGGTCACAGAAACCGGCGATGAGGGGCCAGC
 TGAAGGCTCTGTGGGTGAGTCTGGGCTGGGTACGCTAGCCCTGGCCTTGGGGATCCTGGGAGCCTGGGA
 GTAGTAGCCCTGCTCGTGGGGTATCCTGTGGCGAAAACGACAACCCAGGCGTGAGGAGAGGAAGGCC
 CGGAAAGCCAGGAGGATGAGGAGGAACGTGCAGAGCTGAATCAGTCAGAGGAAGCGGAGATGCCAGAGAA
 TGGTGCCGGGGACCG

ACGCGTACGCGGCCGCTCGAG - GFP Tag - GTTTAA



[View online >](#)

Protein Sequence: >MG206306 representing NM_007425
 Red=Cloning site Green=Tags(s)

MPAGTAARAWVLVLALWGAVAGGQNITARIGEPLVLSCKGAPKKPPQQLWKLNTGRTEAWKVLSPQGGP
 WDSVARILPNGSLLLIPATGIVDEGTFRCRATNRRGKEVKSNYRVRVYQIPGKPEIVDPASELTASVPNKV
 GTCVSEGSYPAGTLSWHLDGKLLIPDGKETLVKEETRRHPETGLFTRSELTVIPTQGGTHPTFSCSFSL
 GLPRRRPLNTAPIQLRVREPGPPEGIQLLVEPEGGIVAPGGTVTLTCAISAQPPPQVHWIKDGAPLPLAP
 SPVLLLPEVGHDEGTYSVATHPSHGPOESPVSIRVTETGDEGPAEGSVGESGLGTLALALGILGGLG
 VVALLVGAILWRKRQPRREERKAPESQEEDEERAEELNQSEEAEMPENGAGGP

TRTRPLE - GFP Tag - V

Restriction Sites: SgfI-MluI

Cloning Scheme:



ACCN: NM_007425

ORF Size: 1206 bp


OTI Disclaimer: Due to the inherent nature of this plasmid, standard methods to replicate additional amounts of DNA in E. coli are highly likely to result in mutations and/or rearrangements. Therefore, OriGene does not guarantee the capability to replicate this plasmid DNA. Additional amounts of DNA can be purchased from OriGene with batch-specific, full-sequence verification at a reduced cost. Please contact our customer care team at custsupport@origene.com or by calling 301.340.3188 option 3 for pricing and delivery.

The molecular sequence of this clone aligns with the gene accession number as a point of reference only. However, individual transcript sequences of the same gene can differ through naturally occurring variations (e.g. polymorphisms), each with its own valid existence. This clone is substantially in agreement with the reference, but a complete review of all prevailing variants is recommended prior to use. [More info](#)

OTI Annotation: This clone was engineered to express the complete ORF with an expression tag. Expression varies depending on the nature of the gene.

Components:	The ORF clone is ion-exchange column purified and shipped in a 2D barcoded Matrix tube containing 10ug of transfection-ready, dried plasmid DNA (reconstitute with 100 ul of water).
Reconstitution Method:	<ol style="list-style-type: none">1. Centrifuge at 5,000xg for 5min.2. Carefully open the tube and add 100ul of sterile water to dissolve the DNA.3. Close the tube and incubate for 10 minutes at room temperature.4. Briefly vortex the tube and then do a quick spin (less than 5000xg) to concentrate the liquid at the bottom.5. Store the suspended plasmid at -20°C. The DNA is stable for at least one year from date of shipping when stored at -20°C.
RefSeq:	<u>NM_007425.3, NP_031451.2</u>
RefSeq Size:	1387 bp
RefSeq ORF:	1209 bp
Locus ID:	11596
UniProt ID:	<u>Q62151</u>
Cytogenetics:	17 B1
Gene Summary:	<p>Mediates interactions of advanced glycosylation end products (AGE). These are nonenzymatically glycosylated proteins which accumulate in vascular tissue in aging and at an accelerated rate in diabetes. Acts as a mediator of both acute and chronic vascular inflammation in conditions such as atherosclerosis and in particular as a complication of diabetes. AGE/RAGE signaling plays an important role in regulating the production/expression of TNF-alpha, oxidative stress, and endothelial dysfunction in type 2 diabetes. Interaction with S100A12 on endothelium, mononuclear phagocytes, and lymphocytes triggers cellular activation, with generation of key proinflammatory mediators. Interaction with S100B after myocardial infarction may play a role in myocyte apoptosis by activating ERK1/2 and p53/TP53 signaling. Can also bind oligonucleotides. Receptor for amyloid beta peptide. Contributes to the translocation of amyloid-beta peptide (ABPP) across the cell membrane from the extracellular to the intracellular space in cortical neurons. ABPP-initiated RAGE signaling, especially stimulation of p38 mitogen-activated protein kinase (MAPK), has the capacity to drive a transport system delivering ABPP as a complex with RAGE to the intraneuronal space. RAGE-dependent signaling in microglia contributes to neuroinflammation, amyloid accumulation, and impaired learning/memory in a mouse model of Alzheimer disease.</p> <p>[UniProtKB/Swiss-Prot Function]</p>

Product images:



Circular map for MG206306