

Product datasheet for **MG206405**

Tacr1 (NM_009313) Mouse Tagged ORF Clone

Product data:

Product Type:	Expression Plasmids
Product Name:	Tacr1 (NM_009313) Mouse Tagged ORF Clone
Tag:	TurboGFP
Symbol:	Tacr1
Synonyms:	NK-; NK1; Nk1r; Spr; Tac; Tac1r
Mammalian Cell Selection:	Neomycin
Vector:	pCMV6-AC-GFP (PS100010)
E. coli Selection:	Ampicillin (100 ug/mL)
Restriction Sites:	SgfI-MluI
Cloning Scheme:	

Cloning sites used for ORF Shuttling:




ACCN:	NM_009313
ORF Size:	1221 bp



[View online »](#)

OTI Disclaimer:	The molecular sequence of this clone aligns with the gene accession number as a point of reference only. However, individual transcript sequences of the same gene can differ through naturally occurring variations (e.g. polymorphisms), each with its own valid existence. This clone is substantially in agreement with the reference, but a complete review of all prevailing variants is recommended prior to use. More info
OTI Annotation:	This clone was engineered to express the complete ORF with an expression tag. Expression varies depending on the nature of the gene.
Components:	The ORF clone is ion-exchange column purified and shipped in a 2D barcoded Matrix tube containing 10ug of transfection-ready, dried plasmid DNA (reconstitute with 100 ul of water).
Reconstitution Method:	<ol style="list-style-type: none">1. Centrifuge at 5,000xg for 5min.2. Carefully open the tube and add 100ul of sterile water to dissolve the DNA.3. Close the tube and incubate for 10 minutes at room temperature.4. Briefly vortex the tube and then do a quick spin (less than 5000xg) to concentrate the liquid at the bottom.5. Store the suspended plasmid at -20°C. The DNA is stable for at least one year from date of shipping when stored at -20°C.
RefSeq:	NM_009313.4 , NP_033339.2
RefSeq Size:	5021 bp
RefSeq ORF:	1224 bp
Locus ID:	21336
UniProt ID:	P30548
Cytogenetics:	6 C3
Gene Summary:	<p>This gene encodes the receptor for the tachykinin, substance P, also referred to as neurokinin 1. This gene belongs to a gene family of tachykinin receptors which are characterized by interactions with G proteins and contain seven hydrophobic transmembrane regions. This receptor has been associated with nitric oxide formation, and it has been localized to cholinergic and nitrenergic neurons as well as on smooth muscle cells. [provided by RefSeq, Mar 2010]</p>

Product images:



Circular map for MG206405