

Product datasheet for **MG206846**

Actl6a (NM_019673) Mouse Tagged ORF Clone

Product data:

Product Type:	Expression Plasmids
Product Name:	Actl6a (NM_019673) Mouse Tagged ORF Clone
Tag:	TurboGFP
Symbol:	Actl6a
Synonyms:	2810432C06Rik; Actl6; AI851094; ARP4; Baf53a; C79802
Mammalian Cell Selection:	Neomycin
Vector:	pCMV6-AC-GFP (PS100010)
E. coli Selection:	Ampicillin (100 ug/mL)
ORF Nucleotide Sequence:	>MG206846 representing NM_019673 Red=Cloning site Blue=ORF Green=Tags(s)

TTTTGTAATACGACTCACTATAGGGCGGCCGGAATTCGTCGACTGGATCCGGTACCGAGGAGATCTGCC
GCC**CGGATCGCC**

ATGAGCGGCGCGTGTACGGCGGAGATGAAGTTGGCGCTCTTGTTTTTGACATTGGATCGTACACAGTGA
GGGCTGGCTATGCTGGCGAGGACTGCCCTAAGGTTGATTTCCCCACGGCTATCGGTGTGGTCTGGAGAG
AGATGACGGAAGTACAATGATGGAGATTGATGGTGACAAAGGCAAGCAGGGCGGCCACCTACTACATA
GACACCAATGCCCTCCGCTGCCAGGGAGAACATGGAGGCCATCTCACCCTCAAGAATGGCATGGTTG
AAGACTGGGATAGTTTCCAGGCCATTTTGGATCATAACACAAGATGCATGTCAAATCCGAAGCCAGCCT
GCATCCTGTTTCATGTCGGAAGCACCGTGAACACCAGGGCGAAGAGAGAGAAAAGTACAGAGTTGATG
TTTGAGCACTACAGCATCCCTGCATTCTTCTTTGCAAAAAGTGCAGTTTTGACGGCATTGCTAATGGTC
GTTCTACTGGGCTGATTTTGGACAGTGGAGCTACCCACACCCTGCGATTCCAGTCCACGATGGCTATGT
TCTTCAACAAGGCATTGTGAAATCCCCTCTGGCTGGAGACTTCATTACCATGCAGTGCAGAGAAGCTTTC
CAGGAAATGAACATAGAAGTCAATCCTCCTTACATGATTGCATCAAAAGAGGCTGTTTCGAGAAGGCTCTC
CAGCCAACTGGAAAAGAAAAGAGAAAAGTGCCTCAGGTTACAAGGCTTGGCACAATTACATGTGCAACTG
CGTCATCCAGGATTTTCAAGCTTCCGTTCTTCAAGTGTGACACTCCACCTACGACGAACAAGTGGCTGCA
CAGATGCCGACCGTTCACTACGAATCCCCAACGGCTACAAGTGGGATTTTGGGGCAGAGCGGCTGAAAA
TTCTGAAGGTTATTTGACCTTCGAACGTAAGGGACTGTCTGGGAACACGATGCTGGGAGTCAGTCA
CGTTGTACAACAGCGTCGGAATGTGTGACATCGACATCAGACCAGGTCTCTACGGCAGTGTGATCGTA
GCAGGAGGAAACAGCTAATACAGAGTTTCACTGACAGGTTAAATAGAGAGCTTTCTCAGAAAACTCCAC
CAAGTATGCGGTTGAAACTGATTGCAAACAACACGACGGTGGAGCGGAGGTTCACTCATGGATTGGTGG
CTCTATCTAGCATCTTTGGGTACCTTTCAACAGATGTGGATTTCTAACAGGAATATGAAGAAGGAGGG
AAGCAGTGTGTAGAAAAGAAAATGCCCT

ACGCGTACGCGGCCGCTCGAG - GFP Tag - GTTTAA



[View online >](#)

Protein Sequence: >MG206846 representing NM_019673
Red=Cloning site Green=Tags(s)

MSGGVYGGDEVGALVFDIGSYTVRAGYAGEDCPKVDFPTAIGVVLERDDGSTMMEIDGDKGKQGGPTYI
 DTNLRVPRENMEAISPLKNGMVEDWDSFQAILDHTYKMHVKSEASLHPVLMSEAPWNTRAKREKLTLM
 FEHYSIPAFFLCKTAVLTAFANGRSTGLILDSGATHHTAIPVHDGYVLQQGIVKSPLAGDFITMQCRELF
 QEMNIELIPPYMIASKEAVREGSPANWKRKEKLPQVTRSWHNYMCNCVIQDFQASVLQVSDSTYDEQVAA
 QMPTVHYEFPNGYNCDFGAERLKIPEGLFDPSNVKGLSGNTMLGVSHVTTSVGMCIDIDIRPGLYGSVIV
 AGGNTLIQSFTDRLNRELSQKTPPSMRCLKLIANNTTVERRFSSWIGGSILASLGTFFQMMWISKQEYEEGG
 KQCVERKCP

TRTRPLE - GFP Tag - V

Restriction Sites:

SgfI-MluI

Cloning Scheme:



ACCN: NM_019673

ORF Size: 1287 bp

OTI Disclaimer: The molecular sequence of this clone aligns with the gene accession number as a point of reference only. However, individual transcript sequences of the same gene can differ through naturally occurring variations (e.g. polymorphisms), each with its own valid existence. This clone is substantially in agreement with the reference, but a complete review of all prevailing variants is recommended prior to use. [More info](#)

OTI Annotation: This clone was engineered to express the complete ORF with an expression tag. Expression varies depending on the nature of the gene.

Components: The ORF clone is ion-exchange column purified and shipped in a 2D barcoded Matrix tube containing 10ug of transfection-ready, dried plasmid DNA (reconstitute with 100 ul of water).

Reconstitution Method:

1. Centrifuge at 5,000xg for 5min.
2. Carefully open the tube and add 100ul of sterile water to dissolve the DNA.
3. Close the tube and incubate for 10 minutes at room temperature.
4. Briefly vortex the tube and then do a quick spin (less than 5000xg) to concentrate the liquid at the bottom.
5. Store the suspended plasmid at -20°C. The DNA is stable for at least one year from date of shipping when stored at -20°C.

RefSeq: [NM_019673.1](#), [NP_062647.1](#)

RefSeq Size: 1770 bp

RefSeq ORF: 1290 bp


Locus ID: 56456

UniProt ID: [Q9Z2N8](#)

Cytogenetics: 3 A3

Gene Summary:

Involved in transcriptional activation and repression of select genes by chromatin remodeling (alteration of DNA-nucleosome topology). Component of SWI/SNF chromatin remodeling complexes that carry out key enzymatic activities, changing chromatin structure by altering DNA-histone contacts within a nucleosome in an ATP-dependent manner. Required for maximal ATPase activity of SMARCA4/BRG1/BAF190A and for association of the SMARCA4/BRG1/BAF190A containing remodeling complex BAF with chromatin/nuclear matrix. Belongs to the neural progenitors-specific chromatin remodeling complex (npBAF complex) and is required for the proliferation of neural progenitors. During neural development a switch from a stem/progenitor to a postmitotic chromatin remodeling mechanism occurs as neurons exit the cell cycle and become committed to their adult state. The transition from proliferating neural stem/progenitor cells to postmitotic neurons requires a switch in subunit composition of the npBAF and nBAF complexes. As neural progenitors exit mitosis and differentiate into neurons, npBAF complexes which contain ACTL6A/BAF53A and PHF10/BAF45A, are exchanged for homologous alternative ACTL6B/BAF53B and DPF1/BAF45B or DPF3/BAF45C subunits in neuron-specific complexes (nBAF). The npBAF complex is essential for the self-renewal/proliferative capacity of the multipotent neural stem cells. The nBAF complex along with CREST plays a role regulating the activity of genes essential for dendrite growth (PubMed:17640523). Component of the NuA4 histone acetyltransferase (HAT) complex which is involved in transcriptional activation of select genes principally by acetylation of nucleosomal histones H4 and H2A. This modification may both alter nucleosome - DNA interactions and promote interaction of the modified histones with other proteins which positively regulate transcription. This complex may be required for the activation of transcriptional programs associated with oncogene and proto-oncogene mediated growth induction, tumor suppressor mediated growth arrest and replicative senescence, apoptosis, and DNA repair. NuA4 may also play a direct role in DNA repair when recruited to sites of DNA damage. Putative core component of the chromatin remodeling INO80 complex which is involved in transcriptional regulation, DNA replication and probably DNA repair (By similarity).[UniProtKB/Swiss-Prot Function]

Product images:

Circular map for MG206846

