

## Product datasheet for **MG207637**

### Igh (BC096617) Mouse Tagged ORF Clone

#### Product data:

Product Type:	Expression Plasmids
Product Name:	Igh (BC096617) Mouse Tagged ORF Clone
Tag:	TurboGFP
Symbol:	Igh
Mammalian Cell Selection:	Neomycin
Vector:	pCMV6-AC-GFP (PS100010)
E. coli Selection:	Ampicillin (100 ug/mL)



[View online »](#)

**ORF Nucleotide Sequence:**

>MG207637 representing BC096617  
 Red=Cloning site Blue=ORF Green=Tags(s)

TTTTGTAATACGACTCACTATAGGGCGGCCGGAATTCGTCGACTGGATCCGGTACCGAGGAGATCTGCC  
 GCC**GCGATCGCC**

ATGGATTTTGGGCTGATTTTTTTTATTGTTGCTCTTTTAAAAGGGGTCCAGTGTGAGGTGAAGCTTCTCG  
 AGTCTGGAGGTGGCTCGGTGCAGCCTGGAGGATCCCTGAAACTCTCCTGTGCAGCCTCAGGATTCGATTT  
 TAGTAAAGACTGGATGAGTTGGGTCCGGCAGGCTCCAGGAAAAGGGCTAGAATGGATTGGAGAAATTAAT  
 CCAGATAGCAGTACGATAAACTATGCACCATCTCTAAAGGATAAAATTCATCATCTCCAGAGAGAACGCCA  
 AAAATACGCTGTACCTGCAAATGAGCAAAGTGAGATCTGAGGACACAGCCCTTTATTACTGTGCAAGATT  
 CCTGGCCTACTGGGGTCAAGGAACCTCAGTCACCGTCTCCTCAGAGCCTGCAAGAGAGCCCACCATCTAC  
 CCACTGACATCCACAAGCTCTGTCAAGTGACCCAGTGATAATCGGCTGCCTGATTCATGATTACTTCC  
 CTTCCGGCAGATGAATGTGACCTGGGAAAGAGTGGGAAGGATATAACCACCGTAAACTCCCACCTGC  
 CCTGGCCTCTGGGGACGGTACACCATGAGCAGCCAGTTGACCCTGCCAGCTGTCGAGTGCCAGAAAGGA  
 GAATCCGTGAAATGTTCCGTGCAACATGACTCTAACCCCGTCCAAGAATTGAACGTGAATTGCCCTGGTA  
 TCTGTTCTCCTCTACTACTCCTCCTCCACCTTCTGCCAGCCCAGCCTGTCACTGCAGCGGCCAGCTCT  
 TGAGGACCTGCTCCTGGGTTTCAGATGCCAGCATCACATGTACTCTGAATGGCCTGAGAGATCCTGAGGGA  
 GCTGTCTTACCTGGGAGCCCTCCACTGGGAAGGATGCAGTGCAGAAGAAAGCTGTGCAGAAATCCTGCG  
 GCTGCTACAGTGTGTCCAGCGTCTGCCTGGCTGTGCTGAGCGCTGGAACAGTGGCGCATCATTCAAGTG  
 CACAGTTACCCATCCTGAGTCTGACACCTTAACTGGCACAAATGCCAAAGTCACAGTGAACACCTTCCCA  
 CCCCAGGTCCACCTGCTACCGCCCGCTCGGAGGAGCTGGCCCTGAATGAGCTCGTGTCCCTGACATGCC  
 TGGTGCAGCTTTCAACCCTAAAGAAGTCTGGTGGCATGGCTGCAATGGAATGAGGAGCTGTCCCAGA  
 AAGCTACCTAGTGTGAGCCCTAAAGGAGCCAGGCGAGGGAGCCACCCTACCTGGTACAAGCGTG  
 TTGCGTGTATCAGCTGAAATCTGAAAACAGGGTGACCACTACTCCTGCATGGTGGCCACGAGGCCCTTGC  
 CCATGAACTTACCCAGAAGACCATCGACCGTCTGTCGGGTAACCCACCAATGTCAGCGTGTCTGTGAT  
 CATGTCAGAGGGAGATGGCATCTGCTAC

**ACGCGTACGCGGCCGCTCGAG** – GFP Tag – GTTTAA

**Protein Sequence:**

>MG207637 representing BC096617  
 Red=Cloning site Green=Tags(s)

MDFGLIFFIVALLKGVQCEVKLLESGLLVQPGGSLKLSAASGFDFSKDWMSWVRQAPGKLEWIGEIN  
 PDSSTINYAPSLKDKFIIISRENAKNTLYLQMSKVRSEDTALYYCARFLAYWQGTSVTVSSEPARPTIY  
 PLTFPQALSSDPVIIGCLIHDFPSGTMNVTWKGSGKDITTVNFPPALASGGRYTMSSQLTLPAVECEP  
 ESVKCSVQHDSNPVQELNVNCPGICSPPTPPPPSCQPSLSLQRPALDLLLLGSDASITCTLNGLRDEP  
 AVFTWEPSTGKDAVQKAVQNSCGCYSVSVLPGCAERWNSGASFKCTVTHPESDTLTGTIAKVTVNTFP  
 PQVHLLPPPSEELALNELVSLTCLVRAFNPKEVLVRWLHGNEELSPESYLVEPLKEPGEATTYLVTSV  
 LRVSAEIKWQGDQYSCMVGHEALPMNFTQKTIDRLSGKPTNVSVSVMSEGDGICY

**TRTRPLE** – GFP Tag – V


**Restriction Sites:**

Sgfl-Mlul

Cloning Scheme:



Plasmid Map:



ACCN: BC096617

ORF Size: 1428 bp

OTI Disclaimer: The molecular sequence of this clone aligns with the gene accession number as a point of reference only. However, individual transcript sequences of the same gene can differ through naturally occurring variations (e.g. polymorphisms), each with its own valid existence. This clone is substantially in agreement with the reference, but a complete review of all prevailing variants is recommended prior to use. [More info](#)

OTI Annotation: This clone was engineered to express the complete ORF with an expression tag. Expression varies depending on the nature of the gene.

Components: The ORF clone is ion-exchange column purified and shipped in a 2D barcoded Matrix tube containing 10ug of transfection-ready, dried plasmid DNA (reconstitute with 100 ul of water).

**Reconstitution Method:**

1. Centrifuge at 5,000xg for 5min.
2. Carefully open the tube and add 100ul of sterile water to dissolve the DNA.
3. Close the tube and incubate for 10 minutes at room temperature.
4. Briefly vortex the tube and then do a quick spin (less than 5000xg) to concentrate the liquid at the bottom.
5. Store the suspended plasmid at -20°C. The DNA is stable for at least one year from date of shipping when stored at -20°C.

**RefSeq:** [BC096617.1](#)

**RefSeq Size:** 1509 bp

**RefSeq ORF:** 1430 bp

**Locus ID:** 111507

**Cytogenetics:** 12 F1-F2 | 12

**Gene Summary:** Summary: Immunoglobulins recognize foreign antigens and initiate immune responses such as phagocytosis and the complement system. Each immunoglobulin molecule consists of two identical heavy chains and two identical light chains. This region represents the germline organization of the heavy chain locus. The locus includes V (variable), D (diversity), J (joining), and C (constant) segments. During B cell development, a recombination event at the DNA level joins a single D segment with a J segment; this partially rearranged D-J gene is then joined to a V segment. The rearranged V-D-J is then transcribed with the IGHM constant region; this transcript encodes a mu heavy chain. Later in development B cells generate V-D-J-Cmu-Cdelta pre-messenger RNA, which is alternatively spliced to encode either a mu or a delta heavy chain. Mature B cells in the lymph nodes undergo switch recombination, so that the V-D-J gene is brought in proximity to one of the IGHG, IGHA, or IGHE genes and each cell expresses either the gamma, alpha, or epsilon heavy chain. Recombination of many different V segments with several J segments provides a wide range of antigen recognition. Additional diversity is attained by junctional diversity, resulting from the random additional of nucleotides by terminal deoxynucleotidyltransferase, and by somatic hypermutation, which occurs during B cell maturation in the spleen and lymph nodes. The RefSeq represents the IGH locus from C57BL/6. Several V and D segments in C57BL/6 are known to be incapable of encoding a protein and are considered pseudogenes. [provided by RefSeq, Jul 2008]