

Product datasheet for **MG209451**

Golga2 (BC011407) Mouse Tagged ORF Clone

Product data:

Product Type:	Expression Plasmids
Product Name:	Golga2 (BC011407) Mouse Tagged ORF Clone
Tag:	TurboGFP
Symbol:	Golga2
Synonyms:	GM130, golgin-95, GM-130, GM 130, mKIAA4150
Mammalian Cell Selection:	Neomycin
Vector:	pCMV6-AC-GFP (PS100010)
E. coli Selection:	Ampicillin (100 ug/mL)



[View online »](#)

ORF Nucleotide Sequence:

>MG209451 representing BC011407
 Red=Cloning site Blue=ORF Green=Tags(s)

TTTTGTAATACGACTCACTATAGGGCGGCCGGAATTCGTCGACTGGATCCGGTACCGAGGAGATCTGCC
 GCC**CGCATCGCC**

ATGGAATCACTGAAGCAGCTCCAGTGGAGAGAGACCAGTATGCGGAGAACCTAAAAGGAGAGAGTGCCA
 TGTGGCAGCAGAGGGTACAGCAAATGGCGGAGCAGGTGCATACACTGAAGGAAGAGAAGGAGCACAGAGA
 ACGGCAGGTACAAGAGCTGGAGACCAGCTTGGCCGCACTCCGGAGTCAGATGGAGGAGCCACCGCCTCCA
 GAGCCTCCAGCTGGGCCCTCCGAGGCGGAAGAGCAGCTGCAGGGAGAGGTCGAGCAGCTGCACAAGGAAC
 TGGAGAGGCTGACGGGACAGCTACGGGCTCAGGTGCAGGACAAATGAGAGCCTGAGCCACCTCAACCGGGA
 GCAGGAGGGCGTCTGCTGGAGCTGGAGCGGGAGGCCAGCGCTGGAGCGAGCAGCCGAGGAGCGCAAG
 CAGATCCTGGAGAGCATGCAGAGTGACCGCACCACCATCAGCAGAGCACTGTCACAGAACCAGAGCTGA
 AGGAGCAGCTGGCCGAGCTGCAGAACGGCTTCGTCAGGCTGACCAATGAGAACATGGAGATCACCAGTGC
 GCTGCAGTCAGAGCAGCATGTCAAGAAAGAGCTAGCCCGAAGCTGGGAGAGCTGCAGGAGAGACTGGGA
 GAGCTGAAGGAGACGGTGGAGCTCAAGAGCCAAGAAGCACAGGGTCTACAGGAGCAGCGAGACCAGTGCC
 TGTCACCTGCAGCAGTACGCTGCTGTTACCAACAGCACTTGGCGGCCTATGAGCAGCTGACCTCTGA
 GAAGGAGGCAATACACAAGCAGCTTCTGCTGCAGACTCAGCTCATGGACCAGCTGCAGCATGAGGAGGTA
 CAGGGCAAGATGGCAGCCGAGCTGGCCCGCAGGAGCTGCAGGAGGCCAGGAGCGCCTAAAAGCTACCA
 GCCAGGAGAACCAACAGCTCCAAGCCAGCTGAGCCTCTTGGTCTCCCTGGGGAAGGAGATGTGGACCA
 GGAGGAAGAAGATGAGGAGGTTCTCAGTCGAGTCTGGCCATCCCAGAAGACCTAGACAGTAGAGAGGCC
 ATGGTGGCATTCTTTAATGCCGCTATAGCCAGAGCAGAGGAAGAACAGGCCCGACTACGTGTGCAGCTGA
 AAGAGCAGAAGGCACGTTGCCGGAGTTTGTCTCACTTGGCAGCCAGTCCAAGCAAGCTTGAAAAGGA
 GGCAGTGGTCCCCAGGAATGTGGATGACTCTGCATCTGAGGAGAGCAATCAGGCCCTACATGTAGCCATG
 GAGAAGCTGCAGAGCCGCTTCTGGAGGTCATGCAGGAGAAAGTGGAGCTCAAGGAGAGGGTAGAGGAGC
 TTGAACATTGCTGCATTAGCTTCTGGAGAGACAGACACCATTGGAGAGTACATTGCCCTTTACAAAAA
 TCAGAGGGCGGTGCTGAAGGCTCGGCATTTGGAGAAGGAAGAATACATCAGCCGGCTGGCTCAGGACAAG
 GAGGAGATGAAGGTGAAGCTGCTGGAGCTCCAGGAGCTGGTGTGAGGCTTGTGAACGAGCGCAATGAAT
 GGCAGGGCAAGTCTCGCCGCTCTCAGAACCTGGTGCAGTGTCCACCCAGTGCCACAGGCTCCCA
 GGAATTTGGTGTCTGACCAGCAGGATGACCTCAGGGAGGTGAGCCTGGCTGACGATATAGAGCCTGCA
 CAAGGAGAGGCAGGGTACCTGCTCCCCATGAGAACCTACTGCACAGCAGATCATGCAGCTGCTGCGTG
 AGATCCAGAACCCCGGAGCGGCCAGGCTGGGTAGCAACCCTTGCATCCCCTTTTCTACCGTGCCGA
 TGAGAACGACGAGGTGAAGATCATGTTGTA

ACGCGTACGCGGCCGCTCGAG – GFP Tag – GTTTAA

Protein Sequence:

>MG209451 representing BC011407
 Red=Cloning site Green=Tags(s)

MESLKQLQVERDQYAENLKGESAMWQQRVQMAEQVHTLKEEKEHRERQVQELETSLAALRSQMEEPPPP
 EPPAGPSEAEELQGEVEQLHKELERLTGQLRAQVQDNESLHNLNREQEGRLELEREAQRWSEAEERK
 QILESMQSDRTTISRALSQNRELKEQLAELQNGFVRLTNENMEITSALQSEQHVKKELARKLGELQERLG
 ELKETVELKSQEAQQLQEQRDQCLSHLQQYAAAYQQHLAAEYQLTSEKEATHKQLLLQTQLMDQLQHEEV
 QGKMAAELARQELQEAQERLKATSQENQQLQAQLSLLVLPGEVDVQEEDEEVPQSSLAIPEDLDSREA
 MVAFFNAAIARAEQARLRVQLKEQKARCRSLSHLAAPVQSKLEKEAVVPRNVDDASSEESNQALHVAM
 EKLQSRFLEVMQEKVELKERVELEHCCIQLSGETDTIGEYIALYQNQRAVLKARHLEKEEYISRLAQDK
 EEMKVKLELQELVLRVNERNEWQKFLAVSQNPGDVLTPVPTGSQFQGAADQQDDLREVSLLADDIEPA
 QGEAGVPAPHENPTAQQIMQLLREIQNPRERPGLGNSNPCIPFFYRADENDEVKIMVV

TRTRPLE – GFP Tag – V

Restriction Sites:

Sgfl-MluI

Cloning Scheme:


ACCN: BC011407

ORF Size: 1853 bp

OTI Disclaimer: The molecular sequence of this clone aligns with the gene accession number as a point of reference only. However, individual transcript sequences of the same gene can differ through naturally occurring variations (e.g. polymorphisms), each with its own valid existence. This clone is substantially in agreement with the reference, but a complete review of all prevailing variants is recommended prior to use. [More info](#)

OTI Annotation: This clone was engineered to express the complete ORF with an expression tag. Expression varies depending on the nature of the gene.

Components: The ORF clone is ion-exchange column purified and shipped in a 2D barcoded Matrix tube containing 10ug of transfection-ready, dried plasmid DNA (reconstitute with 100 ul of water).

Reconstitution Method:

1. Centrifuge at 5,000xg for 5min.
2. Carefully open the tube and add 100ul of sterile water to dissolve the DNA.
3. Close the tube and incubate for 10 minutes at room temperature.
4. Briefly vortex the tube and then do a quick spin (less than 5000xg) to concentrate the liquid at the bottom.
5. Store the suspended plasmid at -20°C. The DNA is stable for at least one year from date of shipping when stored at -20°C.

RefSeq: [BC011407](#), [AAH11407](#)

RefSeq Size: 4174 bp


RefSeq ORF: 1853 bp

Locus ID: 99412

Cytogenetics: 2 B

Gene Summary:

Peripheral membrane component of the cis-Golgi stack that acts as a membrane skeleton that maintains the structure of the Golgi apparatus, and as a vesicle tether that facilitates vesicle fusion to the Golgi membrane (PubMed:28028212). Required for normal protein transport from the endoplasmic reticulum to the Golgi apparatus and the cell membrane (PubMed:28028212). Together with p115/USO1 and STX5, involved in vesicle tethering and fusion at the cis-Golgi membrane to maintain the stacked and inter-connected structure of the Golgi apparatus. Plays a central role in mitotic Golgi disassembly: phosphorylation at Ser-37 by CDK1 at the onset of mitosis inhibits the interaction with p115/USO1, preventing tethering of COPI vesicles and thereby inhibiting transport through the Golgi apparatus during mitosis. Also plays a key role in spindle pole assembly and centrosome organization (By similarity). Promotes the mitotic spindle pole assembly by activating the spindle assembly factor TPX2 to nucleate microtubules around the Golgi and capture them to couple mitotic membranes to the spindle: upon phosphorylation at the onset of mitosis, GOLGA2 interacts with importin-alpha via the nuclear localization signal region, leading to recruit importin-alpha to the Golgi membranes and liberate the spindle assembly factor TPX2 from importin-alpha. TPX2 then activates AURKA kinase and stimulates local microtubule nucleation. Upon filament assembly, nascent microtubules are further captured by GOLGA2, thus linking Golgi membranes to the spindle (By similarity). Regulates the meiotic spindle pole assembly, probably via the same mechanism (PubMed:21552007). Also regulates the centrosome organization (By similarity). Also required for the Golgi ribbon formation and glycosylation of membrane and secretory proteins (By similarity).[UniProtKB/Swiss-Prot Function]

Product images:


Circular map for MG209451