

Product datasheet for **MG209646**

Pip5k1c (BC019138) Mouse Tagged ORF Clone

Product data:

Product Type:	Expression Plasmids
Product Name:	Pip5k1c (BC019138) Mouse Tagged ORF Clone
Tag:	TurboGFP
Symbol:	Pip5k1c
Synonyms:	A1115456; A1835305; Pip5klgamma
Mammalian Cell Selection:	Neomycin
Vector:	pCMV6-AC-GFP (PS100010)
E. coli Selection:	Ampicillin (100 ug/mL)



[View online »](#)

ORF Nucleotide Sequence:

>MG209646 representing BC019138
 Red=Cloning site Blue=ORF Green=Tags(s)

TTTTGTAATACGACTCACTATAGGGCGGCCGGGAATTCGTCGACTGGATCCGGTACCGAGGAGATCTGCC
 GCC**CGCATCGCC**

ATGGAGCTAGAGGTGCCGACGAGGCGGAGAGCGCCGAAGCGGGGCGGTGACGGCGGAAGCGCCCTGGT
 CTGCGGAGAGTGGGGCGGCGCAGGTATGACCCAGAAGAAGGCTGGCCTTGCAGAGGCCCTTGGTGAC
 AGGGCAGCCAGGCCCTGGCCATGGGAAGAAGCTGGGCCACCGAGGCGTGGATGCGTCGGGAGAGACTACG
 TATAAGAAGACCACCTCGTCCACCTGAAGGGCGCCATCCAGCTGGGGATCGGGTACACGGTGGGCAACC
 TGAGCTCTAAGCCGAGCGGGACGTGCTCATGCAGGACTTCTACGTGGTGGAGAGCATCTTCTCCCCAG
 TGAAGGGAGCAACCTCACCCCTGCCACCATTCCAGGATTTCCGCTTCAAGACCTATGCACCTGTTGCC
 TTCGCTACTTCCGGGAGCTCTTTGGCATCCGTCAGATGACTATTTGACTCCCTGTGCAATGAGCCAC
 TCATCGAGCTCTCAACCCCGGGGCCAGCGGCTCTGTCTTCTACGTCACCAGCGACGACGAGTTCATCAT
 CAAGACTGTCATGCACAAGGAGGACAGATTCTGCAGAAGCTGCTGCCTGGCTACTACATGAATCTCAAC
 CAGAACCACGGACGCTGCTGCCAAGTCTATGGGCTGTACTGCGTGCAGTCTGGTGGCAAGAATCC
 GCGTGGTGGTCATGAACAATGTGCTGCCCGTGTGCTCAAATGCACCTTAAGTTCGACCTCAAGGGCTC
 CACGTACAAGCGCAGGGCCAGCAAGAAGGAGAAGGAGAGCCTGCCACCTACAAGGACCTGGACTTC
 ATGCAGGACATGCCCGAGGGGCTGCTGCTGGACTCCGACACCTTTGGCGCCCTGGTCAAGACGCTGCAGC
 GAGACTGCCTGGTGTGGAGAGCTTCAAGATAATGGACTACAGCCTGCTGCTGGGCGTGCACAACATCGA
 TCAGCAGGAGCGAGAGCGCCAGGCCGAGGGCGCCAGAGCAAGGCGGATGAGAAGCGGCCCTGGCCAG
 AAGGCTCTGTATCCACGGCCATGGAGTCTATCCAGGGCGGAGCTGCCCGTGGGGAGGCCATTGAGACAG
 ATGACACGATGGGTGGGATCCAGCAGTGAATGGGCGCGGGGAGCGACTGCTCCACATCGGGATCAT
 TGATATTCTGCAGTCTACAGTTTCATCAAGAAGTTAGAACACACCTGGAAGGCCCTCGTCCATGATGGG
 GACACTGTCTCAGTCCACCGGCCAGTCTTATGCAGAGCGCTTCTTCAAGTTCATGAGCAGCACGGTGT
 TCCGGAAGAGTTCCTCCCTGAAGTCTCTCCATCCAAGAAAGGACGTGGTGCCTGCTGGCGGTCAAACC
 CCTGGGGCCACTGCTGCCTTCTCAGCCAGCCAGATCCCAGCGAGAGAGAGGATGTGCAGTATGACCTG
 CGGGGGCCCGCAGTACCCACGCTTGAGGATGAAGGCCGGCTGACCTCCTGCCCTGCACCCACCGT
 CCTTTGAGGAAGCCACCACGCCTCCATCGCCACCACCCTGTCGTCACCTCCCTCTCCATCCCAGAGCG
 GTCCTTTCAGATACATCGGAGCAGCCCGGTACAGGCGGCGCACGAGTCTTACAGCCAGGATGGCCGG
 CCCAGGAGGAGCCCATGCGGAAGACCTGCAGAAGATAACTGTGCAGGTGGAGCCAGTGTGCGGTGTGG
 GGGTTGTCCCAAGGAGGAGGGTGCAGGAGTGGAGGTCCCCCATGTGGGCATCGGCTGCAGCCTCTGT
 GGAATAGACGCTGCCAGCCAGGCTCAGAGCCTGCCAGCCAGGCTCAGATGAGGAGGATGCACCCCTCT
 ACAGACATCTATTTT

ACGCGTACGCGGCCGCTCGAG - GFP Tag - GTTTAA

Protein Sequence:

>MG209646 representing BC019138
 Red=Cloning site Green=Tags(s)


MELEVPDEAESAEAGAVTAEAAWSAESGAAAGMTQKKAGLAEAPLVTGQPGPGHGKKGHRGVDASGETT
 YKKTSSSTLKGAIQLGIGYTVGNLSSKPERDVLMDYFVVESEIFFPSEGSNLTPAHHFQDFRKYAPVA
 FRYFRELFGIRPDDYLYSLCNEPLIELSNPGASGSVFYVTSDEFI IKTVMHKEAEFLQKLLPGYMNLN
 QNPRTLLPKFYGLYCVQSGGKNIRVVVMNNVLPRVVKMHLKFDLKGSTYKRRASKKEKESLPTYKDLDF
 MQDMPEGLLLDSDTFGALVKTLQRDCLVLESFKIMDYSLLLGVHNIQDQERERQAEGAQSKADEKRPAQ
 KALYSTAMESIQGAARGEAIETDDTMGGIPAVNGRGERLLLHIGIIDILQSYRF IKKLEHTWKALVHDG
 DTVSVHRPSFYAERFFKMSSTVFRKSSSLKSSPSKKGGRGALLAVKPLGPTAAF SASQIPSEREDVQYDL
 RGARSYPTLEDEGRPDLLPCTPPSFEEATTASIATTLSSLSIPERSPSDTSEQPRYRRRTQSSGQDGR
 PQEEPHAEDLQKITVQVEPVCVGVVPEEGAGVEVPPCGASAAAASVEIDAASQASEPASQASDEEDAPS
 TDIYF

TRTRPLE - GFP Tag - V


Restriction Sites: SgfI-MluI

Cloning Scheme:

Cloning sites used for ORF Shuttling:



Plasmid Map:



ACCN: BC019138

ORF Size: 1907 bp

OTI Disclaimer:	The molecular sequence of this clone aligns with the gene accession number as a point of reference only. However, individual transcript sequences of the same gene can differ through naturally occurring variations (e.g. polymorphisms), each with its own valid existence. This clone is substantially in agreement with the reference, but a complete review of all prevailing variants is recommended prior to use. More info
OTI Annotation:	This clone was engineered to express the complete ORF with an expression tag. Expression varies depending on the nature of the gene.
Components:	The ORF clone is ion-exchange column purified and shipped in a 2D barcoded Matrix tube containing 10ug of transfection-ready, dried plasmid DNA (reconstitute with 100 ul of water).
Reconstitution Method:	<ol style="list-style-type: none">1. Centrifuge at 5,000xg for 5min.2. Carefully open the tube and add 100ul of sterile water to dissolve the DNA.3. Close the tube and incubate for 10 minutes at room temperature.4. Briefly vortex the tube and then do a quick spin (less than 5000xg) to concentrate the liquid at the bottom.5. Store the suspended plasmid at -20°C. The DNA is stable for at least one year from date of shipping when stored at -20°C.
RefSeq:	BC019138 , AAH19138
RefSeq Size:	4226 bp
RefSeq ORF:	1907 bp
Locus ID:	18717
Cytogenetics:	10 C1

Gene Summary:

Catalyzes the phosphorylation of phosphatidylinositol 4-phosphate (PtdIns4P) to form phosphatidylinositol 4,5-bisphosphate (PtdIns(4,5)P₂). PtdIns(4,5)P₂ is involved in a variety of cellular processes and is the substrate to form phosphatidylinositol 3,4,5-trisphosphate (PtdIns(3,4,5)P₃), another second messenger. The majority of PtdIns(4,5)P₂ is thought to occur via type I phosphatidylinositol 4-phosphate 5-kinases given the abundance of PtdIns4P. Participates in a variety of cellular processes such as vesicle mediated transport, cell adhesion, cell polarization and cell migration. Together with PIP5K1A is required for phagocytosis, but they regulate different types of actin remodeling at sequential steps. Promotes particle attachment by generating the pool of PtdIns(4,5)P₂ that induces controlled actin depolymerization to facilitate Fc-gamma-R clustering. Mediates RAC1-dependent reorganization of actin filaments. Required for synaptic vesicle transport. Controls the plasma membrane pool of PtdIns(4,5)P₂ implicated in synaptic vesicle endocytosis and exocytosis. Plays a role in endocytosis mediated by clathrin and AP-2 (adaptor protein complex 2). Required for clathrin-coated pits assembly at the synapse. Participates in cell junction assembly. Modulates adherens junctions formation by facilitating CDH1 trafficking. Required for focal adhesion dynamics. Modulates the targeting of talins (TLN1 and TLN2) to the plasma membrane and their efficient assembly into focal adhesions. Regulates the interaction between talins (TLN1 and TLN2) and beta-integrins. Required for uropodium formation and retraction of the cell rear during directed migration. Has a role in growth factor- stimulated directional cell migration and adhesion. Required for talin assembly into nascent adhesions forming at the leading edge toward the direction of the growth factor. Negative regulator of T-cell activation and adhesion. Negatively regulates integrin alpha-L/beta-2 (LFA-1) polarization and adhesion induced by T-cell receptor. Together with PIP5K1A has a role during embryogenesis and together with PIP5K1B may have a role immediately after birth. [UniProtKB/Swiss-Prot Function]