

## Product datasheet for **MG210511**

### **MIh1 (NM\_026810) Mouse Tagged ORF Clone**

#### **Product data:**

Product Type:	Expression Plasmids
Product Name:	MIh1 (NM_026810) Mouse Tagged ORF Clone
Tag:	TurboGFP
Symbol:	MIh1
Synonyms:	1110035C23Rik; AI317206; AI325952; AI561766
Mammalian Cell Selection:	Neomycin
Vector:	pCMV6-AC-GFP (PS100010)
E. coli Selection:	Ampicillin (100 ug/mL)



[View online »](#)

**ORF Nucleotide Sequence:**

>MG210511 representing NM\_026810  
 Red=Cloning site Blue=ORF Green=Tags(s)

TTTTGTAATACGACTCACTATAGGGCGGCCGGAATTCGTCGACTGGATCCGGTACCGAGGAGATCTGCC  
 GCC**CGCATCGCC**

ATGGCGTTTGTAGCAGGAGTTATTTCGGCGTCTGGACGAGACGGTGGTGAACCGCATAGCGCGGGGAAG  
 TCATTCAGCGGCCGCCAATGCTATCAAAGAGATGATAGAAAACGTGTTAGATGCAAAATCTACAAATAT  
 TCAAGTGGTTGTTAAGGAAGGTGGCCTGAAGCTAATTCAGATCCAAGACAATGGCACTGGAAATCAGGAAG  
 GAAGATCTGGATATTGTGTGTGAGAGGTTCACTACGAGTAACTGCAGACTTTTGAGGATTTAGCCAGTA  
 TTTCTACCTATGGCTTTTCGTGGTGGCATTGGCAAGCATAAGCCATGTGGCCCATGTCATTACAAC  
 CAAAACAGCTGATGGAAATGTGCGTACAGAGCAAGTTACTCAGATGGAAAGCTGCAAGCCCTCTCTAAA  
 CCCTGTGCAGGCAACCAGGGCACCTGATCACGGTGAAGACCTTTTTTACAACATAATCACAAGGAGGA  
 AAGCTTTAAAAATCCAAGTGAAGAGTACGAAAAATTTTGAAGTTGTTGGCAGGTATTCAATACACAA  
 TTCAGGCATTAGTATCTCAGTTAAAAACAAGGTGAGACAGTATCTGATGTCAGAACACTGCCAATGCC  
 ACAACCGTGGACAACATTCGCTCCATCTTTGAAATGCGGTTAGTCGAGAAGTATAGAAGTTGGGTGTG  
 AGGATAAAACCTAGCTTTCAAATGAATGGCTATATATCGAATGCAAACTATTCAGTGAAGAAGTGCAT  
 TTTCTACTCTTCATCAACCACCGTCTGGTAGAATCAGCTGCCTTGAGAAAAGCCATTGAAACTGTATAT  
 GCAGCATATTTGCCAAAAACACACACCCATTCTGTACCTCAGTTTGGAAATCAGCCCTCAGAAGCTGG  
 ACGTCAATGTACACCCCAAGCAGCAAGTTCCTTTCTGCATGAGGAGAGCATTCTGCAGCGTGTGCA  
 GCAGCATTGAGAGCAAGCTGCTGGGCTCCAATTCCTCCAGGATGTATTTACCCAGACCTTGCTTCCA  
 GGACTTGCTGGCCCTCTGGGGAGGCAGCTAGACCCACGACAGGGGTGGCTTCTCATCCACTAGTGAA  
 GTGGCGACAAGGTCTACGCTTACCAGATGGTCCGTACGGACTCCCGGGATCAGAAGCTTGACGCTTTCT  
 GCAGCCTGTAAGCAGCCTTGTGCCAGCCAGCCAGGACCTGCCCTGTCCGAGGGGCCAGGACAGAG  
 GGCTCTCTGAAAGGGCCACGCGGGAGGATGAGGAGATGCTTGCTCTCCAGCCCCGCTGAAGCAGCTG  
 CTGAGAGTGAGAACTTGAGAGGGAATCACTAATGGAGACTTCAGACACAGCCAGAAAAGCGGCACCCAC  
 TTCCAGTCCAGGAAGCTCCAGAAAAGAGACATCGGGAGGACTCTGATGTGGAAATGGTGGAAAATGCTTCC  
 GGGAAAGGAAATGACAGCTGCTTGCTACCCAGGAGGAGGATCATTAACTCACCAGCGTCTTGAGTCTCC  
 AGGAAGAGATTAGTGAGCGGTGCCATGAGACTCTCCGGGAGATGCTCCGTAACCATTCCTTTGTGGGCTG  
 TGTGAATCCTCAGTGGGCTTGGCACAGCACCAGACCAAGCTATACCTCCTCAACACTACCAAGCTCAGT  
 GAAGAGCTGTTCTACCAGATACTATTTATGATTTTGCCAACTTTGGTGTCTGAGGTTATCGGAACCAG  
 CGCCACTCTTCGACCTGGCCATGCTGGCCTTAGACAGTCCCTGAAAGTGGCTGGACAGAGGACGACGCCC  
 GAAGGAAGGGCTTGCAGAGTACATTGTCGAGTTTCTGAAGAAGAAAGCGGAGATGCTTGCAGACTATTTT  
 TCTGTGGAGATCGATGAGGAAGGAACTGATCGGATTACCTCTTCTGATTGACAGCTATGTGCCACCTT  
 TGGAGGGACTGCCTATCTTCATTCTTCGACTGGCCACTGAGGTGAATTGGGATGAAGAAAAGGAGTGTTT  
 TGAAAGTCTCAGTAAAGAATGTGCTATGTTTTACTCCATTCGGAAGCAGTATATACTGGAGGAGTCGACC  
 CTCTCAGGCCAGCAGAGTGACATGCCTGGCTCCACGTCAAAGCCCTGGAAGTGGACTGTGGAGCACATTA  
 TCTATAAAGCCTTCGGCTCACACCTCTACCTCCGAAGCATTTCACAGAAGATGGCAATGTCTGACGCT  
 TGCCAACCTGCCAGATCTATACAAGTCTTTGAGCGGTGT

**ACGCGT**ACGCGGCCGCTCGAG – GFP Tag – GTTTAA

Protein Sequence: >MG210511 representing NM\_026810  
 Red=Cloning site Green=Tags(s)

MAFVAGVIRRLDETVVNRIAAGEVIQRPANAIEKEMIEENCLDAKSTNIQVVVKEGGLKLIQIQDNGTGIRK  
 EDLDIVCERFTTSKLTQFEDLASISTYGRGEALASISHVAHVTTITKTADGKCAVRASYSDGKLQAPPK  
 PCAGNQGLITVEDLFYNIITRRKALKNPSEYEGKILEVVGRYSIHNSGISISVKKQGETVSDVRTLPA  
 TTVDNIRSIIFGNAVSRELIEVGCEDKTLAFKMNGYISNANYSVKKCIFLLFINHRLVESAAALRKAJETVY  
 AAYLPKNTHPFLYLSEISPNVDVNVHPTKHEVHFLHEESILQRVQQHIESKLLGSNSSRMFTQTLLP  
 GLAGPSGEAARPTTGVAASSTSGSGDKVYAYQMVRTDSRDQKLD AFLQPVSLLVPSPQDPAPVARGARTE  
 GSPERATREDEEMALPAPAEAAAEESENLERESLMETSDTAQKAAPTSSPGSSRKRHRESDVEMVENAS  
 GKEMTAACYPRRRIINLTSVLSLQEEISERCHETLREMLRNHSFVGCVNPQWALAQHQTKLYLLNNTKLS  
 EELFYQILYDFANFVLRRLSEAPLFDLAMLALDSPESGWTEDDGPKEGLAEYIVEFLKKAEMLADYF  
 SVEIDEEGNLIGLPLIDSYPPEGLPIFILRLATEVNWDEEKECFESLSKECAMFYSIRKQYILEEST  
 LSGQQSDMPGSTSKPWKWTVEHIIYKAFRSHLLPPKHFTEDGNVLQLANLPDLYKVFERC


TRTRPLE - GFP Tag - V

Restriction Sites: SgfI-MluI

Cloning Scheme:

Cloning sites used for ORF Shutting:



**Plasmid Map:**


**ACCN:** NM\_026810

**ORF Size:** 2280 bp

**OTI Disclaimer:** The molecular sequence of this clone aligns with the gene accession number as a point of reference only. However, individual transcript sequences of the same gene can differ through naturally occurring variations (e.g. polymorphisms), each with its own valid existence. This clone is substantially in agreement with the reference, but a complete review of all prevailing variants is recommended prior to use. [More info](#)

**OTI Annotation:** This clone was engineered to express the complete ORF with an expression tag. Expression varies depending on the nature of the gene.

**Components:** The ORF clone is ion-exchange column purified and shipped in a 2D barcoded Matrix tube containing 10ug of transfection-ready, dried plasmid DNA (reconstitute with 100 ul of water).

**Reconstitution Method:**

1. Centrifuge at 5,000xg for 5min.
2. Carefully open the tube and add 100ul of sterile water to dissolve the DNA.
3. Close the tube and incubate for 10 minutes at room temperature.
4. Briefly vortex the tube and then do a quick spin (less than 5000xg) to concentrate the liquid at the bottom.
5. Store the suspended plasmid at -20°C. The DNA is stable for at least one year from date of shipping when stored at -20°C.

**RefSeq:** [NM\\_026810.1](#), [NP\\_081086.1](#)

**RefSeq Size:** 2598 bp

**RefSeq ORF:** 2283 bp

**Locus ID:** 17350

UniProt ID: [Q9JK91](#)

Cytogenetics: 9 60.92 cM

**Gene Summary:** Heterodimerizes with Pms2 to form MutL alpha, a component of the post-replicative DNA mismatch repair system (MMR). DNA repair is initiated by MutS alpha (Msh2-Msh6) or MutS beta (MSH2-MSH3) binding to a dsDNA mismatch, then MutL alpha is recruited to the heteroduplex. Assembly of the MutL-MutS-heteroduplex ternary complex in presence of RFC and PCNA is sufficient to activate endonuclease activity of Pms2. It introduces single-strand breaks near the mismatch and thus generates new entry points for the exonuclease EXO1 to degrade the strand containing the mismatch. DNA methylation would prevent cleavage and therefore assure that only the newly mutated DNA strand is going to be corrected. MutL alpha (Mlh1-Pms2) interacts physically with the clamp loader subunits of DNA polymerase III, suggesting that it may play a role to recruit the DNA polymerase III to the site of the MMR. Also implicated in DNA damage signaling, a process which induces cell cycle arrest and can lead to apoptosis in case of major DNA damages. Heterodimerizes with Mlh3 to form MutL gamma which plays a role in meiosis (By similarity).[UniProtKB/Swiss-Prot Function]