

Product datasheet for **MG210764**

Itch (BC064678) Mouse Tagged ORF Clone

Product data:

Product Type:	Expression Plasmids
Product Name:	Itch (BC064678) Mouse Tagged ORF Clone
Tag:	TurboGFP
Symbol:	Itch
Synonyms:	A130065M08, mKIAA4011
Mammalian Cell Selection:	Neomycin
Vector:	pCMV6-AC-GFP (PS100010)
E. coli Selection:	Ampicillin (100 ug/mL)



[View online »](#)

ORF Nucleotide
Sequence:

>MG210764 representing BC064678
Red=Cloning site Blue=ORF Green=Tags(s)

TTTTGTAATACGACTCACTATAGGGCGGCCGGAATTCGTCGACTGGATCCGGTACCGAGGAGATCTGCC
GCC**CGCATCGCC**

ATGTCTGACAGTGGACCACAGCTTGATTCCATGGGTAGTCTGACCATGAAATCTCAACTTCAGATCACTG
TCATCTCAGCCAACTTAAAGAAAAATAAAGAATTGGTTTGGACCAAGTCTTATGTAGAAGTCACAGT
AGATGGACAGTCAAAGAAGACAGAAAAATGCAATAATACAACAGTCCCAAGTGGAAAGCAGCCCTCACA
GTTATTGTTACCCTACGAGTAAATTATGTTTTCGTGTGTGGAGTACCAGACCTGAAGTCTGATGTTT
TATTGGAACTGCTGGATTAGATATTTATGAAACATTAAGTCAAACAATATGAACTTGAAGAAGTAGT
TATGACTTTGCAGCTTGTAGGTGACAAAGACCAACAGAGACGATGGGAGATTTGCAGTTTGTCTTGAT
GGGCTGCAAGTAGAAGCTGAGGTGTTACTAACGGTGAACGTCATGCTCCGAGAGTACTACACAGAATG
ATGATGGCTGCAGAACAGAGATGATACAAGAGTGAGCACAATGGATCAGAGGACCTGAGGTTGCAGC
GTCAGGGGAAAAACAAGAGGGCCAAATGGGAACAATTCTCCGTCCCTTCAAATGGTGGTTTTAAGCCTTCT
AGACCTCTAGACCTTCGAGACCACCTCCACCCACCCACGAAGACCAGCTTCTGTCAATGGCTCACCAT
CCACGAATTCTGACAGTATGGATCTAGTACAGGCTCCTTGCCACCAACAATAAATGTAATACAAG
TACATCTGAAGGAGCAACATCTGGATTAATAATTCCTTACTATATCTGGAGGCTCGGGCCCTAGGCCT
CTGAATACTGTAAGCCAAGCTCCCTACCACCTGGGTGGGAGCAGAGAGTAGACCAGCATGGGCGTGTTT
ACTATGTAGACCATGTTGAAAAGCGAACAAATGGGATAGACCAGAACCTTACCTCCTGGCTGGGAACG
GCGTGTGGACAATATGGGACGTATTTATATGTTGATCATTTACAAGAACCACAACATGGCAGAGGCCA
ACCTTGGAACTGTCCGGAATGTAACAGTGGCAGCTACAGCGTAGTCAGCTTACAGGAGCAATGCAGC
AGTTTAAACCAGAGATTCATCTATGGGAATCAAGATTTGTTTGTACATCACAACAAGAAATTTGATCC
GCTTGGTCCGTTGCCCTGGATGGGAGAAGAGAAGTACTGATAGCAACGGCAGAGTGTATTTGTCAACCAC
AACACTCGGATTACTCAGTGGGAAGACCCAGAAGCAAGGTGAGTTAAATGAAAAGCCCTTACCAGAAG
GCTGGGAATGAGATTACAGTGGATGGAATTCATATTTTGGGACCACAATAGAAGAGCAACTACTTAA
TATAGATCCACGAACGGGAAAAATCAGCCTTAGACAATGGGCCCCAGATAGCCTATGTGCGGGACTTCAAG
GCAAAAAGTTCAGTATTTCCGTTTCTGGTCCAGCAACTGGCCATGCCACAGCATATAAAGATCACAGTGA
CAAGAAAAACATTGTTTGGAGTTCCTTTCAGCAGATCATGAGCTTACAGCCGCAAGACCTGAGAAGACG
TTTGTGGGTGATTTCCAGGAGAAGAAGTTTAGATTATGGAGGTGTAGCAAGAGAATGGTTCTTTCTT
TTGTCACATGAAGTGTGAACCAATGTATTGCCTGTTTGAATATGCAGGGAAGGATAACTACTGCCTGC
AGATAAACCCCGCTTCTTACATCAATCCAGACCCTGAAATACTTTTCGTTTTATTGGCAGATTTATTGC
CATGGCTCTGTTTCATGGGAAATTCATAGATACCGGATTTTCTTTACCATTCTATAAACGTATCCTGAAT
AAACCAGTTGGACTGAAGGATTTAGAATCTATTGATCCAGAATTTTATAATTCGCTCATCTGGGTTAAAG
AAAACAACATTGAAGAATGTGGTTTGGAAATGACTTCTCAGTTGATAAAGAAATTTAGGTGAAATTAAG
GAGTCATGATTTGAAACCAATGGTGGCAATATCTTGTGACAGAAGAAAAACAAGGAGGAATACATCAGG
ATGGTAGCTGAGTGGAGTTGTCTCGAGGTGTTGAAGAACAGACACAAGCTTTCTTTGAGGGCTTTAATG
AAATCTTCTCAGCAATATTTGCAATATTTTGTGCGAAGGAATTAGAGGTTCTTCTGTGTGGGATGCA
GGAGATTGATTTGAATGACTGGCAGAGACATGCCATCTACCGCCACTACACCAGAACAAGCAAGCAGATC
ATGTGGTTTTGGCAGTTTGTAAAGAAATGATAATGAGAAGAGGATGAGACTTCTGCAGTTTGTACTG
GAACCTGCCGATTGCCAGTAACAACTGCAGTTCCAG

ACGCGTACGCGGCCGCTCGAG - GFP Tag - GTTTAA

Protein Sequence:


>MG210764 representing BC064678
Red=Cloning site Green=Tags(s)

MSDSGPQLDSMGSLTMKSQLQITVISAKLKENKKNWFGSPYVEVTVDGQSKKTEKCNNTNSPKWKQPLT
VIVTPTSCLKFRVWSHQTLKSDVLLGTAGLDIYETLKSNNMKLEEVVMTLQLVGDKEPETMGDLSVCLD
GLQVEAEVVTNGETSCSESTTQNDGCRTRDDTRVSTNGSEDPEVAASGENKRANGNNSPSLSNGGFKPS
RPPRPSRPPPTPRRPASVNGSPSTNSDSDGSSTGSLPPTNTNVNTSTSEGATSGLIIPLTISGGSGPRP
LNTVSQAPLPPGWEQRVDQHGRVYYVDHVEKRRTTWRPEPLPPGWERRVDNMGRYYYYVDHFTRTTTWQRP
TLESVRNYEQWQLQRSQLQGAMQQFNQRFIYGNQDLFATSQNKEFDPLGPLPPGWEKRTDSNGRVYFVNH
NTRITQWEDPRSQQQLNEKPLPEGWEMRFTVDGIPYFVDHNRRATTYIDPRTGKSALDNGPQIAYVRDFK
AKVQYFRFWCQQLAMPQHIIKITVTRKTLFEDSFQQIMSFSPQDLRRRLWVIFPGEEGLDYGGVAREWFFL
LSHEVLNPMYCLFEYAGKDNVCLQINPASYINPDHLKYFRFIGRFIAMALFHGKFIDTGFSLPFYKRILN
KPVGLKDLESIDPEFYNSLIWVKENNIEECGLEMYFSVDKEILGEIKSHDLKPNGGNILVTEENKEEYIR
MVAEWRLSRGVEEQTQAFFEGFNEILPQQYLQYFDAKELEVLLCGMQEIDLNDWQRHAIYRHYTRTSKQI
MWFVQFVKEIDNEKRMRLQLQFVTGTCRLPVTNCRFQ

TRTRPLE - GFP Tag - V

Restriction Sites:

Sgfl-MluI

Plasmid Map:


ACCN: BC064678

ORF Size: 2420 bp

OTI Disclaimer: The molecular sequence of this clone aligns with the gene accession number as a point of reference only. However, individual transcript sequences of the same gene can differ through naturally occurring variations (e.g. polymorphisms), each with its own valid existence. This clone is substantially in agreement with the reference, but a complete review of all prevailing variants is recommended prior to use. [More info](#)

OTI Annotation: This clone was engineered to express the complete ORF with an expression tag. Expression varies depending on the nature of the gene.

Components: The ORF clone is ion-exchange column purified and shipped in a 2D barcoded Matrix tube containing 10ug of transfection-ready, dried plasmid DNA (reconstitute with 100 ul of water).

Reconstitution Method:

1. Centrifuge at 5,000xg for 5min.
2. Carefully open the tube and add 100ul of sterile water to dissolve the DNA.
3. Close the tube and incubate for 10 minutes at room temperature.
4. Briefly vortex the tube and then do a quick spin (less than 5000xg) to concentrate the liquid at the bottom.
5. Store the suspended plasmid at -20°C. The DNA is stable for at least one year from date of shipping when stored at -20°C.

RefSeq: [BC064678](#), [AAH64678](#)

RefSeq Size: 5182 bp

RefSeq ORF: 2420 bp

Locus ID: 16396

Cytogenetics: 2 76.94 cM

Gene Summary:

Acts as an E3 ubiquitin-protein ligase which accepts ubiquitin from an E2 ubiquitin-conjugating enzyme in the form of a thioester and then directly transfers the ubiquitin to targeted substrates (PubMed:15358865, PubMed:16446428, PubMed:17592138, PubMed:18628966, PubMed:20392206, PubMed:25632008). It catalyzes 'Lys-29', 'Lys-48' and 'Lys-63'-linked ubiquitin conjugation (By similarity). Involved in the control of inflammatory signaling pathways (By similarity). Is an essential component of a ubiquitin-editing protein complex, comprising also TNFAIP3, TAX1BP1 and RNF11, that ensures the transient nature of inflammatory signaling pathways (By similarity). Promotes the association of the complex after TNF stimulation (By similarity). Once the complex is formed, TNFAIP3 deubiquitinates 'Lys-63' polyubiquitin chains on RIPK1 and catalyzes the formation of 'Lys-48'-polyubiquitin chains (By similarity). This leads to RIPK1 proteasomal degradation and consequently termination of the TNF- or LPS-mediated activation of NFkB1 (By similarity). Ubiquitinates RIPK2 by 'Lys-63'-linked conjugation and influences NOD2-dependent signal transduction pathways (By similarity). Regulates the transcriptional activity of several transcription factors involved in immune response (PubMed:15358865, PubMed:11828324). Ubiquitinates NFE2 by 'Lys-63' linkages and is implicated in the control of the development of hematopoietic lineages (By similarity). Mediates JUN ubiquitination and degradation (PubMed:15358865). Mediates JUNB ubiquitination and degradation (PubMed:11828324, PubMed:15358865). Critical regulator of type 2 helper T (Th2) cell cytokine production by inducing JUNB ubiquitination and degradation (PubMed:11828324). Involved in the negative regulation of MAVS-dependent cellular antiviral responses (By similarity). Ubiquitinates MAVS through 'Lys-48'-linked conjugation resulting in MAVS proteasomal degradation (By similarity). Following ligand stimulation, regulates sorting of Wnt receptor FZD4 to the degradative endocytic pathway probably by modulating PI42KA activity (By similarity). Ubiquitinates PI4K2A and negatively regulates its catalytic activity (By similarity). Ubiquitinates chemokine receptor CXCR4 and regulates sorting of CXCR4 to the degradative endocytic pathway following ligand stimulation by ubiquitinating endosomal sorting complex required for transport ESCRT-0 components HGS and STAM (By similarity). Targets DTX1 for lysosomal degradation and controls NOTCH1 degradation, in the absence of ligand, through 'Lys-29'-linked polyubiquitination (PubMed:18628966). Ubiquitinates SNX9 (By similarity). Ubiquitinates MAP3K7 through 'Lys-48'-linked conjugation (PubMed:25632008). Involved in the regulation of apoptosis and reactive oxygen species levels through the ubiquitination and proteasomal degradation of TXNIP (By similarity). Mediates the antiapoptotic activity of epidermal growth factor through the ubiquitination and proteasomal degradation of p15 BID (PubMed:20392206). Ubiquitinates BRAT1 and this ubiquitination is enhanced in the presence of NDFIP1 (By similarity).[UniProtKB/Swiss-Prot Function]