

## Product datasheet for **MG210996**

### **Rps6ka5 (NM\_153587) Mouse Tagged ORF Clone**

#### **Product data:**

Product Type:	Expression Plasmids
Product Name:	Rps6ka5 (NM_153587) Mouse Tagged ORF Clone
Tag:	TurboGFP
Symbol:	Rps6ka5
Synonyms:	3110005L17Rik; 6330404E13Rik; AI854034; MSK1; MSPK1; RLPK; RLSK; S6K-alpha-5
Mammalian Cell Selection:	Neomycin
Vector:	pCMV6-AC-GFP (PS100010)
E. coli Selection:	Ampicillin (100 ug/mL)



[View online »](#)

**ORF Nucleotide Sequence:**

>MG210996 representing NM\_153587  
 Red=Cloning site Blue=ORF Green=Tags(s)

TTTTGTAATACGACTCACTATAGGGCGGCCGGGAATTCGTCGACTGGATCCGGTACCGAGGAGATCTGCC  
 GCC**CGGATCGCC**

ATGGAGGGGGAGGGCGGCCGAGCGGGCGCGGGCACCAGCGGGACAGCGGCCAGCGGGCGAGCAGC  
 TCCTTACTGTCAAGCACGAGCTGCGGACCGCCAACTTAACAGGACACGCAGAGAAGGTGGGATTGAAAA  
 CTTTGAGCTGCTGAAGGTCTTAGGAAGTGGAGCTTATGGGAAAGTATTTCTAGTTCTGTAATAAAGCGGC  
 CATGATGCTGAAAGCTGTATGCTATGAAGTTTTAAAAAAGGCAACAATAGTTCAGAAGGCTAAAACTA  
 CAGAGCATACAAGGACAGAACGACAGGTTCTGGAGCACATTAGACAGTCGCCATTTTGGTGACATTGCA  
 CTATGCTTTCCAGACAGAGACCAAGCTTCATCTCATCTTAGATTATAAATGGTGGAGAAGCTTTTACT  
 CATCTTTCTCAAAGAGAGCGTTTACAGAGCACGAAGTGCAGATCTATGTTGGAGAGATTGTGCTTGCCC  
 TGGAACACCTGCACAAGTTGGGGATTATATACCGTGACATTAAGCTTGAAAAATTTCTCCTTGATTCTAA  
 CGCCCATGTGGTCTGACAGATTTCCGCTGAGTAAGGAGTTTGTGGCCGATGAAACTGAAAGAGCATAT  
 TCTTTTGTGGAAGTATTGAATACATGGCTCCAGATATTGTCAGAGGGGGTGATTCGGGACACGACAAGG  
 CAGTTGACTGGTGGAGCTTAGGTGTTCTAATGTATGAGCTACTAACTGGAGCATCTCCTTCACTGTCTGA  
 TGGAGAAAAAATTTCCAGGCTGAGATATCTAGGAGAATCTTAAAAAGTGAAGCTCCGATCCCCAGGAA  
 ATGAGCACAGTAGCTAAAGACCTACTTCAGCGTCTCTTGATGAAAGATCCCAAGAAGCGATTGGGATGTG  
 GCCACGGGATGCAGAAGAAATCAAAGAATCTCTTCTTTGAGAAGATAAAGTGGGATGACTTAGCTGC  
 CAAAAAGTGCCTGCGCCATTCAAGCCGGTATCCGGGATGAGTTAGATGTGAGTAACTTCGCTGAAGAG  
 TTCACAGAGATGGACCCACCTACTCTCCCGCTGCCTTGCCACAGAGCTCCGAGCGACTGTTTCAGGGCT  
 ATTCCTTCGTTGCTCCATCTATTCTTCAAGCGTAACGACGCTGTTATAGATCCTTTCAGTCCATAT  
 GGGAGTTGACCGTCTGGAGTGACAAATGTTGCCAGGAGTGAATGATGAAGGACTCACCCCTCTATCAA  
 CACTATGACCTAGATCTGAAGGACAAACCTTTGGGAGAAGGAAGTTTTTCAATTTGTCGCAAGTGTGTAC  
 ACAAAAAAATAACCAAGCATTGCTGTCAAATAATCAGCAAAAGGATGGAAGCCAATACTCAGAAAGA  
 AATAACAGCTCTGAAACTCTGCGAAGGACACCCCAATATTGTGAAACTACATGAAGTTTTTCACGATCAG  
 GTGGCAGCGTCTGCTCAGCCCCAGGACAGGTTGTTCTATGCTCTTCTACTCTTAGCTCTGCTTTTCA  
 ACAGAAGTCTCACACGGAAGCCTGTCACCTGGACATGGCTGGTCCATAGCACCTCTCAGCTCCCTCCATT  
 GCCACCTCCCATGCCTGAAATTGTCTATTTATCTTCTCTCTGACAATGGACAGCTTCATACGTTTCTT  
 TGATGGAAGTCTGAATGGCGGAGAGCTGTTGAGCGCATCAAGAGAAAGAAGCACTTCAGCGGAGACTG  
 AGGCCAGCTACATCATGCGGAAGCTTGTGTCAGCTGTGAGCCACATGCACGATGTTGGAGTGGTGCACAG  
 AGATCTGAAGCCTGAGAATCTGTTGTTCACTGATGAAAACGACAATTTGGAAATTAAGTGAATGACTTT  
 GGGTTTGACGCTCAAGCCACCTGACAATCAGCCCTCAAGACCCCGTCTTACCCTCCACTACGCGAG  
 CACCGGAGCTCTTGACGCACAATGGCTACGATGAGTCTGTGACCTGTGGAGCTTGGGTGTCATCTGTA  
 TACAATGCTGTGAGGCGAGGTGCCATTCCAGTCTCATGACAGAAGTTAACATGCACCAAGTGCAGTAGAA  
 ATCATGAAGAAAATTAAGGGGAGACTTCTCCTTTGAAGGAGAAGCGTGAAGAATGTATCCCAAGAGG  
 CTAAGATTTGATTCAAGGGCTTCTTACAGTTGATCCAATAAAAGGCTGAAAATGTCTGGTTTGAGATA  
 CAATGAATGGCTTCAGGATGGCAGTCAGCTGTCCTCCAATCCTCTGATGACTCCAGACATCTGGGATCG  
 TCAGGAGCAGCTGTGCACCTGTGTGAAAGCAACCTTTTCATGCCTTTAACAAGTATAAGAGAGAAGGAT  
 TTTGCCTGCAGAATGTGATAAAGCCCCCTTTGGCTAAGAGGAGGAAAAATGAAGAGGACCAGCACCAGCAC  
 AGAGACTCGCAGCAGCTCCAGCGAGAGCTCCCGCTCCTCCTCCTCCAGTCCCACGGCAAGACAACGCT  
 ACCAAGACGCTGCAGCCAGCAATCCCACCGAAGGCAAGTAAACCAGACACCCTCTCCAGTTCTCAGAC

**ACCGTACGCGGCCGCTCGAG** - GFP Tag - GTTTAA

**Protein Sequence:** >MG210996 representing NM\_153587  
Red=Cloning site Green=Tags(s)

MEGEGGSGGAGTSGDSGDGGEQLLTVKHELRTANLTGHAEKVGIENFELLKVLGTGAYGKVFVLRKISG  
HDAGKLYAMKVLKATIVQKAKTTEHTRTERQVLEHIRQSPFLVTLHYAFQTETKLHLILDYINGGELFT  
HLSQRERFTEHEVQIYVGEIVLALHLHKLGIYRDIKLENILLDSNGHVVL TDFGLSKEFVADETERAY  
SFCGTIEYMAPDIVRGGDSGHDKAVDWWSLGVLMYELLTGASPF TVDGEKNSQAEISRRILKSEPPYPQE  
MSTVAKDLLQRLLMKDPKRLGCGPRDAEEIKEHLFFEKIKWDDLAACKVPAPFKPVIRDEL DVSNFAEE  
FTMDPTYSPAALPQSSERLFQGYSFVAPSILFKRNAVIDPLQFHMGVDRPGVTNVARSAMKDSPFYQ  
HYDLDLKDKPLGEGSF SICRKCVHKKTNQAFVKIISKRMEANTQKEITALKLCEGHPNIVKLHEVFHDQ  
VAASAQPPGQVVLCSLLLLALLFNRSLTRKPVTTWLVHSTSQLPPLPPMPEIVLFILLSDNGQLHTFL  
VMELLNGGELFERIKRKKHFSETEASYIMRKLVS AVSHMHDVGVVHRDLKPENLLFTDENDNLEIKVIDF  
GFARLKPPDNQPLKTPCF TLHYAAPELL THNGYDESCDLWSLGVILYTMLSGQVPFQSHDRSLTCTSAVE  
IMKKIKKGD SFEGEAWKNVSQEAKDLIQGLL TVDPNKRLKMSGLRYNEWLQDGSQSSNPLMTPDILGS  
SGAAVHTCVKATFHAFNKYKREGFCLQNVDKAPLAKRRKMKRTSTSTETRSSSESSRSSSSQSHGKTP  
TKTLQPSNPTEGSPDTLFQFSD

TRTRPLE - GFP Tag - V

**Restriction Sites:** Sgfl-MluI



<b>ACCN:</b>	NM_153587
<b>ORF Size:</b>	2589 bp
<b>OTI Disclaimer:</b>	The molecular sequence of this clone aligns with the gene accession number as a point of reference only. However, individual transcript sequences of the same gene can differ through naturally occurring variations (e.g. polymorphisms), each with its own valid existence. This clone is substantially in agreement with the reference, but a complete review of all prevailing variants is recommended prior to use. <a href="#">More info</a>
<b>OTI Annotation:</b>	This clone was engineered to express the complete ORF with an expression tag. Expression varies depending on the nature of the gene.
<b>Components:</b>	The ORF clone is ion-exchange column purified and shipped in a 2D barcoded Matrix tube containing 10ug of transfection-ready, dried plasmid DNA (reconstitute with 100 ul of water).
<b>Reconstitution Method:</b>	<ol style="list-style-type: none"><li>1. Centrifuge at 5,000xg for 5min.</li><li>2. Carefully open the tube and add 100ul of sterile water to dissolve the DNA.</li><li>3. Close the tube and incubate for 10 minutes at room temperature.</li><li>4. Briefly vortex the tube and then do a quick spin (less than 5000xg) to concentrate the liquid at the bottom.</li><li>5. Store the suspended plasmid at -20°C. The DNA is stable for at least one year from date of shipping when stored at -20°C.</li></ol>
<b>RefSeq:</b>	<a href="#">NM_153587.1</a>
<b>RefSeq Size:</b>	4406 bp
<b>RefSeq ORF:</b>	2592 bp
<b>Locus ID:</b>	73086
<b>UniProt ID:</b>	<a href="#">Q8C050</a>
<b>Cytogenetics:</b>	12 E

**Gene Summary:**

Serine/threonine-protein kinase that is required for the mitogen or stress-induced phosphorylation of the transcription factors CREB1 and ATF1 and for the regulation of the transcription factors RELA, STAT3 and ETV1/ER81, and that contributes to gene activation by histone phosphorylation and functions in the regulation of inflammatory genes (By similarity) (PubMed:11553624, PubMed:11909979, PubMed:16806820). Phosphorylates CREB1 and ATF1 in response to mitogenic or stress stimuli such as UV-C irradiation, epidermal growth factor (EGF) and anisomycin (PubMed:11909979). Plays an essential role in the control of RELA transcriptional activity in response to TNF and upon glucocorticoid, associates in the cytoplasm with the glucocorticoid receptor NR3C1 and contributes to RELA inhibition and repression of inflammatory gene expression (PubMed:12628924, PubMed:16806820). In skeletal myoblasts is required for phosphorylation of RELA at 'Ser-276' during oxidative stress (PubMed:12628924). In erythropoietin-stimulated cells, is necessary for the 'Ser-727' phosphorylation of STAT3 and regulation of its transcriptional potential (PubMed:11553624). Phosphorylates ETV1/ER81 at 'Ser-191' and 'Ser-216', and thereby regulates its ability to stimulate transcription, which may be important during development and breast tumor formation (By similarity). Directly represses transcription via phosphorylation of 'Ser-1' of histone H2A (By similarity). Phosphorylates 'Ser-10' of histone H3 in response to mitogenics, stress stimuli and EGF, which results in the transcriptional activation of several immediate early genes, including proto-oncogenes c-fos/FOS and c-jun/JUN (PubMed:15870105, PubMed:16517600). May also phosphorylate 'Ser-28' of histone H3 (PubMed:11441012, PubMed:15870105). Mediates the mitogen- and stress-induced phosphorylation of high mobility group protein 1 (HMGN1/HMG14) (By similarity). In lipopolysaccharide-stimulated primary macrophages, acts downstream of the Toll-like receptor TLR4 to limit the production of pro-inflammatory cytokines (PubMed:18690222). Functions probably by inducing transcription of the MAP kinase phosphatase DUSP1 and the anti-inflammatory cytokine interleukin 10 (IL10), via CREB1 and ATF1 transcription factors (PubMed:18690222). Plays a role in neuronal cell death by mediating the downstream effects of excitotoxic injury (PubMed:12807421). Phosphorylates TRIM7 at 'Ser-106' in response to growth factor signaling via the MEK/ERK pathway, thereby stimulating its ubiquitin ligase activity (By similarity). [UniProtKB/Swiss-Prot Function]