

## Product datasheet for **MG211587**

### Tek (BC050824) Mouse Tagged ORF Clone

#### Product data:

Product Type:	Expression Plasmids
Product Name:	Tek (BC050824) Mouse Tagged ORF Clone
Tag:	TurboGFP
Symbol:	Tek
Synonyms:	AA517024; Cd202b; Hyk; STK1; Tie-2; Tie2
Mammalian Cell Selection:	Neomycin
Vector:	pCMV6-AC-GFP (PS100010)
E. coli Selection:	Ampicillin (100 ug/mL)
ORF Nucleotide Sequence:	>MG211587 representing BC050824 Red=Cloning site Blue=ORF Green=Tags(s)

TTTTGTAATACGACTCACTATAGGGCGCCGGGAATTCGTCGACTGGATCCGGTACCGAGGAGATCTGCC  
GCC**CGATCGCC**

ATGGACTCTTTAGCCGGCTTAGTCTCTGTGGAGTCAGCTTGCTCCTTTATGGAGTAGTAGAAGGCGCCA  
TGGACCTGATCTTGATCAATTCCTACCTCTTGCTGTGATGCCGAAACATCCCTCACCTGCATTGCCTC  
TGGGTGGCACCCCATGAGCCCATCACCATAGGAAGGGACTTTGAAGCCTAATGAACCAGCACCAAGAT  
CCACTGGAGGTTACTCAAGATGTGACCAGAGAATGGGCGAAAAAAGTTGTTTGAAGAGAGAAAAAGGCCA  
GTAAGATTAATGGTGCTTATTTCTGTGAAGGTCGAGTTCGAGGACAGGCTATAAGGATACGGACCATGAA  
GATGCGTCAACAAGCATCCTTCCCTACCTGCTACTTTAACTATGACCGTGGACAGGGGAGATAATGTGAAC  
ATATCTTTCAAAAAGGTGTTAATTAAGAAGAAGATGCAGTGATTTACAAAAATGGATGTGAAGCTCAGA  
AGTGGGGGCCGACTGTAGCCGCTTGTACTACTTGAAGAACAATGGAGTCTGCCATGAAGATACCGG  
GGAATGCATTTGCCCTCCTGGGTTATGGGGAGAACAATGTGAGAAAGCTTGTGAGCCGCACACATTTGGC  
AGGACCTGTAAAGAAAGGTGTAGTGGACCAGAAGGATGCAAGTCTTATGTGTTCTGTCTCCAGACCCTT  
ACGGGTGTTCTGTGCCACAGGCTGGAGGGGTGTCAGTGAATGAAGCATGCCATCTGGTTACTACGG  
ACCAGACTGTAAAGTCAAGTGCAGTGTGAGAAAGAAGGCAGGCCAAGGATGACTCCACAGATAGAGGATT  
TCTCAAGGATGGCAAGGGCTGCAGTGTGAGAAAGAAGGCAGGCCAAGGATGACTCCACAGATAGAGGATT  
TGCCAGATCACATTGAAGTAAACAGTGGAAAATTTAACCCTATCTGCAAAGCCTCTGGGTGGCCACTACC  
TACTAGTGAAGAAATGACCCTAGTGAAGCCAGATGGGACAGTGTCCAAACCAATGACTTCAACTATACA  
GATCGTTTCTCAGTGGCCATATTCAGTGTCAACCGAGTCTTACCTCCTGACTCAGGAGTCTGGGTGCA  
GTGTGAACACAGTGGCTGGGATGGTGGAAAAGCCTTTCAACATTTCCGTCAAAGTTCTTCCAGAGCCCT  
GCACGCCCAATGTGATTGACTGGACATAACTTTGCTATCATCAATATCAGCTCTGAGCCTTACTTT  
GGGGATGGACCCATCAAATCCAAGAAGCTTTTCTATAAACCTGTCAATCAGGCCTGGAAATACATTGAAG  
TGACGAATGAGATTTTCACTCTCAACTACTTGGAGCCCGGACTGACTACGAGCTGTGTGTGACGCTGCC  
CCGTCCTGGAGAGGGTGGAGAAGGCATCTGGGCTGTGAGACGATTTACAACAGCGTCTATCGGACTC



[View online »](#)

CCTCCTCAAGAGGTCTCAGTCTCTGCCCCAAAAGCCAGACAGCTCTAAATTTGACTTGGCAACCGATAT  
 TTCAAACCTCAGAAGATGAATTTTATGTGGAAGTCGAGAGGCGATCCCTGCAAACAAGTGTATCAGCA  
 GAACATCAAAGTGCCTGGGAACCTGACCTCGGTGCTACTGAGCAACTTAGTCCCCAGGGAGCAGTACACA  
 GTCCGAGCTAGAGTCAACACCAAGGCGCAGGGGAGTGGAGTGAAGAAGTCAAGGCTGGACCCTTAGTG  
 ACATTCTCCCTCCTCAACCAGAAAACATCAAGATCTCCAACATCACTGACTCCACAGCTATGGTTTCTTG  
 GACAATAGTGGATGGCTATTCGATTTCTCCATCATATCCGGTATAAGGTTCAAGGCAAAAATGAAGAC  
 CAGCACATTTGATGTGAAGATCAAGAATGCTACCGTTACTCAGTACCAGCTCAAGGCGCTAAGACCGAGA  
 CTACATACCATGTGGATATTTTTGCTGAGAACAACATAGGATCAAGCAACCCAGCCTTTTCTCATGAACT  
 GAGGACGCTTCCACATTCCTCCAGCCTCTGCAGACCTCGGAGGGGAAAAGTGTACTCATAGCCATCCTT  
 GGGTCGGCTGGAATGACTTGCATCACCGTGTGTTGGCGTTTCTGATTATGTTGCAACTGAAGAGAGCAA  
 ATGTCCAAAGGAGAATGGCTCAGGCATTCCAGAACGTGAGAGAAGAACCAGCTGTGCAGTTAACTCAGG  
 AACTCTGGCCCTAACAGGAAGGCCAAAAACAATCCGGATCCCACAATTTATCCTGTGCTGACTGGAAT  
 GACATCAAGTTTCAAGACGTGATCGGAGAGGGCAACTTTGGCCAGTTCTGAAGGCACGCATCAAGAAGG  
 ATGGGTTACGGATGGATGCCCCATCAAGAGGATGAAAGAGTATGCCTCAAAGATGATCACAGGGACTT  
 CGCAGGAGAACTGGAGGTTCTTTGTAACCTTGGACACCATCCAAACATCATTAACTCTTGGGAGCATGT  
 GAACACCGAGGCTATTTGTACCTAGCTATTGAGTATGCCCGCATGGAACCTCCTGGACTTCTGCGTA  
 AGAGCAGAGTGTAGAGACAGACCTGCTTTTGCATCGCCAACAGTACAGCTTCCACACTGTCCTCCCA  
 ACAGCTTCTTATTTGCTGCAGATGTGGCCCGGGGATGGACTACTTGAGCCAGAAACAGTTTATCCAC  
 AGGGACCTGGCTGCCAGAAACATTTTAGTTGGTAAAACACATAGCCAAAATAGCAGATTTTGGATTGT  
 CACGAGGTCAAGAAGTGTATGTAAAAAGACAATGGGAAGGCTCCAGTGCCTGGATGGCAATCGAATC  
 ACTGAACTATAGTGTCTATAACAACAACAGTGTGTCTGGTCTATGGTGTATTGCTCTGGGAGATTGTT  
 AGCTTAGGAGGCCACCCCTACTGCGGCATGACGTGCGCGGAGCTCTATGAGAAGTACCCAGGGCTACA  
 GGCTGGAGAAGCCCTGAACCTGTGATGATGAGGTTGATGATCTAATGAGACAGTGTGGAGGAGACGCC  
 TTATGAGAGACCATCATTGCCCCAGATTTGGTGTCTTAAACAGGATGCTGGAAGAACGGAAGACATAC  
 GTGAACACCACACTGTATGAGAAGTTACCTATGCAGGAATTGACTGCTCTGCGGAAGAAGCAGCC

ACGCGTACGCGGCCGCTCGAG - GFP Tag - GTTTAA

**Protein Sequence:**

>MG211587 representing BC050824

Red=Cloning site Green=Tags(s)

MDSL AGLVLCGVSLLLYGVVEGAMDILINSLPLVSDAETSLTCIASGWHPHEPITIGRDFEALMNQHQD  
 PLEVTDVDTREWAKKVVWKREKASKINGAYFCEGRVVRGQAIRIRTMKMRQASFLPATLTMTVDRGDNVN  
 ISFKKVLIKEEAVIYKNGCEAQKWGPDCSRPCTTCKNNGVCHEDTGEICPPGFMGRTCEKACEPHTFG  
 RTCKERCSGPEGCKSYVFLPDPYGCSCATGWRGLQCNEACPSGYYGPDCKLRCHCTNEEICDRFQGLC  
 SQGWQGLQCEKEGRPRMTPQIEDLPDHIEVNSGKFNPIKASGWPLPTSEEMTLVKPDGTVLQPNDFNYT  
 DRFSVAIFTVNRVLPDPSGVVWCSVNTVAGMVEKPFNISVKVLPPELHAPNVIDTGHNFAIINISSEPYF  
 GDGPIKSKKLFYKPVNQAWKYIEVTNEIFTLNYLEPRTDYELCVQLARPGEGGEGHPGPVRRFTTASIGL  
 PPRGLSLLPKSQTALNLTWQPIFTNSEDEFYVEVERSLQTTSDQQNIKVPGNLTSVLLSNLVPREQYT  
 VRRARVNTKAQGEWSEELRAWLSDILPPQENIKISNITDSTAMVSWTIVDGYSISSIIIRYKVGKQKNE  
 QHIDVKIKNATVTQYQLKLEPETTYHVDIFAENNISSNPAFSELRLTLPSPASADLGGGKMLLIAIL  
 GSAGMTCITVLLAFLIMLQLKRANVQRMAQAFQNVREEPVQFNSGTLALNRKAKNPDPTIYPVLDWN  
 DIKFQDVI GEGNFQVLKARIKKDGLRMDAAIKRMKEYASKDDHRDFAGELEVLCCKLGHHPNIIINLLGAC  
 EHRGYLYLAIEYAPHGNLLDFLRKSRVLETDPAFAIANSTASTLSSQQLLHFAADVARGMDYLSQKQFIH  
 RDLAARNILVGENYIAKIADFGLSRGQEVYVVKTMGRLPVRWMAIESLNYSVYTTNSDVWSYGVLLWEIV  
 SLGGTPYCGMTCAELYEKLPQGYRLEKPLNCDDEVYDLMRQCWREKPYERPSFAQILVSLNRMLEERKTY  
 VNTTLYEKFTYAGIDCSAEAAA

TRTRPLE - GFP Tag - V


**Restriction Sites:**

Sgfl-Mlul

Cloning Scheme:



Plasmid Map:



ACCN: BC050824

ORF Size: 3216 bp

OTI Disclaimer: The molecular sequence of this clone aligns with the gene accession number as a point of reference only. However, individual transcript sequences of the same gene can differ through naturally occurring variations (e.g. polymorphisms), each with its own valid existence. This clone is substantially in agreement with the reference, but a complete review of all prevailing variants is recommended prior to use. [More info](#)

OTI Annotation: This clone was engineered to express the complete ORF with an expression tag. Expression varies depending on the nature of the gene.

<b>Components:</b>	The ORF clone is ion-exchange column purified and shipped in a 2D barcoded Matrix tube containing 10ug of transfection-ready, dried plasmid DNA (reconstitute with 100 ul of water).
<b>Reconstitution Method:</b>	<ol style="list-style-type: none"><li>1. Centrifuge at 5,000xg for 5min.</li><li>2. Carefully open the tube and add 100ul of sterile water to dissolve the DNA.</li><li>3. Close the tube and incubate for 10 minutes at room temperature.</li><li>4. Briefly vortex the tube and then do a quick spin (less than 5000xg) to concentrate the liquid at the bottom.</li><li>5. Store the suspended plasmid at -20°C. The DNA is stable for at least one year from date of shipping when stored at -20°C.</li></ol>
<b>RefSeq:</b>	<u><a href="#">BC050824.1</a></u>
<b>RefSeq Size:</b>	4549 bp
<b>RefSeq ORF:</b>	3218 bp
<b>Locus ID:</b>	21687
<b>Cytogenetics:</b>	4 43.34 cM
<b>Gene Summary:</b>	<p>Tyrosine-protein kinase that acts as cell-surface receptor for ANGPT1, ANGPT2 and ANGPT4 and regulates angiogenesis, endothelial cell survival, proliferation, migration, adhesion and cell spreading, reorganization of the actin cytoskeleton, but also maintenance of vascular quiescence. Has anti-inflammatory effects by preventing the leakage of proinflammatory plasma proteins and leukocytes from blood vessels. Required for normal angiogenesis and heart development during embryogenesis. Required for post-natal hematopoiesis. After birth, activates or inhibits angiogenesis, depending on the context. Inhibits angiogenesis and promotes vascular stability in quiescent vessels, where endothelial cells have tight contacts. In quiescent vessels, ANGPT1 oligomers recruit TEK to cell-cell contacts, forming complexes with TEK molecules from adjoining cells, and this leads to preferential activation of phosphatidylinositol 3-kinase and the AKT1 signaling cascades. In migrating endothelial cells that lack cell-cell adhesions, ANGPT1 recruits TEK to contacts with the extracellular matrix, leading to the formation of focal adhesion complexes, activation of PTK2/FAK and of the downstream kinases MAPK1/ERK2 and MAPK3/ERK1, and ultimately to the stimulation of sprouting angiogenesis. ANGPT1 signaling triggers receptor dimerization and autophosphorylation at specific tyrosine residues that then serve as binding sites for scaffold proteins and effectors. Signaling is modulated by ANGPT2 that has lower affinity for TEK, can promote TEK autophosphorylation in the absence of ANGPT1, but inhibits ANGPT1-mediated signaling by competing for the same binding site. Signaling is also modulated by formation of heterodimers with TIE1, and by proteolytic processing that gives rise to a soluble TEK extracellular domain. The soluble extracellular domain modulates signaling by functioning as decoy receptor for angiopoietins. TEK phosphorylates DOK2, GRB7, GRB14, PIK3R1, SHC1 and TIE1.[UniProtKB/Swiss-Prot Function]</p>