

## Product datasheet for **MG211651**

### Upf1 (NM\_030680) Mouse Tagged ORF Clone

#### Product data:

Product Type:	Expression Plasmids
Product Name:	Upf1 (NM_030680) Mouse Tagged ORF Clone
Tag:	TurboGFP
Symbol:	Upf1
Synonyms:	B430202H16Rik; NORF1; PNORF-1; Rent1; Upflp
Mammalian Cell Selection:	Neomycin
Vector:	pCMV6-AC-GFP (PS100010)
E. coli Selection:	Ampicillin (100 ug/mL)
ORF Nucleotide Sequence:	>MG211651 representing NM_030680 Red=Cloning site Blue=ORF Green=Tags(s)

TTTTGTAATACGACTCACTATAGGGCGGCCGGAATTCGTCGACTGGATCCGGTACCGAGGAGATCTGCC  
GCC**CGATCGCC**

ATGAGTGTGGAGGCGTACGGCCCCAGCTCGAAACACTCACCTTCTTGGACACTGAGGAGGCCGAGCTGC  
TCGGCGCCGACACCCAGGGCTCCGAGTTCGAATTCACCGACTTACCCTTCCCAGCCAGACGCAGACGCC  
CCCCGGCGGCCCGGGCGCGGGAGGCCCGGGCGGAGCGGGCGCAGGCGCGCGGCCGCGCCAGCTCGAC  
GCACAAGTTGGACCAGAGGGCATCTTGCAAAATGGGGCTGTGGATGACAGTGTGGCAAGACCAGCCAGC  
TGCTAGCTGAGCTGAACCTCGAGGAAGATGAAGAGGACACATACTACACTAAGGACTCCCAGTCCACGC  
CTGCAGTACTGTGGAATCCATGATCCTGCCTGCGTGGTTTACTGTAATACCAGCAAGAAGTGGTTCTGC  
AATGGCCGAGGAAATACTTCTGGCAGCCACATTGTGAATCACCTCGTGAGGGCAAAATGCAAGGAAGTGA  
CGCTGCACAAGGACGGGCCTCTGGGCGAGACCGTGTGGAGTGCTACAAGTGTGGCTGCCCAACGTCTT  
CCTGCTGGGCTTCATCCCTGCGAAGGCCGACTCTGTGGTGGTGTGTTGTGCAGGCAGCCCTGTGCCAGC  
CAGAGCAGCCTGAAGGACATCAACTGGGACAGCTCACAGTGGCAGCCCTAATCCAGGACCGGTGCTTTC  
TGTCATGGCTGGTCAAGATTCCTCTGAGCAGGAGCAGCTGCGAGCACGGCAGATCACGGCACAGCAGAT  
CAACAAGCTGGAAGAGCTCTGGAAGGAAAATCCTTCAGCCACTCTGGAGGACCTGGAGAAGCCAGGCGTA  
GACGAGGAGCCACAGCAGTGTCTCTGCGTTACGAGGATGCTTACCAGTACCAGAATCTTCGGGCCAC  
TGGTCAAGCTGGAGGCTGACTATGACAAGAAGTTGAAGGAGTACAGACTCAAGATAACATCACGGTCAG  
GTGGGACCTGGGCCTTAACAAGAAGAGAATCGCCTTCTTCACTTTGCCAAGACTGACTCTGACATGCGG  
CTCATGCAGGGTATGAGATCTGTCTGCGGTACAAAGGGGATCTGGCGCCCTGTGGAAGGGGATTGGCC  
ACGTCATCAAGGTTCTGATAATTATGGTATGAGATTGCTATTGAGCTCCGCAGCAGCGTGGTGCCCC  
TGTGGAAGTGACCCACAACCTTCCAAGTGGATTTGTGTGGAAGTCAACCTCTTTTATAGGATGCAGAGT  
GCACTGAAGACCTTCGCTGTGGACGAGACCTCTGTGTGAGGATATTTACCACAAGCTGCTGGGCCAGC  
AGGTGGAGGATGGTTCATCAAGTGCCAGCTGCCAAAGCGCTTACAGCTCAGGGGCTCCCTGACCTCAA  
CCACTCTCAGGTGATGTGTGAAGACCGTGTGCAGAGACCACTCAGCCTCATCCAGGGCCCTCCAGCC



[View online »](#)

ACAGGCAAGACTGTGACATCAGCCACTATTGTCTACCACCTTGCTCGGCAGGGCAATGGCCTGTACTGG  
 TTTGTGCTCCAAGTAACATCGCTGTGGACCAGCTCACAGAGAAGATCCACCAGACAGGACTGAAGGTCTG  
 ACGCCTCTGTGCCAAGAGCCGTGAGGCCATTGACTCCCCAGTGTCTTCTCGGCTTTGCACAACCAGATC  
 AGGAACATGGACAGCATGCTGAGCTGCAGAAGCTGCAGCAGCTAAAGGATGAGACAGGCCGAGCTGTCAT  
 CTGCAGATGAGAAGCGGTACCGGGCGCTTAAGCGCACAGCTGAGAGAGAACTTTCATGAATGCAGATGT  
 CATATGCTGCACATGTGTGGTGTGTTGACCCGAGGCTGGCCAAGATGCAGTTCCGTTCCATCCCTCATT  
 GACGAGAGCACCCAGGCCACTGAGCCTGAGTGCATGGTGCCTGTAGTCCTTGGGGCCAAGCAGCTAATCC  
 TCGTCCGTGACCACTGCCAGCTGGGCCAGTGGTGTGCAAGAAGGCAGCCAAGGCCGACTGTCAACA  
 ATCGCTCTTCGAGCGCTTGGTGGTGTGGGCATCCGGCCATCCGCTGCAGGTGAATACCGCATGCAC  
 CCTGCACTCAGCGCCTTCCGTCCAACATCTTCTACGAGGGCTCATTGCAGAATGGCGTCACTGCAGCGG  
 ATCGTGTCAAAAAGGCTTTGACTTCCAGTGGCCACAACCTGACAAGCCTATGTTCTTCTACGTGACGCA  
 GGGCCAGGAGGATTGCCAGCTCTGGCACATCTACCTCAACAGGACGGAGGCAGCCAATGTGGAGAAG  
 AATACTACGAAGCTGTTGAAGGCAGGTGCAAAGCCTGACCAGATCGGCATCATCACCCCCTACGAGGGCC  
 AGCGCTTACTTGGTGCAGTACATGCAGTTCAGCGGCTCCCTGCACACAAGCTCTACCAGGAAGTGA  
 AATTGCCAGTGTGGACGCTTCCAGGGCCGGGAGAAGGACTTCATCATTCTGTCTCGTGCAGGCCAAT  
 GAACATCAGGGCATTGGGTTCTAAACGACCCCGGGCTCTGAATGTGGCTCTACCAGAGCAAGATATG  
 GCGTGATCATTGTGGTAACCCAAGGCCCTGTGGAAGCAGCCCTGTGGAATCACCTGTGAGCTACTA  
 CAAGGAACAGAAGGGCTAGTGAAGGGCCGCTCAACAACCTACGTGAGAGCCTCATGCAGTTACAGAAG  
 CCTCGAAACTTGTCAACACTGTCAACCCGGGTGCCCGCTTCAATGACTACTGCCATGTACGATGCCCGTG  
 AGGCCATCATCCCGGCTGTCTATGACCGCAGCAGCCAGGGCCCGCCCTCGAACATGTACTTCCAGAC  
 CCATGACCAGATTAGTATGATCAGCGCAGGCCACGCCAGTGGCTGCCATGAACATCCCTATCCCTTC  
 AACTTGGTCAATGCCTCCCATGCCCCACCTGGCTACTTCGGACAGGCCAACGGCCGCGCAGTGGTCCGG  
 GCACCCCAAAAACCAAGACTGGCCGTGGGGCCGCCAGAAGAACCCTTGGGCTTCTGGGCCCTCGGA  
 GACCACCTTCCCAACAGCCAGGCCAGGAGGTGGCCCTCCAGCCCTTTTTCACAGGTTGCCCTCACA  
 CAGGGTTACGTGTCCATGAGCCAGCCCTCTCAGATGAGCCAGCCTGGCCTCTCCAGCCAGAAGTGTCC  
 AGGACAGTACTCGGTGATGAGTTAAATCACAGATTGACGTGGCACTCTCACAAAGACTCCACATACCA  
 GGGAGAGCGGGCATACCAGCACGGCGGGTCCAGGGCTGTCCAGTAC

ACGCGTACGCGGGCCGCTCGAG - GFP Tag - GTTTAA

**Protein Sequence:**

>MG211651 representing NM\_030680  
 Red=Cloning site Green=Tags(s)

MSVEAYGPSSQTLTFLDTEEAELLGADTQGSEFEFTDFTLPSQTQTPGGPGGAGGPGGAGAGGAAGQLD  
 AQVGP EGILQNGAVDDSAKTSQLLAELNFEDEEDTYTKDLPVHACSYCGIHPACVVCNTSKKWFC  
 NGRGNTSGSHIVNHLVRAKCKEVLHKDGPLGETVLECYNCGCRNVFLGFIPAKADSVVLLCRQPCAS  
 QSSLKIDINWDSSQWQPLIQDRFCSWLVKIPSEQELRQITAQQINKLEELWKENPSATLEDEKPGV  
 DEEPQHVLRLRYEDAYQYQNIIFGPLVKLEADYDKKLKESQTDNIIVRWDLGLNKKRIAFFTLPKTDSMR  
 LMQGDEICLRYKGLAPLWKGIGHVIKVPDNYGDEIAIELRSSVGAPEVTHNFQVDFVWKSTSFDRMQS  
 ALKTFVAVDETSVSGYIYHKLLGHEVEDVVIKCKLQPKRFTAQGLPDLNHSQYVAVKTVLQRPLSLIQGPPG  
 TGKTVTSATIVYHLARQGNPVLVCAPSNIIVDQLTEKIHQTGLKVVRLCAKSREAIIDSPVSLALHNQI  
 RNMDMPELQKLQQLKDETGELSSADEKRYRALKRTERELLMNADVICCTCVGADPRLAKMQFRSIL  
 DESTQATEPECMVPVVLGAKQLILVGDHCQLGPVVMCKKAAGLSQSLFERLVVLRIPRLQVQYRMH  
 PALSAFPSNIFYEGSLQNGVTAADRKKGDFQWPQPKPMFFYVTQGQEEIASSGTSYLNRTAANVEK  
 ITTKLLKAGAKPDQIGIITPYEQRSYLQYMQFSGSLHTKLYQEVEIASVDAFQGREKDFIILSCVRAN  
 EHQIGFLNDPRRLNVALTRARYGVIIIVGNPKALSKQPLWNHLLSYYKEQKALVEGPLNNLRESLMQFSK  
 PRKLVNTVNPGARFMTTAMYDAREAIIPGSVYDRSSQGRPSNMVFQTHDQISMISAGPSHVAAMNIPF  
 NLVMPPMPPPGYFGQANGPAAGRTPKTKTGRGGRQKNRFLGPGPSQTTL PNSQASQDVASQPF SQALT  
 QGYVMSQPSQMSQPLSQPELSQDSYLGDEFKSQIDVALSQDSTYQGERAYQHGGVTGLSQQ

TRTRPLE - GFP Tag - V


**Restriction Sites:**

Sgfl-MluI

Cloning Scheme:



Plasmid Map:



ACCN: NM\_030680

ORF Size: 3339 bp

OTI Disclaimer: The molecular sequence of this clone aligns with the gene accession number as a point of reference only. However, individual transcript sequences of the same gene can differ through naturally occurring variations (e.g. polymorphisms), each with its own valid existence. This clone is substantially in agreement with the reference, but a complete review of all prevailing variants is recommended prior to use. [More info](#)

OTI Annotation: This clone was engineered to express the complete ORF with an expression tag. Expression varies depending on the nature of the gene.

<b>Components:</b>	The ORF clone is ion-exchange column purified and shipped in a 2D barcoded Matrix tube containing 10ug of transfection-ready, dried plasmid DNA (reconstitute with 100 ul of water).
<b>Reconstitution Method:</b>	<ol style="list-style-type: none"><li>1. Centrifuge at 5,000xg for 5min.</li><li>2. Carefully open the tube and add 100ul of sterile water to dissolve the DNA.</li><li>3. Close the tube and incubate for 10 minutes at room temperature.</li><li>4. Briefly vortex the tube and then do a quick spin (less than 5000xg) to concentrate the liquid at the bottom.</li><li>5. Store the suspended plasmid at -20°C. The DNA is stable for at least one year from date of shipping when stored at -20°C.</li></ol>
<b>RefSeq:</b>	<u>NM_030680.3</u>
<b>RefSeq Size:</b>	4598 bp
<b>RefSeq ORF:</b>	3342 bp
<b>Locus ID:</b>	19704
<b>UniProt ID:</b>	<u>Q9EPU0</u>
<b>Cytogenetics:</b>	8 34.15 cM
<b>Gene Summary:</b>	<p>RNA-dependent helicase and ATPase required for nonsense-mediated decay (NMD) of mRNAs containing premature stop codons. Is recruited to mRNAs upon translation termination and undergoes a cycle of phosphorylation and dephosphorylation; its phosphorylation appears to be a key step in NMD. Recruited by release factors to stalled ribosomes together with the SMG1C protein kinase complex to form the transient SURF (SMG1-UPF1-eRF1-eRF3) complex. In EJC-dependent NMD, the SURF complex associates with the exon junction complex (EJC) (located 50-55 or more nucleotides downstream from the termination codon) through UPF2 and allows the formation of an UPF1-UPF2-UPF3 surveillance complex which is believed to activate NMD. Phosphorylated UPF1 is recognized by EST1B/SMG5, SMG6 and SMG7 which are thought to provide a link to the mRNA degradation machinery involving exonucleolytic and endonucleolytic pathways, and to serve as adapters to protein phosphatase 2A (PP2A), thereby triggering UPF1 dephosphorylation. UPF1 can also activate NMD without UPF2 or UPF3, and in the absence of the NMD-enhancing downstream EJC indicative for alternative NMD pathways. Plays a role in replication-dependent histone mRNA degradation at the end of phase S; the function is independent of UPF2. For the recognition of premature termination codons (PTC) and initiation of NMD a competitive interaction between UPF1 and PABPC1 with the ribosome-bound release factors is proposed. The ATPase activity of UPF1 is required for disassembly of mRNPs undergoing NMD (By similarity). Essential for embryonic viability.[UniProtKB/Swiss-Prot Function]</p>