

Product datasheet for **MG216017**

Ddi2 (NM_001017966) Mouse Tagged ORF Clone

Product data:

Product Type:	Expression Plasmids
Tag:	TurboGFP
Symbol:	Ddi2
Synonyms:	1110056G13Rik; 1700027M01Rik; 9130022E05Rik; AI604911; AU040698
Mammalian Cell Selection:	Neomycin
Vector:	pCMV6-AC-GFP (PS100010)
E. coli Selection:	Ampicillin (100 ug/mL)

ORF Nucleotide Sequence: >MG216017 representing NM_001017966
Red=Cloning site Blue=ORF Green=Tags(s)

TTTTGTAATACGACTCACTATAGGGCGCCGGGAATTCGTCGACTGGATCCGGTACCGAGGAGATCTGCC
GCC**GCGATCGCC**

ATGCTGCTCACCGTGTACTGTGTGCGGAGGGACCTCTCCGAGGTGACCTTTCCCTCCAGGTGACGCGCC
ACTTCGAGCTGCACAACCTCCGCGCGCTGTGCGAGCTCGAGTCCGGCATCCCCGCGCCGAGAGCCAGAT
TGTCTATGCAGAAAGCCCTCTCACCACAACACAGGTCACTGGCTTCTTACGGTTTAAAAGATGGTGAT
GTTGTGATTCTACGACAGAAGGAGAATGCAGACCCTCGCCCTGCAGTGCAGTTCTCAAACCTACCCCGGA
TTGACTTCAGTAGCATAGCTGTGCCTGGCACGTCAAACCCCAACAGCCAGCTGCCAGAACCCAGGC
TCAGCACTCATCTCCTGGGAAATGGCATCTTCTCCTCAAGGCCTAGACAACCCAGCCCTGCTCCGAGAC
ATGCTGCTGGCCAACCCGCATGAGTTGCTTGTGCTGAAGGAGCGCAACCCACCTTGGCGGAAGCTCTGC
TCAGTGGAGATCTTGAGAAATCTCTAGGGTCTTGGTGGAGCAGCAGCAGGACCGAGCCGGAGAGAACA
AGAACGGATTTCGTCTGTTTCTGCTGACCCCTTTGATCTCGAAGCTCAGGCAAAGATAGAAGAAGATATA
AGGCAGCAGAACATCGAGGAGAACATGACGATAGCTATGGAGGAGGCCCGGAGAGCTTCGGCCAGGTGCG
CCATGCTGTACATCAACTGCAGAGTGAACGGGCATCCTGTGAAAGCCTTTGTCGACTCAGGAGCCAGAT
GACTATTATGAGCCAGGCCTGTGCAGAAAGGTGTAACATAATGAGACTGGTAGATCGTCGGTGGGCTGGC
ATTGCCAAAGGAGTAGGCACACAGAAGATTATTGGAAGGGTGCATCTAGCTCAGGTTTCAGATTGAAGCGG
ATTTTCTGGCGTGTCTTCTCAATTCTGGAAGAACAGCCTATGGACATGCTTCTGGGTCTGGACATGCT
GAAACGGCACCAAGTGTCTATTGACTTGAAGAAAAATGTTCTGGTGATTGGCACACGGGCTCACAGACC
ACCTTCTTCTGAGGGGGAGTTACCAGAGTGTGCTCGTTTGGCATATGGAACCGGGGAGAGGACATAC
GGCCAGAGGAAATTGCAGACCAAGAATTAGCAGAAGCCATTCAAAAATCAGCCGAGGATGCAGAGCGTCA
GAAGCCA

ACGCGTACGCGGCCGCTCGAG - GFP Tag - GTTTAA




Protein Sequence: >MG216017 representing NM_001017966
 Red=Cloning site Green=Tags(s)

MLLTVYCVRRDLSEVTFSLQVDADFELHNFRLCELESGIPAAESQIVYAERPLTDNHRSLASYGLKDG
 VVILRQKENADPRPAVQFNSLPRIDFSSIAVPGTSSNPQQRQLPRTQAQHSSPGEMASSPQGLDNPALLR
 DMLLANPHEL SLLKERNPLAEALLSGDLEKFSRVLVEQQQDRARREQERIRLFSADPFLEAQAKIEEDI
 RQQNIEENMTIAMEEAPE SFGQVAMLYINCRVNGHPVKAFVDSGAQMTIMSQACAERCNIMRLVDRRWAG
 IAKGVGTQKIIGRVHLAQVQIEGDFLACFSILEEQPMDMLLGLDMLKRHQCSIDLKKNVLVIGTTGSQT
 TFLPEGELPECARLAYGTGREDIRPEEIAQELAEAIQKSAEDAERQKP

TRTRPLE - GFP Tag - V

Restriction Sites: SgfI-MluI

Cloning Scheme:



ACCN: NM_001017966

ORF Size: 1197 bp


OTI Disclaimer: The molecular sequence of this clone aligns with the gene accession number as a point of reference only. However, individual transcript sequences of the same gene can differ through naturally occurring variations (e.g. polymorphisms), each with its own valid existence. This clone is substantially in agreement with the reference, but a complete review of all prevailing variants is recommended prior to use. [More info](#)

OTI Annotation: This clone was engineered to express the complete ORF with an expression tag. Expression varies depending on the nature of the gene.

Components: The ORF clone is ion-exchange column purified and shipped in a 2D barcoded Matrix tube containing 10ug of transfection-ready, dried plasmid DNA (reconstitute with 100 ul of water).

Reconstitution Method:	<ol style="list-style-type: none">1. Centrifuge at 5,000xg for 5min.2. Carefully open the tube and add 100ul of sterile water to dissolve the DNA.3. Close the tube and incubate for 10 minutes at room temperature.4. Briefly vortex the tube and then do a quick spin (less than 5000xg) to concentrate the liquid at the bottom.5. Store the suspended plasmid at -20°C. The DNA is stable for at least one year from date of shipping when stored at -20°C.
Note:	Plasmids are not sterile. For experiments where strict sterility is required, filtration with 0.22um filter is required.
RefSeq:	NM_001017966.2 , NP_001017966.1
RefSeq Size:	3539 bp
RefSeq ORF:	1200 bp
Locus ID:	68817
UniProt ID:	A2ADY9
Cytogenetics:	4 E1
Gene Summary:	<p>Aspartic protease that mediates the cleavage of NFE2L1/NRF1 at 'Leu-104', thereby promoting release of NFE2L1/NRF1 from the endoplasmic reticulum membrane. Ubiquitination of NFE2L1/NRF1 is a prerequisite for cleavage, suggesting that DDI2 specifically recognizes and binds ubiquitinated NFE2L1/NRF1. Seems to act as a proteasomal shuttle which links the proteasome and replication fork proteins like RTF2. Required, with DDI1, for cellular survival following replication stress. Together or redundantly with DDI1, removes RTF2 from stalled forks to allow cell cycle progression after replication stress and maintains genome integrity. [UniProtKB/Swiss-Prot Function]</p>

Product images:



Circular map for MG216017