

Product datasheet for **MG216624**

Pkig (NM_001164055) Mouse Tagged ORF Clone

Product data:

Product Type: Expression Plasmids
Product Name: Pkig (NM_001164055) Mouse Tagged ORF Clone
Tag: TurboGFP
Symbol: Pkig
Synonyms: PKIgamma
Mammalian Cell Selection: Neomycin
Vector: pCMV6-AC-GFP (PS100010)
E. coli Selection: Ampicillin (100 ug/mL)
ORF Nucleotide Sequence: >MG216624 representing NM_001164055
Red=Cloning site Blue=ORF Green=Tags(s)

TTTTGTAATACGACTCACTATAGGGCGGCCGGAATTCGTCGACTGGATCCGGTACCGAGGAGATCTGCC
GCC**CGATCGCC**

ATGATGGAAGTCGAGTCCTCTACTCGGACTTCATCTCCTGCGACCGGACAGGCCGTCGGAATGCAGTCC
CTGACATCCAGGGTGACTCGGAGCCGTGAGTGTGCGGAAGCTCGCTGGAGACATGGCGAGCTCGCACT
TGAGGGAGCAGAAGGACAGGCAGAGGGAAGTACCCCGACAAGGAAGCCAGCAGCCAGCCTGAGAGCAGT
GATCGAACACCTCATCC

ACGCGTACGCGGCCGCTCGAG - GFP Tag - GTTTAA

Protein Sequence: >MG216624 representing NM_001164055
Red=Cloning site Green=Tags(s)
MMEVESSYSDFISCDRTGRRNAVPIQGDSEAVSVRKLAGDMGELALEGAEGQAEGSTPDKEASSQPES
DANTSS

TRTRPLE - GFP Tag - V

Restriction Sites: SgfI-MluI



[View online »](#)

OTI Annotation:	This clone was engineered to express the complete ORF with an expression tag. Expression varies depending on the nature of the gene.
Components:	The ORF clone is ion-exchange column purified and shipped in a 2D barcoded Matrix tube containing 10ug of transfection-ready, dried plasmid DNA (reconstitute with 100 ul of water).
Reconstitution Method:	<ol style="list-style-type: none">1. Centrifuge at 5,000xg for 5min.2. Carefully open the tube and add 100ul of sterile water to dissolve the DNA.3. Close the tube and incubate for 10 minutes at room temperature.4. Briefly vortex the tube and then do a quick spin (less than 5000xg) to concentrate the liquid at the bottom.5. Store the suspended plasmid at -20°C. The DNA is stable for at least one year from date of shipping when stored at -20°C.
RefSeq:	<u>NM_001164055.1</u> , <u>NP_001157527.1</u>
RefSeq Size:	983 bp
RefSeq ORF:	231 bp
Locus ID:	18769
UniProt ID:	<u>O70139</u>
Cytogenetics:	2 84.4 cM
Gene Summary:	Extremely potent competitive inhibitor of cAMP-dependent protein kinase activity, this protein interacts with the catalytic subunit of the enzyme after the cAMP-induced dissociation of its regulatory chains.[UniProtKB/Swiss-Prot Function]